

# Comprehensive Conservation Plan

## *Rainwater Basin Wetland Management District*

**August 2007**

**Prepared by**

*U.S. Fish and Wildlife Service*

Rainwater Basin Wetland Management District  
2610 Avenue Q  
Kearney, NE 68848  
308/236 5015

*and*

Region 6, Mountain-Prairie Region  
Division of Refuge Planning  
134 Union Boulevard, Suite 300  
Lakewood, CO 80228  
303/236 4305

**Approved by**



J. Mitch King  
Regional Director, Region 6  
U.S. Fish and Wildlife Service  
Lakewood, CO

*8/22/07*  
Date




# Comprehensive Conservation Plan

## *Rainwater Basin Wetland Management District*

---

### Submitted by




Gene Mack  
Project Leader  
Rainwater Basin Wetland Management District  
Kearney, NE

8/21/07

Date

### Concurred with by



Rod Krey  
Refuge Supervisor (KS, NE, ND, SD)  
U.S. Fish and Wildlife Service  
Lakewood, CO

8/21/07

Date

*and*



Richard A. Coleman, PhD  
Assistant Regional Director, Region 6  
National Wildlife Refuge System  
U.S. Fish and Wildlife Service  
Lakewood, CO

8/21/07

Date



# Contents

<i>Abbreviations</i> .....	ix
<i>Summary</i> .....	xi
<b>1 Introduction</b> .....	1
1.1 Purpose and Need for the Plan .....	1
1.2 The U.S. Fish and Wildlife Service and the Refuge System .....	4
1.3 National and Regional Mandates .....	5
1.4 District Contributions to National and Regional Plans .....	5
1.5 Ecosystem Description and Threats .....	11
1.6 The Planning Process .....	14
<b>2 The District</b> .....	19
2.1 Establishment, Acquisition, and Management History .....	19
2.2 Special Values of the District .....	19
2.3 Purposes for the District.....	21
2.4 Vision.....	34
2.5 Goals .....	34
2.6 Planning Issues.....	34
<b>3 Affected Environment</b> .....	39
3.1 Physical Environment .....	39
3.2 Biological Resources .....	49
3.3 Cultural Resources .....	54
3.4 Special Management Areas .....	54
3.5 Visitor Services .....	54
3.6 Partnerships.....	55
3.7 Socioeconomic Environment .....	55
3.8 Operations .....	58
<b>4 Management Direction</b> .....	59
4.1 Goals, Objectives, Strategies, and Rationale .....	59
<i>Wetland Goal</i> .....	59
<i>Upland Goal</i> .....	68
<i>Water Rights Goal</i> .....	71
<i>Invasive Plant Species Goal</i> .....	72
<i>Wildlife Diseases Goal</i> .....	72
<i>Research and Science Goal</i> .....	73
<i>Cultural Resources Goal</i> .....	74
<i>Visitor Services Goal</i> .....	75
<i>Partnership Goal</i> .....	77
<i>Socioeconomics Goal</i> .....	78
<i>Operations Goal</i> .....	78
4.2 Staff and Funding.....	79
4.3 Step-down Management Plans.....	79
4.4 Monitoring and Evaluation .....	79
4.5 Plan Amendment and Revision.....	80
<i>Glossary</i> .....	81

*Appendixes*

A. Key Legislation and Policies.....	89
B. Compatibility Determinations.....	95
C. Preparers .....	101
D. Public Involvement .....	103
E. Environmental Compliance .....	113
F. Black-tailed Prairie Dog Management Plan.....	115
G. Section 7 Biological Evaluation .....	131
H. List of Plants .....	135
I. List of Amphibians, Reptiles, and Mammals.....	149
J. List of Birds.....	153
K. Rainwater Basin Vegetation Classification .....	159
L. Fire Management Program .....	173
M. Rainwater Basin Joint Venture Private Lands Program: Best Management Practices Eligibility Criteria and Cost Share.....	177
N. Wetland Management District Ditch and Tile Maintenance Policy.....	185
O. RONS and SAMMS Projects.....	187

<i>Bibliography</i> .....	191
---------------------------	-----

# Figures and Tables

## FIGURES

1	Vicinity map for the Rainwater Basin Wetland Management District, Nebraska .....	2
2	Waterfowl production areas in the Rainwater Basin, Nebraska .....	3
3	Physiographic areas of the United States .....	6
4	“Hourglass” flight path of migratory birds .....	7
5	U.S. Fish and Wildlife Service ecosystem map .....	12
6	Platte-Kansas rivers ecosystem and Arkansas-Red rivers ecosystem map.....	13
7	Steps in the planning process .....	14
8	Waterfowl production areas in Adams County, Nebraska.....	22
9	Waterfowl production areas in Clay County, Nebraska.....	23
10	Waterfowl production areas in Fillmore County, Nebraska.....	24
11	Waterfowl production areas in Franklin County, Nebraska.....	25
12	Waterfowl production areas in Gosper County, Nebraska .....	26
13	Waterfowl production areas in Hall County, Nebraska.....	27
14	Waterfowl production areas in Hamilton County, Nebraska .....	28
15	Waterfowl production areas in Kearney County, Nebraska.....	29
16	Waterfowl production areas in Phelps County, Nebraska .....	30
17	Waterfowl production areas in Saline County, Nebraska .....	31
18	Waterfowl production areas in Seward County, Nebraska.....	32
19	Waterfowl production areas in York County, Nebraska .....	33
20	Land cover in the western portion of the Rainwater Basin, Nebraska.....	41
21	Land cover in the northeastern portion of the Rainwater Basin, Nebraska .....	42
22	Land cover in the southeastern portion of the Rainwater Basin, Nebraska .....	43
23	Groundwater-level changes in Nebraska—predevelopment to spring 2006.....	46
24	Historical wetlands in the Rainwater Basin, Nebraska.....	47
25	Current wetlands in the Rainwater Basin, Nebraska .....	48
26	Populations of Nebraska and the 17 counties in the Rainwater Basin .....	56
27	Age composition within the 17 counties in the Rainwater Basin .....	57
28	Farm count for the 17 counties in the Rainwater Basin .....	57
29	Rainwater Basin duck use-days .....	63
30	Time required for replenishment of endogenous fat reserves following an 8-hour migratory move .....	63

## TABLES

1	Planning process summary .....	15
2	Waterfowl production areas in the Rainwater Basin Wetland Management District.....	20
3	2004 Land cover within the Rainwater Basin.....	45
4	Habitat types at the district in 2004.....	50
5	Nebraska’s state-listed and federally listed species .....	53
6	Current, preferred, and achievable plant composition at WPA wetlands.....	60
7	Current, preferred, and achievable plant composition at WPA uplands .....	69
8	Habitat requirements for selected grassland birds .....	70
9	Key grassland species found in the Rainwater Basin.....	70
10	Current and proposed staff for Rainwater Basin Wetland Management District, Nebraska.....	80
11	Step-down management plans for Rainwater Basin Wetland Management District, Nebraska .....	80



# Abbreviations

<b>Administration Act</b>	National Wildlife Refuge System Administration Act
<b>ATV</b>	all-terrain vehicle
the <b>basin</b>	Rainwater Basin
<b>CAFO</b>	concentrated animal-feeding operation
<b>CCP</b>	comprehensive conservation plan
<b>CFR</b>	Code of Federal Regulations
<b>COMLG</b>	Conservation Order for Mid-continent Light Geese
<b>CP/PLR</b>	Central Plains/Playa Lakes Region
<b>CWCS</b>	comprehensive wildlife conservation strategy
the <b>district</b>	Rainwater Basin Wetland Management District
<b>DUD</b>	duck use-day
<b>EA</b>	environmental assessment
<b>FGDC</b>	Federal Geographic Data Committee
<b>FMP</b>	fire management plan
<b>GIS</b>	Geographic Information System
<b>GS</b>	general pay schedule
<b>Improvement Act</b>	National Wildlife Refuge System Improvement Act of 1997
<b>IPM</b>	integrated pest management
<b>ME</b>	metabolized energy
<b>NASS</b>	Natural Agricultural Statistics Service
<b>NAWCP</b>	North American Waterbird Conservation Plan
<b>NAWMP</b>	North American Waterfowl Management Plan
<b>NDEQ</b>	Nebraska Department of Environmental Quality
<b>NEPA</b>	National Environmental Policy Act
<b>NGPC</b>	Nebraska Game and Parks Commission
<b>NOI</b>	notice of intent
<b>NVCS</b>	National Vegetation Classification System
<b>NWR</b>	national wildlife refuge
<b>ORP</b>	outdoor recreation planner
<b>PILT</b>	payments in lieu of taxes
<b>PL</b>	public law
<b>PM</b>	particulate matter
<b>RWBJV</b>	Rainwater Basin Joint Venture
<b>refuge</b>	national wildlife refuge
<b>Refuge System</b>	National Wildlife Refuge System
<b>RONS</b>	Refuge Operating Needs System
<b>SAMMS</b>	Service Asset Maintenance Management System
<b>Service</b>	U.S. Fish and Wildlife Service
<b>SWG</b>	state wildlife grant
<b>UNL</b>	University of Nebraska–Lincoln
<b>USC</b>	United States Code
<b>USDA</b>	U.S. Department of Agriculture
<b>USDOE</b>	U.S. Department of Energy
<b>USEPA</b>	U.S. Environmental Protection Agency
<b>USFWS</b>	U.S. Fish and Wildlife Service
<b>USGS</b>	U.S. Geological Survey
<b>USGS–BRD</b>	U.S. Geological Survey’s biological resources division
<b>WG</b>	wage grade pay schedule
<b>WMA</b>	wildlife management area
<b>WMD</b>	wetland management district
<b>WPA</b>	waterfowl production area



# Summary

Millions of ducks and geese glide into the flat plains of south-central Nebraska each spring. These migrants take their rest at the few remaining oases that dot the Rainwater Basin, where they find sanctuary and nourishment before continuing their annual migration.

Conserving the integrity of these important stopovers is the role of the Rainwater Basin Wetland Management District. The foundation for management, over the next 15 years, of these public wetlands known as waterfowl production areas is this comprehensive conservation plan approved in August 2007.

## THE DISTRICT

The Rainwater Basin Wetland Management District is a part of the National Wildlife Refuge System of the U.S. Fish and Wildlife Service. The district, which covers 13 Nebraska counties, manages 230 tracts of land (more than 24,000 acres) that form 61 waterfowl production areas. In addition, the district manages 35 conservation easements (2,476 acres).



Each waterfowl production area typically contains wetlands that are managed for waterfowl and shorebirds. Most of the wetlands in the Rainwater Basin are small and surrounded by cropland. Extensive wetland drainage and alteration have reduced the number of wetlands in the basin to a level that threatens waterbird populations. Upland areas are managed for a high diversity of native vegetation to sustain grassland birds.

Nebraska's Rainwater Basin is internationally known for its spectacular bird migrations—329 species of birds have been observed. Common waterbirds include snow goose, northern pintail, green-winged teal, and solitary sandpiper. Grasshopper sparrow, bobolink, and ring-necked pheasant are common grassland species.



*Due to its unique location on the Central Flyway, millions of birds funnel into the district's waterfowl production areas.*

Of the estimated 80,000 visitor days per year at the district, about 60% come for hunting and 40% for wildlife viewing. Hunting seasons include opportunities to harvest waterfowl, pheasant, and deer. Every spring and fall, thousands of tourists and locals visit the district to observe and enjoy the semiannual migration of cranes, waterfowl, and other birds.

## VISION FOR THE DISTRICT

*The Rainwater Basin provides critical habitat for millions of migratory birds.*

*The basin's name reflects both the basis of its wetland hydrology and natural precipitation cycles. A network of functioning wetland and prairie plant ecosystems provides a native grassland mosaic that gives the local community a sense of pride and connection to the Great Plains flora and fauna. The lands managed by the wetland management district serve as an example of land stewardship mimicking natural processes, and they provide an array of wildlife-dependent educational and recreational opportunities.*

*It is only through partnerships with individuals, agencies, and organizations that this vision can be achieved and maintained.*

## **GOALS FOR THE DISTRICT**

The following goals reflect the vision for the district—providing for healthy ecosystems and compatible opportunities for the public to appreciate and enjoy the natural environment.

### ***WETLAND HABITAT GOAL***

Restore, enhance, and maintain the hydrology and early successional vegetation conditions essential to the conservation of migratory birds.

### ***UPLAND HABITAT GOAL***

Reestablish and maintain native grassland communities of the Rainwater Basin.

### ***WATER RIGHTS GOAL***

Develop partnerships to protect the natural hydrology of WPA watersheds and ensure the necessary water rights are in place to protect future use of both ground and surface water.

### ***WILDLIFE DISEASES GOAL***

Work with partners to prevent or control the outbreak and spread of wildlife-borne diseases to protect human and migratory bird populations.

### ***INVASIVE PLANT SPECIES GOAL***

Reduce and control the spread of nondesirable, nonnative plant species within wetland and upland habitats for the benefit of native plant and wildlife communities.

### ***RESEARCH AND SCIENCE GOAL***

Encourage and support research that substantially contributes to the understanding and management of the Rainwater Basin wetland and grassland ecosystem.

### ***CULTURAL RESOURCES GOAL***

Identify and evaluate the cultural resources in the district and protect those that are determined to be significant.

### ***VISITOR SERVICES GOAL***

Provide quality wildlife-dependent recreation and educational opportunities by instilling an understanding of basic ecological processes, purpose of the Rainwater Basin Wetland Management District, and mission of the Service for persons of all abilities and cultural backgrounds.

### ***PARTNERSHIP GOAL***

Promote and develop partnerships with adjacent landowners, public and private organizations, Native American tribes, and other interested individuals to protect, restore, enhance, and maintain a diverse and productive ecosystem.

### ***SOCIOECONOMICS GOAL***

Obtain a better understanding of the social and economic contribution WPAs make to the people and communities within the Rainwater Basin.

### ***OPERATIONS GOAL***

Safely and efficiently use funding, staffing, infrastructure, and partnerships to achieve the purpose and objectives of the Rainwater Basin Wetland Management District.

## **THE COMPREHENSIVE CONSERVATION PLAN**

The Service prepared this final comprehensive conservation plan in cooperation with the Rainwater Basin Joint Venture, the Nebraska Game and Parks Commission, and the Biological Resources Division of the U.S. Geological Survey—along with public participation.

### ***MANAGEMENT DIRECTION***

The management direction of the district will be to manage the district's resources in a holistic manner. There will be a focus on cooperation, coordination, and better exchange of information. An expanded district staff will work with partners to improve the waterfowl production areas across the landscape of the Rainwater Basin. The emphasis will be on adaptive management—as more information is known, management will be changed to improve effects on the environment.

# 1 Introduction



© Mike Vesey

*Northern pintails are abundant at the waterfowl production areas.*

In August 2007, the regional director of region 6 of the U.S. Fish and Wildlife Service (Service) approved this comprehensive conservation plan (CCP) for the Rainwater Basin Wetland Management District (district), Nebraska.

Over the next 15 years, this CCP will serve as the working guide for management programs and use of public lands within the district, which is located in south-central Nebraska (see figure 1, vicinity map).

Chapter 4, “Management Direction,” of this CCP specifies the necessary actions to achieve the vision and purposes of the Rainwater Basin Wetland Management District, which Congress established to manage waterfowl production areas (WPAs) in the Rainwater Basin (basin) (see figure 2, waterfowl production areas). Wildlife and their habitats is the first priority in refuge management and public use (wildlife-dependent recreation) is allowed and encouraged as long as it is compatible with the district’s purposes.

This CCP was developed in compliance with the National Wildlife Refuge System Improvement Act of 1997 (Improvement Act) and Part 602 of “The Fish and Wildlife Service Manual.” The actions described within this CCP meet the requirements of the National Environmental Policy Act of 1969 (NEPA). Appendix A contains more detail on these laws and policies. Public involvement and the planning process are described in “1.6 The Planning Process.”

## 1.1 PURPOSE AND NEED FOR THE PLAN

The purpose of the planning process that led to development of this CCP is to identify the role that the district will play in support of the mission of the National Wildlife Refuge System (Refuge System), and to provide long-term guidance for management of district programs and activities.

The CCP is needed

- to communicate with the public and other partners in efforts to carry out the mission of the Refuge System;
- to provide a clear statement of direction for management of the district’s WPAs;
- to provide neighbors, visitors, and government officials with an understanding of the Service’s management actions on and around the district’s WPAs;
- to ensure that the Service’s management actions are consistent with the mandates of the Improvement Act;
- to ensure that the management of the district’s WPAs is consistent with federal, state, and county plans;
- to provide a basis for the development of budget requests for the district’s operation, maintenance, and capital improvement needs.

Sustaining the nation’s fish and wildlife resources is a task that can be accomplished only through the




Figure 1. Vicinity map for the Rainwater Basin Wetland Management District, Nebraska.




Figure 2. Waterfowl production areas in the Rainwater Basin, Nebraska.

combined efforts of governments, businesses, and private citizens.

## 1.2 THE U.S. FISH AND WILDLIFE SERVICE AND THE REFUGE SYSTEM

The Service is the principal federal agency responsible for fish, wildlife, and plant conservation. The Refuge System is one of the Service's major programs.

### U.S. FISH AND WILDLIFE SERVICE

*The mission of the U.S. Fish and Wildlife Service, working with others, is to conserve, protect, and enhance fish and wildlife and their habitats for the continuing benefit of the American people.*

Over a century ago, America's fish and wildlife resources were declining at an alarming rate. Concerned citizens, scientists, and hunting and angling groups joined together to restore and sustain America's national wildlife heritage. This was the genesis of the U.S. Fish and Wildlife Service.

Today, the Service enforces federal wildlife laws, manages migratory bird populations, restores nationally significant fisheries, conserves and restores vital wildlife habitat, protects and recovers endangered species, and helps other governments with conservation efforts. In addition, the Service administers a federal aid program that distributes hundreds of millions of dollars to states for fish and wildlife restoration, boating access, hunter education, and related programs across America.

The Service manages the National Wildlife Refuge System including thousands of WPAs and other special management areas. It also operates 66 national fish hatcheries and 78 ecological services field stations.

### Service Activities in Nebraska

Service activities in Nebraska contribute to the state's economy, ecosystems, and education programs. The Rainwater Basin Wetland Management District contributes to the economic benefits of hunting, wildlife observation, and photography in Nebraska. A report titled, "Banking on Nature 2004: The Economic Benefits to Local Communities of National Wildlife Refuge Visitation," evaluated the effects of refuges on local economies.

Based on figures from 2004, the district is estimated to have generated \$900,000 in local economic effects from recreation visits (BBC Research and Consulting 2006). The majority of effects were associated with expenditures by nonresident visitors. In addition, the

district's budget contributes a stimulus to the local economy with a significant portion of payroll, maintenance, and operation expenditures spent locally.

The district employs 12 full-time employees, has a current budget of \$1.8 million, and has an annual visitation of 80,000. The budget includes funds for the fire program and the Partners for Fish and Wildlife Program. In addition, volunteers contribute 240 hours to the district's operations.

The Nebraska Sport Fish and Wildlife Restoration Program is a source of federal excise taxes paid by hunters, anglers, and boaters on fishing and hunting equipment. The monies generated from this tax have economic benefits to Nebraska. In 2001, the economic impact of angler expenditures was \$146 million and hunters contributed \$198 million to the overall economy (U.S. Fish and Wildlife Service [USFWS] 2000).

### THE NATIONAL WILDLIFE REFUGE SYSTEM

In 1903, President Theodore Roosevelt designated the 5.5-acre Pelican Island in Florida as the nation's first wildlife refuge for the protection of brown pelicans and other native, nesting birds. This was the first time the federal government set aside land for wildlife. This small but significant designation was the beginning of the Refuge System.

One hundred years later, the Refuge System has become the largest collection of lands in the world specifically managed for wildlife, encompassing over 96 million acres within 546 refuges and over 3,000 small areas for waterfowl breeding and nesting. Today, there is at least one refuge in every state including Puerto Rico and the U.S. Virgin Islands.

In 1997, the Improvement Act established a clear mission for the Refuge System.

*The mission of the National Wildlife Refuge System is to administer a national network of lands and waters for the conservation, management, and where appropriate, restoration of the fish, wildlife and plant resources and their habitats within the United States for the benefit of present and future generations of Americans.*

The Improvement Act states that each national wildlife refuge (that is, each unit of the Refuge System, which includes wetland management districts) shall be managed

- to fulfill the mission of the Refuge System;
- to fulfill the individual purposes of each refuge and district;

- to consider the needs of fish and wildlife first;
- to fulfill the requirement of developing a CCP for each unit of the Refuge System, and fully involve the public in the preparation of these plans;
- to maintain the biological integrity, diversity, and environmental health of the Refuge System;
- to recognize that wildlife-dependent recreation activities including hunting, fishing, wildlife observation, photography, environmental education, and interpretation are legitimate and priority public uses;
- to retain the authority of refuge managers to determine compatible public uses.

In addition to the mission for the Refuge System, the wildlife and habitat vision for each unit of the Refuge System stresses the following principles:

- Wildlife comes first.
- Ecosystems, biodiversity, and wilderness are vital concepts in refuge and district management.
- Habitats must be healthy.
- Growth of refuges and districts must be strategic.
- The Refuge System serves as a model for habitat management with broad participation from others.

Following passage of the Improvement Act, the Service immediately began to carry out the direction of the new legislation, including preparation of CCPs for all national wildlife refuges and wetland management districts. Consistent with the Improvement Act, the Service prepares all CCPs in conjunction with public involvement. Each unit of the Refuge System is required to complete its CCP within the 15-year schedule (by 2012).

### People and the Refuge System

The nation's fish and wildlife heritage contributes to the quality of American lives and is an integral part of the country's greatness. Wildlife and wild places have always given people special opportunities to have fun, relax, and appreciate the natural world.

Whether through bird watching, fishing, hunting, photography, or other wildlife pursuits, wildlife recreation contributes millions of dollars to local economies. In 2002, approximately 35.5 million people visited the Refuge System, mostly to observe wildlife in their natural habitats. Visitors are most often accommodated through nature trails, auto tours, interpretive programs, and hunting and fishing opportunities. Significant economic benefits are being generated to the local communities that surround refuges and wetland management districts. Economists report that Refuge System visitors contribute more than \$792 million annually to local economies.

## 1.3 NATIONAL AND REGIONAL MANDATES

Refuge System units are managed to achieve the mission and goals of the Refuge System, along with the designated purpose of the refuges and districts (as described in establishing legislation, executive orders, or other establishing documents). Key concepts and guidance of the Refuge System are in the Refuge System Administration Act of 1966 (Administration Act), Title 50 of the Code of Federal Regulations (CFRs), "The Fish and Wildlife Service Manual," and the Improvement Act.

The Improvement Act amends the Administration Act by providing a unifying mission for the Refuge System, a new process for determining compatible public uses on refuges and districts, and a requirement that each refuge and district be managed under a CCP. The Service has made compatibility determinations (see appendix B) for the following uses at the district: haying, grazing, farming, environmental education, interpretation, wildlife observation, photography, recreational fishing, recreational hunting, and timber harvest.

The Improvement Act states that wildlife conservation is the priority of Refuge System lands and that the Secretary of the Interior will ensure that the biological integrity, diversity, and environmental health of refuge lands are maintained. Each refuge and district must be managed to fulfill the Refuge System's mission and the specific purposes for which it was established. The Improvement Act requires the Service to monitor the status and trends of fish, wildlife, and plants in each refuge and district.

A detailed description of these and other laws and executive orders that may affect the CCP or the Service's implementation of the CCP is in appendix A. Service policies on planning and day-to-day management of refuges and districts are in the "Refuge System Manual" and "The Fish and Wildlife Service Manual."

## 1.4 DISTRICT CONTRIBUTIONS TO NATIONAL AND REGIONAL PLANS

The Rainwater Basin Wetland Management District contributes to the conservation efforts described here.

### *FULLFILLING THE PROMISE*

A 1999 report, "Fulfilling the Promise, The National Wildlife Refuge System" (USFWS 1999), is the culmination of a yearlong process by teams of Service employees to evaluate the Refuge System nationwide. This report was the focus of the first national Refuge System conference (in 1998)—attended by refuge managers, other Service employees, and representatives from leading conservation organizations.

The report contains 42 recommendations packaged with three vision statements dealing with wildlife and habitat, people, and leadership. This CCP deals with all three of these major topics. The planning team looked to the recommendations in the document for guidance during CCP planning.

***PARTNERS IN FLIGHT***

The “Partners in Flight” program began in 1990 with the recognition of declining population levels of many migratory bird species. The challenge, according to the program, is managing human population growth while maintaining functional natural ecosystems. To meet this challenge, Partners in Flight worked to identify priority land-bird species and habitat types. Partners in Flight activity has resulted in 52 bird conservation plans covering the continental United States.

The primary goal of Partners in Flight is to provide for the long-term health of the bird life of this continent. The first priority is to prevent the rarest species from going extinct. The second priority is to prevent uncommon species from descending into threatened status. The third priority is to “keep common birds common.”


There are 58 physiographic areas, defined by similar physical geographic features, wholly or partially

contained within the contiguous United States and several others wholly or partially in Alaska. The Rainwater Basin Wetland Management District lies within physiographic area 34 (see figure 3, physiographic areas).

The source of the following description is from the Partners in Flight website (Butcher, no date).

Physiographic area 34, known as the “Central Mixed-grass Prairie,” includes the central portion of Nebraska and Kansas, and a small portion occurs in southern South Dakota. The Nebraska Sandhills cover the northern and western portions of the area. The sandhills are an area of rolling, irregular dunes interspersed with gently sloping valleys and numerous small wetlands. The remainder of the physiographic area is a dissected loess plain, drained by several major rivers. All of the uplands are natural mixed- and tall-grass prairie communities, and the larger river valleys support northern floodplain forests.

The Nebraska Sandhills is one of the few, large, productive areas for grassland birds on the continent. It remains in excellent condition due to long-term use of virtually 100% of private lands for grazing livestock. Historical grazing practices have been, largely, beneficial. To keep the area healthy for birds, it is important to maintain the health of the ranching economy.



**Figure 3. Physiographic areas of the United States.**  
(Source: Partners in Flight.)

Priority bird species and habitats of the Central Mixed-grass Prairie are listed below:

*Grassland*

lesser prairie-chicken  
greater prairie-chicken  
Swainson's hawk  
dickcissel  
long-billed curlew  
Bell's vireo  
Smith's longspur

*Big River Sandbars*

pipin plover

*Wetlands*

American white pelican  
black rail

Large wetland-grassland complexes benefit all of the high-priority birds and are essential to some. It is important to maintain all existing complexes. The black rail is a species that uses wet meadows; its ecology remains largely unknown and more survey work and retention of potential habitat are needed.

One of the most important features of the physiographic area is the close proximity of the Platte River to the district's wetlands, which combine to form a large and diverse habitat complex. This complex provides midlatitudinal, migrational habitat for midcontinental populations of sandhill cranes (86%), snow geese (90%), white-fronted geese (90%), and mallards (50%). About 30% of the continental population of northern pintails migrate through the district's habitats. In addition, impressive numbers of shorebirds annually stop in the area.

Key areas are receiving attention through the RWBJV and other endeavors. The efforts—some of which involve repeated removal of woody vegetation from sandbars that have stabilized with altered hydrology—are important to continue to keep the area attractive for these birds.

Maintenance of large, unfragmented, grassland ecosystems is the conservation objective for areas such as the Missouri Coteau where agriculture is not dominant. On the drift prairie and other agricultural areas, it is important to conserve discrete blocks of grassland-wetland complexes.

## ***NORTH AMERICAN WATERFOWL MANAGEMENT PLAN***

The Rainwater Basin is located in the Central Flyway, which is one of four administrative waterfowl flyways in North America.

Due to its unique location in the Central Flyway, millions of birds—including sandhill cranes, Canada geese, snow geese, and mallards—funnel into the district's WPAs to rest and eat before continuing on their journey (see figure 4, "hourglass" flight path of migratory birds).



**Figure 4. "Hourglass" flight path of migratory birds.**

The Central Flyway occurs in the following states and provinces:

- United States—Colorado, Kansas, Montana, Nebraska, New Mexico, North Dakota, Oklahoma, South Dakota, Texas, and Wyoming
- Mexico—Aguascalientes, Chihuahua, Coahuila, Colima, Distrito Federal, Durango, Guanajuato, Guerrero, Jalisco, Nayarit, Nuevo León, México, Morelos, Oaxaca, Puebla, Queretaro, San Luis Potosi, Sinaloa, Sonora, Tamaulipas, Veracruz, and Zacatecas
- Canada—Alberta and Saskatchewan

Federal, state, and provincial representatives from the United States and Canada make up the Central Flyway Council. The council meets regularly to coordinate population surveys, regulate and set hunting seasons, and plan for management of the migratory bird resource.

Canada and the United States united in 1986 to form the North American Waterfowl Management Plan (NAWMP) (USFWS et al. 1998), designed to restore diminishing continental waterfowl populations to the levels of the 1970s. The NAWMP envisioned a 15-year effort to achieve landscape conditions that could sustain waterfowl populations. Specific NAWMP objectives are to increase and restore duck populations to the average levels of the 1970s: 62 million breeding ducks and a fall flight of 100 million birds.

By 1985, waterfowl populations had plummeted to record lows. Habitat that waterfowl depend on was disappearing at a rate of 60 acres per hour. Recognizing the importance of waterfowl and wetlands to North

Americans and the need for international cooperation to help in the recovery of a shared resource, the United States and Canada governments developed a strategy to restore waterfowl populations through habitat protection, restoration, and enhancement. Mexico became a signatory to the plan in 1994.

The plan is innovative because of its international scope, plus its implementation at the regional level. Its success depends on the strength of partnerships called “joint ventures,” involving federal, state, provincial, tribal, and local governments; businesses; conservation organizations; and individual citizens.

Joint ventures are regional, self-directed partnerships that carry out science-based conservation through a wide array of community participation in the United States, Canada, and Mexico. Joint ventures develop implementation plans focusing on areas of concern identified in the plan. The Rainwater Basin Wetland Management District lies within the administrative boundary of the Rainwater Basin Joint Venture.

### **Rainwater Basin Joint Venture**

The Rainwater Basin Joint Venture (Gersib et al. 1992) is one of 14 joint ventures formed to undertake conservation projects. The joint venture was founded in 1992 with a goal to restore and permanently protect 37,000 acres of high-quality wetlands and 25,000 acres of associated uplands with adequate water and distribution to meet the needs of waterfowl and other migratory birds.

#### **Location**

Although the RWBJV’s focus is the basin, its boundary also encompasses that portion of “Bird Conservation Region 19” in Nebraska (see figure 1). Three prominent geographic features occur within the joint venture—the basin in south-central Nebraska, the Nebraska Sandhills in north-central Nebraska, and the central portion of the Platte River.

The Playa Lakes Joint Venture bounds the RWBJV on the west and south. On the east, the RWBJV borders the Upper Mississippi–Great Lakes Region Joint Venture. The RWBJV’s northern boundary is the Nebraska state line and it borders the Prairie Pothole Region and the Northern Great Plains Region joint ventures.

#### **Description**

Land use in the basin portion of the RWBJV is almost entirely agriculture, with corn and soybeans being the dominant crops. The topography of the basin is flat and it is poorly drained—forming thousands of shallow wetlands. Most of the wetlands are small and incorporated into cropland. Extensive wetland drainage and alteration has reduced the number of wetlands to a level that threatens populations of waterfowl and other waterbirds. The area is part of the tall- and mixed-grass prairie region of the Great Plains.

The sandhills portion is native, mixed-grass prairie that is used for livestock production. The topography is hilly, grass-covered, sand dunes. The porous sand allows for rapid percolation, forming a large groundwater reservoir. The groundwater is exposed in the low valleys and depressions—creating over a million acres of lakes, wetlands, and wet meadows.

The Platte River is a flat, braided river that has become forested in the last century. It is historically significant for settlement and for wildlife migrations. An approximately 150-mile stretch of the river transects the Central Flyway. Each spring nearly one-half million sandhill cranes and millions of ducks and geese use the river. River use by spring-migrating waterfowl increases dramatically when the basin’s wetlands are dry or frozen.

#### **Conservation**

Each joint venture includes the participation of individuals, corporations, conservation organizations, and government agencies (USFWS et al. 1998). The district contributes to and participates in the RWBJV through its Partners for Fish and Wildlife Program, participation on various committees, and management of WPAs.

### ***UNITED STATES SHOREBIRD CONSERVATION PLAN***

Partners from state and federal agencies and nongovernmental organizations from across the country pooled their resources and expertise to develop a conservation strategy for migratory shorebirds and the habitats upon which they depend. The “United States Shorebird Conservation Plan” provides a scientific framework to determine species, sites, and habitats that most urgently need conservation action.

Main goals of the plan, completed in 2000, are to ensure that adequate quantity and quality of shorebird habitat is maintained at the local level and to maintain or restore shorebird populations at the continental and hemispheric levels. Separate technical reports were developed for a conservation assessment, research needs, a comprehensive monitoring strategy, and education and outreach. These national assessments were used to “step down” goals and objectives into 11 regional conservation plans.

The Rainwater Basin Wetland Management District lies within the “Central Plains/Playa Lakes Region” (CP/PLR), which includes Texas (excluding the coast); eastern New Mexico and Colorado; western Oklahoma, Kansas, and Nebraska; and the southeastern corner of Wyoming. Within the CP/PLR, there are five bird conservation regions as part of the North American Bird Conservation Initiative. Shorebird habitat types within the CP/PLR include ephemeral wetlands such as playa lakes, semipermanent wetlands, seasonally flooded wetlands, mud and alkali flats, wet meadows, short-grass prairie, agricultural fields, reservoirs, rivers, and other shallow water sources such as



ditches and farm ponds. All shorebird habitats within the CP/PLR are important to their conservation, especially scattered and ephemeral wetlands. The most critical habitats in the CP/PLR for those species experiencing population declines are grasslands and other upland areas with low vegetation structure, rivers, and salt flats. The salt flats and river sandbars within the CP/PLR are used by migrating and nesting piping plovers and nesting snowy plovers. Short-stature grasslands, mowed areas, alfalfa fields, plowed agricultural fields, and sod farms in Oklahoma, Kansas, and Nebraska are important to migrating buff-breasted sandpipers. Short-grass prairies, grazed grasslands, and fallow agricultural fields in western Kansas and Nebraska and in the eastern portions of New Mexico, Colorado, and Wyoming are breeding sites for mountain plovers. Upland habitats, recently burned rangelands, and salt flats within the CP/PLR are used as staging areas for American golden-plovers.

A particular challenge is that a high proportion of potentially important playa lakes, salt plains, and grasslands are either wholly or partly under private ownership (approximately 75%). The CP/PLR working group supports the hemispheric and national goals of the “United States Shorebird Conservation Plan,” to “restore and maintain populations of all species of shorebirds in the Western Hemisphere” and to “stabilize populations of all shorebird species known or suspected of being in decline due to limiting factors occurring within the U.S., while ensuring that common species remain common” (Brown et al. 2000). However, land managers and biologists within the CP/PLR cannot control some of the factors that affect population sizes such as conditions on the wintering and, in most species, on the breeding grounds. Therefore, the CP/PLR can best contribute to the above goals by concentrating on protecting, improving, and expanding the habitat available for migrating shorebirds within the region and protecting, improving, and expanding the habitat available for the species that breed in the region.

Within the Rainwater Basin, shorebird habitat can be exceptional if rainfall coincides with shorebird migration. The basin is important to large numbers of migratory Hudsonian godwits; long-billed dowitchers; buff-breasted, white-rumped, stilt, least, and pectoral sandpipers; American avocets; lesser yellowlegs; semipalmated plovers; common snipe; Wilson’s phalaropes; and killdeer. However, shorebird numbers and stopover habitat may not be fully understood. With the cooperation of landowners, state and federal natural resource agencies are conducting wetland habitat restoration and improvement work on private lands as part of the RWBJV.

## ***NORTH AMERICAN WATERBIRD CONSERVATION PLAN***

The “North American Waterbird Conservation Plan” (NAWCP) is the product of an independent partnership of individuals and institutions having interest and responsibility for conservation of waterbirds and their habitats in the Americas. This partnership—Waterbird Conservation for the Americas—was created to support a vision in which the distribution, diversity, and abundance of populations and habitats of breeding, migratory, and nonbreeding waterbirds are sustained or restored throughout the lands and waters of North America, Central America, and the Caribbean. The NAWCP provides a continental-scale framework for the conservation and management of 210 species of waterbirds including seabirds, coastal waterbirds, wading birds, and marsh birds. These birds use aquatic habitats in 29 nations throughout North America, Central America, the islands and pelagic (open ocean) waters of the Caribbean Sea and western Atlantic, the United States–associated Pacific Islands, and pelagic waters of the Pacific. The waterbirds’ dependence on aquatic habitats—such as wooded swamps, stream corridors, salt marshes, barrier islands, continental shelf waters, and pelagic waters—make them especially vulnerable to the threats that face water and wetland resources globally.

The conservation of waterbirds faces significant challenges. Eighty percent of the species considered in the NAWCP are colonial nesters—congregating at breeding sites in numbers ranging up to hundreds of thousands of birds. Of this group, the plan finds that one-third are considered to be at risk of serious population loss. Waterbird populations are subject to numerous threats, many of which are habitat-based and affect all aquatic birds and other aquatic resources. The threats that the NAWCP identifies as requiring remedial action include destruction of inland and coastal wetlands, introduced predators and invasive species, pollutants, mortality from fisheries and other human industries, disturbance, and conflicts arising from abundant species.

The NAWCP identifies strategies and opportunities for achieving its vision. The plan documents a

dynamic process for species status assessment for use in setting conservation priorities at a regional scale and has identified many of the key issues that require conservation action. In addition, the plan promotes habitat- and site-based conservation actions throughout the Americas, especially via the “Important Bird Areas” programs and similar efforts. Regional waterbird conservation working groups will “step down” the continental-level aspects of the plan to the regional and local levels. At all scales, the NAWCP advocates integration of waterbird conservation with other bird conservation initiatives, when appropriate, to efficiently provide the best management options for local wildlife and habitat managers.

### **RECOVERY PLANS FOR FEDERALLY LISTED THREATENED OR ENDANGERED SPECIES**

Where federally listed threatened or endangered species occur at the Rainwater Basin Wetland Management District’s WPAs, management goals and strategies in their respective recovery plans will be followed. The list of threatened or endangered species that occur at the district will change as species are listed or delisted, or as listed species are discovered on district lands.

The district lies within the historical range of the whooping crane, least tern (interior population), American burying beetle, and western prairie fringed orchid. All of these species have recovery plans. If these species are found in the district, the staff will follow recovery plan guidelines.

### **STATE COMPREHENSIVE CONSERVATION WILDLIFE STRATEGY**

Over the past several decades, documented declines of wildlife populations have occurred nationwide. Congress created the State Wildlife Grant (SWG) program in 2001. This program provides states and territories with federal dollars to support conservation aimed at preventing wildlife from becoming endangered and in need of protection under the Endangered Species Act. The SWG program represents an ambitious endeavor to take an active hand in keeping species from becoming threatened or endangered in the future.

According to the SWG program, each state, territory, and the District of Columbia must have a completed comprehensive wildlife conservation strategy (CWCS) by October 1, 2005 to receive future funding.

These strategies will help define an integrated approach to the stewardship of all wildlife species, with additional emphasis on species of concern and habitats at risk. The goal is to shift focus from single-species management and highly specialized individual efforts to a geographically based, landscape-oriented,

fish and wildlife conservation effort. The Service approves CWCSs and administers SWG program funding.

In 2005, the NGPC developed a statewide CWCS called the Nebraska Natural Legacy Plan. The planning team reviewed the legacy plan and the information obtained was used during the development of the CCP. Implementation of the habitat goals and objectives in this CCP will support the goals and objectives of the legacy plan.

### **Nebraska Natural Legacy Plan**

The planning process for the legacy plan solicited public input and the help of state, federal, and nongovernmental agencies. One of the plan’s purposes was to identify areas in the state that have unique wildlife and habitat characteristics. These unique areas—“biologically unique landscapes”—are focus areas for the conservation of the state’s rarest species and natural habitats.

Nebraska’s mission to “develop and implement a blueprint for conserving Nebraska’s flora, fauna, and natural habitats” provides the state with a way to address pressing natural resource issues. When formulating proposed actions, planners must take into account the state’s strong agricultural background. Farms and ranches cover 93% of the total land area and support a significant share of Nebraska’s overall biological diversity. Maintenance of biological diversity throughout the state requires that conservation efforts be directed at a broad range of land issues and management practices on public and private lands. Maintaining and improving existing habitat on working farms and ranches is essential to conserving biological diversity and offers the greatest hope for success.

Nebraska plans to improve the efficiency and effectiveness of conservation by taking a more systematic approach to identifying and prioritizing the components of biological diversity through a “course filter/fine filter” approach. Monitoring of Nebraska lands becomes a priority as the state begins implementation of the plan. Monitoring of management actions is conducted at two levels: (1) response of individual species, and (2) response of habitats or ecological communities. Monitoring trends in abundance and distribution of different habitat types can be used to detect land use changes and can help direct conservation action toward those types that are showing the steepest decline.

Nebraska divides the state into four ecoregions for management purposes: tall-grass prairie, mixed-grass prairie, sandhills, and short-grass prairie. The ecoregions that occur in the basin—mixed-grass prairie and tall-grass prairie—are further described.

#### **Mixed-grass Prairie Ecoregion**

The mixed-grass prairie ecoregion lies between the tall-grass prairie to the east and the short-grass

prairie to the west, acting as a transition zone for the two. The region's climate is semiarid with annual average precipitation ranging from 28 inches in the east to 20 inches in the west. Average annual temperatures range from 52°F to 57°F. Starting in 1940, the government put in place policies to subsidize and facilitate conversion of marginal land such as playa wetlands to croplands. Center-pivot irrigation facilitated cultivation of steeper slopes and lands isolated from surface irrigation sources. Two-thirds of the land in the ecoregion is engaged in cropland production with most of the remaining grasslands used for livestock grazing.

### Tall-grass Prairie Ecoregion

The tall-grass prairie ecoregion covers primarily the eastern quarter of the state, with parts extending further westward. Loess and organic matter form the basis for the deep, fertile soils that typify eastern Nebraska. Annual precipitation ranges from 25–36 inches, with summer temperatures reaching highs of 90°F and dropping to lows of 10°F in winter. The tall-grass prairie ecoregion is considered to have more diversified farming operations than the western part of the state.

### Ecoregion Threats

Native animal species and ecoregion threats are essentially the same for all four regions in Nebraska. More than 300 species of resident and migratory birds have been found in the area. Most of the 55 mammal species are widespread with no distinct affiliation to the regions. Native, large predators have become extremely rare or extirpated from the regions. The 75 species of fish present in the ecoregions are “big river” generalists that can withstand a wide variation of environmental extremes. Wetlands are used for breeding by all the amphibians and reptiles. Insects are the most diverse and perhaps the most important group ecologically and economically because they play vital roles as herbivores, predators, pollinators, decomposers, soil aerators, and as food for other wildlife.

Several stresses face and affect the ecoregions, as follows:

- Conversion and fragmentation of natural habitats.
- Wetland drainage.
- Wetland sedimentation.
- Altered hydrology of wetlands.
- Fire plays an important role in prairie maintenance by promoting nutrient cycling, creating microhabitats, and increasing plant vigor and native plant diversity. Currently, less than 1% of the state's grasslands and woodlands are burned annually. Loss of fire has resulted in the degradation of thousands of acres of prairie by invasive plant species.
- Most grazing takes place in the absence of fire and with relatively little variation in timing and

intensity. Overgrazing can severely impact the composition of grasslands, and increase the amount of sediment and other pollutants entering waterbodies. Grazing systems used on prairie remnants cause losses of plant and animal diversity and ecological functions.

- Spread of invasive plants has threatened the ecoregions' biological diversity.
- Altered hydrology and channel degradation of rivers and streams cause reductions in natural flows and reduce habitat available.
- Large-scale habitat fragmentation from conversion of native habitats to crop fields, housing developments, and roads has occurred over most of the state with the exception of the Nebraska Sandhills.

## 1.5 ECOSYSTEM DESCRIPTION AND THREATS

The Service has adopted watersheds as the basic building blocks for carrying out ecosystem conservation (see figure 5, ecosystem map). The district is located within the Platte-Kansas rivers ecosystem. In addition, the Nebraska Natural Legacy Plan (2005) identifies the Rainwater Basin as one of 40 “biologically unique landscapes.”

### *PLATTE-KANSAS RIVERS ECOSYSTEM*

The Platte-Kansas rivers ecosystem includes almost all of Nebraska, southeast Wyoming, northeast Colorado, and northern Kansas (see figure 6). This ecosystem encompasses approximately 182,000 square miles and is home to the Nebraska Sandhills, the largest sand dune complex in the Western Hemisphere. The sandhills and many other areas provide vital habitat for numerous threatened and endangered wildlife and plant species.

The ecosystem spans from snow-capped, barren mountain peaks in Colorado to lowland riparian cottonwood forests along the Missouri River in eastern Nebraska and Kansas. The mountainous regions are predominately a mixture of coniferous forests comprised of Douglas-fir, ponderosa pine, lodgepole pine, Engelmann spruce, and subalpine fir. Pinyon pine and juniper woodlands and aspen communities are common throughout. Alpine meadows and lakes, willow shrub lands, and barren rocky areas are common at high elevations. Forests generally transition into shrub communities dominated by sagebrush with short grasses and forbs in eastern Wyoming and western Nebraska. Farther to the east, trees give way to short-grass prairie dominated by buffalograss, blue grama, hairy grama, and western wheatgrass. The short-grass prairie turns into mixed-grass prairie, due primarily to greater annual rainfall, in central Nebraska and Kansas.

Many federally listed endangered and threatened species including the piping plover, whooping crane,




Figure 5. U.S. Fish and Wildlife Service ecosystem map.




Figure 6. Platte-Kansas rivers ecosystem and Arkansas-Red rivers ecosystem map.

and Eskimo curlew have sought out this area as a refuge.

Threats to the Platte-Kansas rivers ecosystem that require attention include overgrazing, invasive plants, population growth and housing development, and groundwater and surface water depletion. To overcome these threats, priorities for the ecosystem are to ensure that (1) natural, healthy ecological processes dominate; and (2) economic development complements environmental protection.

The district contributes to the accomplishment of goals and objectives for this ecosystem through its Partners for Fish and Wildlife Program and existing partnerships.

### 1.6 THE PLANNING PROCESS

The steps involved in the development of this CCP are intended to comply with the Improvement Act, the NEPA, and the implementing regulations of

the acts. The Service issued a final refuge planning policy in 2000. This policy established requirements and guidance for refuge and district plans—including CCPs and step-down management plans—to ensure that planning efforts comply with the Improvement Act. The planning policy identified several steps of the CCP and environmental analysis process (see figure 7, steps in the planning process). Table 1 displays the planning process used by the Service in the development of CCPs for all units of the Refuge System.

This CCP was prepared by a planning team composed of representatives from various Service programs including district and regional office staffs and from the Rainwater Basin Joint Venture (RWBJV), U.S. Geological Survey’s biological resources division (USGS–BRD), Nebraska Game and Parks Commission (NGPC), and region 6’s division of refuge planning (see appendix C, preparers).




Figure 7. Steps in the planning process.

**Table 1. Planning process summary for Rainwater Basin Wetland Management District, Nebraska.**

<i>Date</i>	<i>Event</i>	<i>Outcome</i>
June 10, 2005	Initial conference call between the district staff and regional office planning staff.	Initiated contacts to organize development of the CCP and an overview of district issues. Started development of a mailing list.
July 13–15, 2005	Site visit to the district and initial meeting with the proposed planning team.	Acquainted regional office staff and state personnel with district activities and issues. Developed a preliminary list of qualities and issues.
October 13–14, 2005	Purposes, vision, and goals workshop; Kearney, NE.	Reviewed purposes for the district. Developed a vision statement and set of goals for the draft CCP and EA.
December 5, 2005 (5–8 p.m.)	Public scoping meeting; Kearney, NE.	Presented the district's background information and the CCP process. The public queried staff and provided comments.
December 6, 2005 (5–8 p.m.)	Public scoping meeting; York, NE.	Presented the district's background information and the CCP process. The public queried staff and provided comments.
December 7, 2005 (5–8 p.m.)	Public scoping meeting; Clay Center, NE.	Presented the district's background information and the CCP process. The public queried staff and provided comments.
December 8, 2005 (5–8 p.m.)	Public scoping meeting; Holdrege, NE.	Presented the district's background information and the CCP process. The public queried staff and provided comments.
December 6–7, 2005	CCP kickoff meeting.	Finalized the planning team. Updated list of issues and qualities. Identified biological and mapping needs. Determined the CCP steps and schedule.
February 28–March 2, 2006	Alternatives development workshop; Grand Island, NE.	Developed a range of alternatives for managing the district.
March 23–24, 2006	Impacts assessment workshop via conference calls: Kearney–Lincoln–Denver.	Assessed environmental impacts, by focus area, from each alternative developed. Recommended a proposed action.
August 1–3, 2006	Biological objectives, strategies, and rationale development workshop; Kearney, NE.	Drafted the biological objectives, strategies, rationale, and bibliography for the proposed action.
August 29–31, 2006	Nonbiological objectives, strategies, and rationale development workshop; Kearney, NE.	Drafted the nonbiological objectives, strategies, rationale, and bibliography for the proposed action.
December–March 2007	First draft CCP and EA preparation.	Prepared the first draft of the CCP and EA.
April–June 2007	Internal Service and state review of the draft CCP and EA.	Collected internal comments about the draft CCP and EA. Addressed comments; prepared the draft CCP and EA for public review.

**Table 1. Planning process summary for Rainwater Basin Wetland Management District, Nebraska.**

<i>Date</i>	<i>Event</i>	<i>Outcome</i>
July 18–August 18, 2007	Public review of the draft CCP and EA; open house meeting in Hastings, NE, about the draft CCP and EA.	Met with interested public and collected public comments about the draft CCP and EA in person and via mail and email.
August 20–September 14, 2007	Analysis of comments and finalization of the CCP.	Briefed the Service’s region 6 directorate on the outcome of the public review period. Finalized the plan with approval of the regional director.


The Service began the preplanning process in September 2005 (see appendix D, public involvement). During preplanning, the team developed a mailing list, internal issues, and a special qualities list. The planning team identified current district program status, compiled and analyzed relevant data, and determined the purpose of the refuge.

Compliance with the NEPA was achieved through involvement of the public. After the notice of intent (NOI) to prepare the draft CCP and EA was published in the *Federal Register* in November 2005, scoping began in December 2005 with public meetings. Scoping is the process of obtaining information from the public for input into the planning process. Over the course of preplanning and scoping, the planning team collected information about resources of the district and the surrounding areas, which is summarized in chapter 3.

The Service developed alternatives for management of the district after reviewing a wide range of public comments and management needs. After careful consideration of the environmental consequences of the alternatives, the Service chose alternative B as the proposed action to be implemented. The evaluation of these alternatives was documented in the “Draft Comprehensive Conservation Plan and Environmental Assessment for the Rainwater Basin Wetland Management District.” The Service published the notice of availability (NOA) for the draft CCP and environmental assessment (EA) on the same day it was released for public review—July 18, 2007. The draft CCP and EA outlined long-term guidance for management decisions, set forth proposed objectives and strategies to accomplish district purposes and meet goals, and identified the Service’s best estimate of future needs.

Public review of the draft CCP and EA concluded on August 18, 2007. After an analysis of the public’s comments, the Service’s regional director of region 6 selected alternative B (the proposed action) as the preferred alternative for the final CCP. Subsequently, the draft CCP was modified in accordance with substantive public comments to produce this final CCP, which the regional director approved in August 2007 after documentation of a “finding of no significant impact” (see appendix E, environmental compliance).

For further details of the planning process including comments and responses, see appendix D, public involvement.



*The public came to four open houses to learn about the district and offer ideas and concerns.*

## **COORDINATION WITH THE PUBLIC**

A mailing list was developed by the planning team, consisting of more than 500 names—private citizens; local, regional, and state government representatives and legislators; other federal agencies; and interested organizations (see appendix D, public involvement).

The Service held four public scoping meetings, in open-house format, during December 2005 (see table 1 for details). Attendees provided written and oral comments and were informed that comprehensive planning was an open process where they could submit their comments at any time and by any means (letter, telephone, or Internet) until the time the CCP was final.

The combined total attendance to these public meetings was 63 persons. The planning team received additional written comments via mail. Seventeen written comments were received throughout the scoping process. Input obtained from meetings and correspondence, including emails, were considered in development of this CCP.

For information about the results of the public review period and the comments received, see appendix D.

## ***STATE COORDINATION***

In November 2005, an invitation letter to participate in the CCP process was sent by the Service's region 6 director to the director of the NGPC. Two representatives from the NGPC actively participated in the development of this CCP as part of the core planning team. Local NGPC wildlife managers and the district staff maintain excellent and ongoing working relations that precede the CCP process.

## ***TRIBAL COORDINATION***

The planning team contacted Native American tribal representatives of the Pawnee Tribe and Otoe-Missouria Tribes. The tribal governments are part of the mailing list.

## ***RESULTS OF SCOPING***

Table 1 summarizes all scoping activities. Comments collected from scoping meetings and correspondence, including comment forms, were used in the development of a final list of issues that are addressed in the CCP.

The Service determined which alternatives could best address these issues. The planning process ensured that issues with the greatest effect on the district were resolved or given priority over the life of the CCP. Identified issues, along with a discussion of effects on resources, are summarized in chapter 2, section 2.6.

In addition, the Service considered suggested changes to current district management presented by the public and other groups.



## 2 The District

A wetland management district includes land that the Refuge System acquires with Federal Duck Stamp and other funds. These districts, including the Rainwater Basin Wetland Management District, are to be restored and managed primarily as prairie wetland habitat, which is critical to waterfowl and other wetland birds.

This chapter describes the establishment, vision and goals, special values, and planning issues of the district.


### 2.1 ESTABLISHMENT, ACQUISITION, AND MANAGEMENT HISTORY

The Rainwater Basin Wetland Management District was established in 1963 with the purchase of land on what is now Massie WPA. An area acquisition office of the Service was located in Hastings from 1962 to 1971, during a time when more than 14,000 acres were acquired to be managed as WPAs. Valentine National Wildlife Refuge, more than 200 miles from the nearest WPA, managed these WPAs from 1963 until 1966 when the wetland management district was staffed by one person. The district held the name of Hastings Wetland Management District until its name was changed to Rainwater Basin Wetland Management District in 1981.

The Rainwater Basin encompasses 17 counties in southern Nebraska. The district boundary includes all or portions of 13 of these counties. The district has land-purchasing authority within Adams, Clay, Fillmore, Franklin, Gosper, Hall, Hamilton, Kearney, Phelps, Polk, Saline, Seward, and York counties. The district manages WPAs in all of these counties except Polk County. In addition, the district manages one WPA that occurs outside the district boundary, in Cuming and Dodge counties.

Through the years, the Service has acquired 230 tracts of land that form 61 WPAs under the district's management. The WPAs managed by the district total 24,210.09 acres and occur across 14 Nebraska counties (see table 2). Figures 8–19 display the WPAs within each county, except for Cuming and Dodge counties, which lie outside the district boundary.

The Service's acquisition goal for the district is 24,000 acres. To date, the Service has acquired 21,703 acres of the 24,210 acres the district manages. The remaining 2,507 acres under district management have been gifted or obtained from other agencies. Three of the areas managed by the district are atypical WPAs. McMurtrey WPA in Clay County was transferred to



*Snow geese congregate at Griess WPA (Fillmore County).*

the Service from the Department of Defense (originally part of the Hastings Ammunition facility). McMurtrey WPA has no public access and is managed as a closed area. In addition, the Farmers Home Administration transferred to the Service the Haseman WPA (in Cuming and Dodge counties, northwest of Omaha) and Schwisow WPA (Saline County). These two properties are different from the other 59—they are not within the basin and they occur within floodplains of the Elkhorn and Big Blue rivers respectively. Although these two WPAs are outside the basin, they are managed in the same manner as the rest of the district's WPAs and are open to public use.

Separate from the WPAs, the district contains 35 conservation easements, which total 2,476 acres. The Farmers Home Administration transferred all these areas to the Service. While the easement restrictions vary, they generally prohibit wetland drainage, grassland conversion, and development. However, 25 of the easements allow livestock grazing. Special use permits are generally required for vegetative manipulation.

### 2.2 SPECIAL VALUES OF THE DISTRICT

The planning team and the public identified special qualities that make the district valuable for wildlife and the American people. Attributes identified included the following:

- The district provides a critical spring staging area for millions of migratory birds that gather from the coastal states, Mexico, and South America.

**Table 2. Waterfowl production areas managed by Rainwater Basin Wetland Management District, Nebraska.**

<i>County Name and WPA Summary</i>	<i>WPA Name</i>	<i>Acreage</i>
Adams: 2 WPAs; 391.56 acres	Kenesaw	231.56
	Weseman	160.00
Clay: 16 WPAs; 6,374.48 acres	Eckhardt	175.21
	Glenvil	119.46
	Green Acres	63.66
	Hansen	320.00
	Harms	60.00
	Harvard (all units)	1,484.00
	Hultine (all units)	1,000.00
	Lange	158.76
	Massie	853.00
	McMurtrey	1,067.00
	Meadowlark	80.00
	Moger	196.70
	Schuck	80.00
	Smith	476.40
Theesen	80.29	
Verona	160.00	
Cuming and Dodge: 1 WPA, 229.11 acres	Haseman	229.11
	Brauning	240.00
Fillmore: 11 WPAs; 3,578.52 acres	County Line	408.00
	Griess	20.00
	Krause	534.26
	Mallard Haven	1,087.00
	Miller's Pond	130.78
	Morphy	89.54
	Rauscher	250.75
	Real	160.00
	Rolland	128.56
	Wilkins	529.63
	Franklin: 3 WPAs; 1,783.04 acres	Macon Lakes
Quadhamer		593.56
Ritterbush		80.87
Gosper: 3 WPAs; 1,453.59 acres	Elley	60.00
	Peterson	1,156.09
	Victor Lakes	237.50
Hall: 1 WPA, 627.81 acres	Hannon	627.81
Hamilton: 3 WPAs; 1,120 acres	Nelson	160.00
	Springer	640.00
	Troester	320.00
Kearney: 9 WPAs; 2,873.75 acres	Bluestem	75.93
	Clark	451.00
	Frerichs	46.50
	Gleason	569.58
	Jensen	465.00
	Killdeer	38.36
	Lindau	152.38
	Prairie Dog	892.00
Youngson	183.00	

**Table 2. Waterfowl production areas managed by Rainwater Basin Wetland Management District, Nebraska.**

<i>County Name and WPA Summary</i>	<i>WPA Name</i>	<i>Acres</i>
Phelps: 6 WPAs; 4,606.89 acres	Atlanta	1,147.08
	Cottonwood	560.00
	Funk	1,996.40
	Johnson	577.44
	Jones	165.97
	Linder	160.00
Polk: 0 WPA	—	—
Saline: 1 WPA, 61 acres	Schwisow	61.00
Seward: 2 WPAs; 471.14 acres	Freeman Lakes	187.76
	Tamora	283.38
York: 3 WPAs, 639.2 acres (County Line WPA occurs in two counties, Fillmore and York. This WPA's occurrence and acreage is reported in total under Fillmore County.)	County Line	(see Fillmore County)
	Heron	320.00
	Sininger	160.00
	Waco	159.20

- Large concentrations of migratory birds use the basin in the spring: 90% of the midcontinental population of white-fronted geese, 50% of the midcontinental population of mallards, 30% of the continental population of northern pintails, and 90% of the midcontinental population of snow geese.
- The basin has the second highest number of confirmed sightings of whooping cranes along their migration route.
- Five of nine grassland species of concern nest in the basin.
- Two ecosystems, tall-grass prairie and mid-grass prairie, occur in the district.
- Most of the remaining prominent wetlands in the basin are under ownership and management of the district.
- The district has the ability to provide supplemental water to wetlands during migration.

Socialist Republics, and with various countries in the Western Hemisphere.”

Migratory Bird Conservation Act (16 USC 715d [2]) —“For use as an inviolate sanctuary, or for any other management purpose, for migratory birds.”

Migratory Bird Conservation Act (16 USC 715i [a]) —“Areas of lands, waters, or interests therein acquired or reserved pursuant to this subchapter shall ... be administered ... to conserve and protect migratory birds in accordance with treaty obligations with Mexico, Canada, Japan and the Union of Soviet Socialist Republics, and other species of wildlife found thereon, including species that are listed ... as endangered or threatened species, and to restore and develop adequate wildlife habitat.”

Migratory Bird Hunting and Conservation Stamp Act (16 USC 718 [c])—“Small areas, to be designated as ‘Waterfowl Production Areas’ may be acquired without regard to the limitations and requirements of the Migratory Bird Conservation Act, but all of the provisions of such Act which govern the administration and protection of lands acquired thereunder, except the inviolate sanctuary provisions of such Act, shall be applicable to areas acquired pursuant to this subsection.”

Public Land Order 6979 (May 25, 1993)—“To protect waterfowl production areas.”

Public Land Order 7206 (June 24, 1996)—“The following described public lands are hereby withdrawn from settlement, sale, location or entry under the general land laws, including the U.S. mining law, but not from leasing under the mineral leasing laws, to protect waterfowl production areas. This withdrawal will expire 50 years from the effective date of this order unless ... the Secretary determines that the withdrawal shall be extended.”

## 2.3 PURPOSES FOR THE DISTRICT

The purposes for the Rainwater Basin Wetland Management District are described in the following legislation and public land orders:

Consolidated Farm and Rural Development Act (7 United States Code [USC] 2002 [a])—“For conservation purposes any real property, or interest therein ... that has marginal value for agricultural production; is environmentally sensitive; or has special management importance.”

Emergency Wetlands Resources Act (16 USC 3901 [b]) —“It is the purpose of this chapter to promote ... the conservation of the wetlands of the nation in order to maintain the public benefits they provide and to help fulfill international obligations in various migratory bird treaties and conventions with Canada, Mexico, Japan, the Union of Soviet




Figure 8. Waterfowl production areas in Adams County, Nebraska.




Figure 9. Waterfowl production areas in Clay County, Nebraska.




Figure 10. Waterfowl production areas in Fillmore County, Nebraska.




Figure 11. Waterfowl production areas in Franklin County, Nebraska.




Figure 12. Waterfowl production areas in Gosper County, Nebraska.




Figure 13. Waterfowl production areas in Hall County, Nebraska.




Figure 14. Waterfowl production areas in Hamilton County, Nebraska.




Figure 15. Waterfowl production areas in Kearney County, Nebraska.




Figure 16. Waterfowl production areas in Phelps County, Nebraska.




Figure 17. Waterfowl production areas in Saline County, Nebraska.




Figure 18. Waterfowl production areas in Seward County, Nebraska.




Figure 19. Waterfowl production areas in York County, Nebraska.

## 2.4 VISION

*The Rainwater Basin provides critical habitat for millions of migratory birds.*

*The basin's name reflects both the basis of its wetland hydrology and natural precipitation cycles. A network of functioning wetland and prairie plant ecosystems provides a native grassland mosaic that gives the local community a sense of pride and connection to the Great Plains flora and fauna. The lands managed by the wetland management district serve as an example of land stewardship mimicking natural processes, and they provide an array of wildlife-dependent educational and recreational opportunities.*

*It is only through partnerships with individuals, agencies, and organizations that this vision can be achieved and maintained.*

## 2.5 GOALS

The following goals reflect the vision for the district—providing for healthy ecosystems and compatible opportunities for the public to appreciate and enjoy the natural environment.

### **WETLAND HABITAT GOAL**

Restore, enhance, and maintain the hydrology and early successional vegetation conditions essential to the conservation of migratory birds.

### **UPLAND HABITAT GOAL**

Reestablish and maintain native grassland communities of the Rainwater Basin.

### **WATER RIGHTS GOAL**

Develop partnerships to protect the natural hydrology of WPA watersheds and ensure the necessary water rights are in place to protect future use of both ground and surface water.

### **INVASIVE PLANT SPECIES GOAL**

Reduce and control the spread of nondesirable, nonnative plant species within wetland and upland habitats for the benefit of native plant and wildlife communities.

### **WILDLIFE DISEASES GOAL**

Work with partners to prevent or control the outbreak and spread of wildlife-borne diseases to protect human and migratory bird populations.

## **RESEARCH AND SCIENCE GOAL**

Encourage and support research that substantially contributes to the understanding and management of the Rainwater Basin wetland and grassland ecosystem.

## **CULTURAL RESOURCES GOAL**

Identify and evaluate the cultural resources in the district and protect those that are determined to be significant.

## **VISITOR SERVICES GOAL**

Provide quality wildlife-dependent recreation and educational opportunities by instilling an understanding of basic ecological processes, purpose of the Rainwater Basin Wetland Management District, and mission of the Service for persons of all abilities and cultural backgrounds.

## **PARTNERSHIP GOAL**

Promote and develop partnerships with adjacent landowners, public and private organizations, Native American tribes, and other interested individuals to protect, restore, enhance, and maintain a diverse and productive ecosystem.

## **SOCIOECONOMICS GOAL**

Obtain a better understanding of the social and economic contribution WPAs make to the people and communities within the Rainwater Basin.

## **OPERATIONS GOAL**

Safely and efficiently use funding, staffing, infrastructure, and partnerships to achieve the purpose and objectives of the Rainwater Basin Wetland Management District.

## 2.6 PLANNING ISSUES

The Service held four public meetings, sent news releases to the local and regional press, published an announcement in the “Federal Register,” and sent numerous mailings to solicit public input on important issues. Following are the most significant issues identified, which are addressed throughout this CCP.

### **HABITAT MANAGEMENT**

The district's primary purpose is to provide optimal migration habitat for waterfowl, waterbirds, and other species that depend on a grassland-wetland ecosystem. To achieve goals and objectives set for the district's habitats, there must be aggressive management. Nearly all lands bought for the district had been drained and farmed and have required extensive restoration.

Restoration work is not achieved by merely plugging a drainage ditch or planting native grasses.



*Kenesaw Waterfowl Production Area (Adams County).*

Restoration requires years of assertive management to establish native grasslands that can compete or withstand the influence of early succession, such as encroachment or dominance of weedy or woody plants. Planned grazing and burning are the two more common treatments used to reach a naturally dynamic grassland-wetland ecosystem. Staff levels are currently at a level that is more in line with managing a native grassland community rather than restoring drained and farmed lands.

There is a gap between the public's perception of wetland ecology and an understanding of how managed disturbance mimics natural disturbance and creates a healthier ecosystem. When a WPA wetland goes dry, a portion of the public expects additional water pumping be done, another portion expects management to be changed to benefit pheasants rather than waterfowl, and another portion simply concludes that no management is being done. Prescribed fire and grazing are perceived by some as habitat destruction rather than a management tool that is beneficial in sustaining these habitats.

## **WATER AND WETLAND MANAGEMENT**

Wetlands within the Rainwater Basin are in multiple ownerships and the district has been unsuccessful in obtaining complete ownership of the wetlands at many WPAs. On some WPAs, the areas were bought with little or no adjacent upland. Without complete ownership, restoration and management of the wetland is difficult or not possible. The WPAs and their watersheds (the surrounding areas that naturally drain into the WPAs' wetlands) are altered by land leveling, diversion of runoff water away from the wetland, and lowered water quality associated with agricultural runoff and sedimentation. Agricultural runoff includes effluents from feedlots. Some of the WPAs contain legal drainage tiles (underground

drainage systems that pre-date the WPAs' existence) that prevent wetland restoration. Nebraska water laws do not protect WPAs from having the natural runoff captured before it reaches the wetland.

Pumping groundwater to provide supplemental water to wetlands is critical to maintain habitat during the spring and fall migrations. However, the growing demand for groundwater is creating a conflict between agricultural needs and wildlife needs.

Increasing energy costs, antiquated equipment, and growing restrictions on groundwater usage are threatening the district's ability to provide adequate water.



*Pumping water to wetlands.*

## **INVASIVE PLANT CONTROL**

Invasive plants, especially those designated as noxious weeds, have the ability to degrade wildlife habitat and to spread into adjacent lands. This has been a significant issue for the district for years. The district directs a large portion of their resources for the control of invasive plants. Integrated pest management (IPM) strategies currently include prescribed fire, grazing, mowing, herbicides, biological control using introduced insects, "interseeding," and farming.

The establishment of new invasive plants—such as salt cedar and purple loosestrife—is a constant threat. Generally, an immediate control response to new invasive plants is most effective in the long term.

## **WILDLIFE DISEASE CONTROL**

Since 1975, the district has had a history of avian cholera. In 1980 and 1998, avian cholera outbreaks killed over 100,000 birds during the spring migrations. The high concentration of birds each spring poses a threat of disease outbreaks and the spread of disease in the Central Flyway. Management actions taken have included increased pumping to improve water quality and quantity and collection of infected carcasses to control the spread of disease.

In 2005, there was a positive case of chronic wasting disease found in Hall County within the Rainwater Basin. The disease is expected to continue to spread eastward across Nebraska.

In the future, the H5N1 strain of avian influenza is expected to migrate from the Eastern Hemisphere to the Western Hemisphere. Because of the high concentration of birds found throughout the basin during spring migration, there is a concern about the spread of the disease.

## **SPECIES OF CONCERN**

Species of concern are the prairie dog and federally listed threatened and endangered species.

In July 1998, the National Wildlife Federation petitioned the Service to list the black-tailed prairie dog as a threatened species under the Endangered Species Act. In the fall of 1999, the Service's regional director (region 6) issued a moratorium on all control of black-tailed prairie dogs on Service lands. In February 2000, the Service concluded that this species warranted listing, but was precluded from being listed due to concerns and resource constraints related to other higher priority species. In August 2004, an updated evaluation of the best available scientific information led the Service to determine that the black-tailed prairie dog should be removed as a candidate for listing.

Prairie dog colonies are located on five WPAs scattered throughout the district. The five sites had historical use, and management has been done to allow the dog towns to continue. The spread of the populations to private lands has been controlled by two factors: (1) the sites are surrounded by cropland; and (2) the precipitation level causes native grasses to grow faster than the prairie dogs can remove it. The tall vegetation keeps the town to a manageable size. A management plan written in 2003 is included in appendix F.

The CCP will not address specifically this species any further because (1) the district currently manages black-tailed prairie dog colonies in accordance with the existing management plan (appendix F), and (2) the existing colonies within the district are stable.

The district staff, in consultation with staff from region 6's ecological services, reviewed all threatened and endangered species with historical ranges on or

near the district to determine if additional actions could be taken to restore or enhance habitat for endangered species. These species are the whooping crane, interior population of the least tern, American burying beetle, and western prairie fringed orchid. No species were identified as requiring actions different from those being taken to meet the purposes of the district (see appendix G, section 7 biological evaluation).

The "Biological Integrity, Diversity, and Environmental Health Policy" (USFWS 2001) guides Service personnel in carrying out the clause of the Improvement Act that directs the Secretary of the Interior to ensure the maintenance of the "biological integrity, diversity, and environmental health" of the Refuge System. This policy guides the Service to consider restoration of lost or severely degraded components of the Refuge System "where appropriate and in concert with refuge purposes and the Refuge System mission"; this includes federally listed species.

## **RESEARCH AND SCIENCE**

The Rainwater Basin serves as a critical staging area during spring migration. While the birds are in the basin, they feed extensively in surrounding croplands and within the wetlands. When they are not feeding, these birds roost on the larger wetlands. There is little information to determine whether the basin is meeting the needs and energetics of waterfowl and other migratory birds. Such information will help direct management actions by the district.

Water-pumping decisions rely heavily on intuition because of the limited scientific information that is available. The district's intent for pumping is to provide water in those wetlands that provide the most food and resting area for birds. It is unknown how the hydrology of individual wetlands in the basin, in combination with water pumping, can provide optimal habitat conditions. For example, is it the wisest use of water to pump a wetland with wet soils and less preferred plant species or a wetland with dry soil and preferred plant species.

The distribution and abundance of amphibians, reptiles, invertebrates, and small mammals on the WPAs is unknown.

As part of the CCP development process, a Service-funded socioeconomic study determined the extent to which the existence and operations of the district benefit the local and state economies. Excerpts from this report are included in chapter 3, section 3.7.

## **VISITOR SERVICES**

Hunting, fishing, wildlife observation, photography, environmental education, and interpretation are all uses allowed on WPAs. The high concentration of birds and the limited public lands available for public recreation in the basin makes this an issue of interest.

There is a demand for increased and improved recreational hunting opportunities on the WPAs. By regulation, lands acquired as WPAs are open to public trapping as well.

The public has a significant misunderstanding about the types and management of public lands in the basin, particularly those managed by the district and the NGPC. This is an issue because both agencies manage public uses differently and their respective missions, while complementary, are not exactly alike. This fact often causes confusion between wildlife enthusiasts and the public in general.

Many people in the local communities are not aware of the district's existence because district personnel are based more than 20 miles away from the WPAs. In addition, the district office is not easily located in an older industrial portion of Kearney.

### ***PARTNERSHIPS***

The scattering of small public areas (WPAs) among privately owned lands increases the need to build partnerships throughout the Rainwater Basin. The district cannot address many of the opportunities without the help of partners. This is especially the case for visitor services, particularly in environmental education. Public involvement will strengthen local interest and increase the district's ability to manage and promote wildlife resources. The district is not fully using the skills and resources of other groups, organizations, and local communities. In turn, the district is not providing the resources and expertise that would help local conservation groups meet their objectives.

### ***OPERATIONS***

The basin encompasses 4,200 square miles in a 17-county area of southern Nebraska. The district extends across 13 of those counties. The distance from the easternmost to the westernmost WPA is 133 miles. The logistics of transporting equipment and traveling to WPAs makes it difficult to effectively manage the properties. It takes the district staff 2 hours each way to reach the farthest WPA, which reduces to 4–6 hours the amount of time to work at an area. Because of the long distances, fuel costs are a major burden. In a

typical year, the staff drives approximately 110,000 miles.

The district faces immediate challenges with its headquarters. The Service leased the office/shop facility in 1977 when the staff consisted of four people. The office portion was an open foyer and three rooms. In 1994, to accommodate the new private lands staff and migratory bird biologist, a portion of the shop was converted to office space. In 2000, to accommodate the new fire staff and station biologist, the library/conference room was converted into a cramped, open-office setting for four desks. There are currently 12 people in the staff.

The heating system for the office was installed in the shop portion of the building. The office area has only one exit or opening leading to the outside. No windows exist in any of the offices. There is no exchange of fresh air, so the furnace recycles office and shop air continually throughout the office portion.

The development of a fire program brought with it two fire engines that have to be stored inside at all times—reducing the amount of working and storage area for other equipment. During the spring fire program, the shop area is crowded with fire engines, water tenders, and all-terrain vehicles (ATVs) to prevent freezing during the cold nights. The shop is the only storage facility on the site and ATVs used for weed spraying have to be stored in the shop during summer months. The high temperatures inside the shop during the summer cause the herbicide mixture to volatilize and enter the office.

The district headquarters facility is located in an urban setting that consists of a mixture of trailer homes and an industrial park. Its location does not invite the public to visit the office and public visitation is very low. Theft and vandalism have increased.

The headquarters site does not include any cold storage facility and some equipment has to be stored at two WPAs—one in the eastern portion of the district and the other in the western portion. All heavy equipment is stored outside in an unsecured area. The storage facility in the east (McMurtrey WPA) has access only through a courtesy agreement with neighboring landowners.



# 3 District Resources and Description



Gary Eslinger/USFWS

*The stiff sunflower is a native forb in the Rainwater Basin.*

The Rainwater Basin Wetland Management District manages 61 noncontiguous tracts of federal land totaling 24,210.09 acres in south-central Nebraska. All of these lands within the Rainwater Basin are WPAs, which each typically contain one large wetland. Together, these WPAs are managed as a grassland ecosystem designed to provide optimal habitat for waterfowl and shorebirds. Uplands (nonwetlands) are managed for a high diversity of native grass species and grassland birds. The district also manages 35 conservation easements totaling 2,476 acres.

This chapter describes the physical environment and biological resources of the basin and the district. In addition, this chapter describes the district's fire and grazing history, cultural resources, special management areas, visitor services, socioeconomic environment, and operations.

## 3.1 PHYSICAL ENVIRONMENT

The Rainwater Basin is located near the center of the Great Plains. Its geographical region encompasses approximately 4,200 square miles covering portions or all of 17 counties in south-central Nebraska. The

widest span is 160 miles across, extending from Gosper County to central Seward County. The northern edge parallels the central Platte River. The southern edge lies about 10 miles from the Kansas border.

## GLOBAL WARMING

The U.S. Department of the Interior issued Order No. 3226 in 2001 requiring federal agencies under its direction that have land management responsibilities to consider potential climate change effects as part of long-range planning endeavors.

The Department of Energy's report, "Carbon Sequestration Research and Development" (USDOE 1999), concluded that ecosystem protection is important to carbon sequestration and may reduce or prevent loss of carbon currently stored in the terrestrial biosphere. The report defines carbon sequestration as "the capture and secure storage of carbon that would otherwise be emitted to or remain in the atmosphere."

The increase of carbon dioxide (CO<sub>2</sub>) within the earth's atmosphere has been linked to the gradual rise in surface temperature commonly referred to as "global

warming.” In relation to comprehensive conservation planning for Refuge System units, carbon sequestration constitutes the primary climate-related effect to be considered in planning.

Vegetated land is a tremendous factor in carbon sequestration. Large, naturally occurring communities of plants and animals that occupy major habitats—grasslands, forests, wetlands, tundra, and desert—are effective both in preventing carbon emission and in acting as biological “scrubbers” of atmospheric CO<sub>2</sub>.

One Service activity in particular—prescribed burning—releases CO<sub>2</sub>, directly to the atmosphere from the biomass consumed during combustion. However, there is no net loss of carbon because new vegetation quickly germinates and sprouts to replace the burned-up biomass. This vegetation sequesters an approximately equal amount of carbon as was lost to the air (Dai et al. 2006).

Several other effects of climate change may need to be considered in the future:

- Habitat available in lakes and streams for cold-water fish such as trout and salmon could be reduced.
- Forests may change, with some plant species shifting their range northward or dying out and other trees moving in to take their place.
- Ducks and other waterfowl could lose breeding habitat because of stronger and more frequent droughts.
- Changes in the timing of migration and nesting could put some birds out of synchronization with the life cycles of their prey.

## **CLIMATE**

The basin has a continental climate characteristic of extreme temperature changes through the seasons and relatively low precipitation rates. The district averages 21–28 inches (west to east) of precipitation annually. Eighty percent of the precipitation occurs between April and September. The change in precipitation amount across the district is primarily responsible for the shift from mixed-grass prairie in the west to tall-grass prairie in the east.

Winter temperatures average 27°F, while summer temperatures average 75°F. Daily minimum temperatures above 32°F occur 136–177 days. Relative humidity averages 55% at midafternoon and 80% at dawn across most of the basin. Annual evaporation in the basin exceeds rainfall accumulations. Evaporative losses in Phelps County were computed by the Thornthwaite method. These losses average more than 5.3 inches per month (June–August). Annual free-water evaporation from small bodies of water average 46 inches; about 77% of that amount is lost from May through October. Evaporative losses can significantly reduce the amount of pooled surface water.

## **PHYSIOGRAPHY, GEOGRAPHY, AND SOILS**

The 59 WPAs found within the basin contain the same geological and soil characteristics. The Rainwater Basin lies in the flat to gently rolling, mixed-grass, loess plains of south-central Nebraska. This area is geologically new and has not developed a complete system of streams to drain surface water. It is from this characteristic that the area received its name—Rainwater Basin.

Wind-deposited Peorian Loess occurs extensively across the basin and has about 10,000 years of stability (Keech and Dreezen 1959). Upland soils that formed in wind-deposited material include Crete, Hastings, Holdrege, Hord, and Uly (Kuzila 1984). The soils are suitable for farming, with about 80% of the land being cropped.

The shallow, flat depressions formed by wind scouring are often referred to as playa wetlands because of their ephemeral (lasting for a brief time) nature. Radiocarbon dating indicates the wetlands were created near the end of the Ice Age, 20,000–25,000 years ago. Some depressions may have been enlarged and new ones created as recently as 3,000 years ago (Farrar 1996a). Over thousands of years, minute clay particles accumulated in the bottoms of the depressions, effectively sealing them off and preventing water from seeping away. The impervious clay layers are 6–72 inches thick. The wetland soils are predominantly Butler, Fillmore, Scott, and Massie (Kuzila 1994, Kuzila and Lewis 1993).

The Service and RWBJV grouped the basin’s land cover into 22 categories (see table 3). Cropland covers 80% of the landscape. Grassland covers 10% and roads cover 2.6%. Nonfarmed wetlands represent 1%. Figures 20–22 show the land cover, grouped into 10 simplified categories, across the basin.

## **WATER RESOURCES**

This section describes the relationships between the hydrology, wetlands, and water quality in the basin in relation to the district’s WPAs. In addition, this section describes the water rights situation.

### **Hydrology**

As a whole, groundwater levels throughout most of the basin have little influence on wetlands. Artificial groundwater mounds occur near irrigation delivery canals and some wetlands do benefit from this water source. Most of the basin has groundwater located more than 50 feet deep; some is 400 feet or deeper (see figure 23, groundwater map). One area east of the Tri-County canal (including Johnson WPA and Funk WPA) has groundwater levels that are less than 50 feet deep (Ekstein and Hygnstrom 1996), showing a rise of more than 50 feet from the period before development to now (see figure 23, groundwater map). Recent (2000–2006) groundwater levels have shown a decline from

**Figure 20. Land cover in the western portion of the Rainwater Basin.**

**INSERT FIGURE 20  
11X17 MAP**

**Figure 21. Land cover in the northeastern portion of the Rainwater Basin.**

**INSERT FIGURE 21  
11X17 MAP**

**Figure 22. Land cover in the southeastern portion of the Rainwater Basin.**

**INSERT FIGURE 22  
11X17 MAP**

**BLANK BACK OF  
FIGURE 22**

**Table 3. 2004 Land cover in the Rainwater Basin.**

<i>Land Cover</i>	<i>Acres</i>	<i>Percent</i>
agriculture (crop)	3,114,972	79.9
Conservation Reserve Program	27,265	0.7
road	102,838	2.6
developed urban area	27,969	0.7
developed rural area	80,873	2.1
other developed area	3,041	0.1
woodland forest	48,325	1.3
range/grass/pasture	384,031	9.8
sand pit	84	0.0
lagoon	486	0.0
irrigation reuse pit	5,487	0.1
reservoir	3,887	0.1
stock pond	11,298	0.3
farmed wetland	7,424	0.2
early successional hydrophytes*	27,189	0.7
late-successional hydrophytes*	8,513	0.2
riparian canopy	45,253	1.2
riparian shrubland	245	0.0
river channel	249	0.0
sandbar	46	0.0
wet meadow	971	0.0
floodplain marsh	189	0.0
<b>Total</b>	<b>3,900,635</b>	<b>100.0</b>

(Source: Bishop and Reker 2006.)

\*A hydrophyte is a plant that is adapted to grow in water.

5–15 feet throughout most of the basin, with a few areas decreasing more than 25 feet (UNL 2006a).

The development of center-pivot irrigation in the last third of the century has placed great demand on the groundwater underlying the basin. Irrigation, compounded by extensive drought conditions, has caused the state legislature to pass legislation to begin to control the declines in groundwater. Eleven river basins within the state have been determined to be overappropriated (that is, permitted uses for the water in the stream or river exceeds the amount of water in the stream or river). Nebraska's natural resource districts in those river basins are required to develop an integrated surface water and groundwater management plan for each river basin. Only the extreme western edge of the Rainwater Basin Wetland Management District lies within an overappropriated river basin. However, the natural resource districts within the basin have placed a moratorium on new wells and require stricter monitoring on water pumping.

There are 71 registered groundwater wells scattered over 39 of the district's WPAs. Only 26 WPAs have water-pumping capability and only 23 of these WPAs can pump water due to ownership issues relating to the "hydric footprint" (soil characteristics that indicate the existence of a wetland basin). These existing wells have the ability to deliver water to 2,230 wetland acres (approximately 20% of the total WPA wetland acres). Each well is metered and complies with state regulations for irrigation wells.

Groundwater recharge may occur in the basin and is being investigated in a 2-year research project. Research from playa lakes in Texas and New Mexico indicates that playas recharge groundwater by playa water percolating through the soils (referred to as "interstitial and macropore flow") (Wood 2000).

### Wetlands and Water Quality


Mapping done in the middle part of the last century indicated that the basin contained about 100,000 acres of wetland habitat in 4,000 basins (Schildman and Hurt 1984). However, modern soil surveys indicate two to three times as many wetlands existed (Gersib et al. 1990). Ducks Unlimited inventoried the hydric soils (those characterized by considerable moisture) and computed the historical number of wetlands to be 204,436 acres (personal communication with Darin Blunk, Bismarck, ND, 2004). Figure 24 displays the historical wetlands in the basin.

In 2004, a spring habitat assessment by the Great Plains GIS office found that only 1,693 basins (16% of the historical number) contained some wetland function in the form of retained water; the area totaled 27,839 total acres. The public wetlands provided 45% of the waterfowl habitat but represented only 9% of the 1,693 wetlands. Figure 25 displays the current wetlands in the basin.

Agricultural development has been responsible for most of the wetland loss (Schildman and Hurt 1984). Modifications within the watersheds and the wetlands have caused many wetlands to become more ephemeral in nature. Land leveling and the diversion of runoff to concentration pits and road ditches are the primary causes (Raines et al. 1990). The Great Plains GIS office has identified 11,859 concentration pits totaling 7,506 acres within the basin. The pits' water storage capacity is about two-thirds of the historical storage capacity of wetlands in the basin. There are 627 pits within individual WPA watersheds. Agricultural runoff has increased siltation and deposited related chemicals, which has resulted in poor water quality and partial filling of many wetlands (Gersib 1991, Frankforter 1996).

In their natural state, the larger wetlands collected runoff from several square miles. If precipitation was adequate, the wetlands probably held water throughout most years. The most common wetlands are small—covering less than 40 acres. The larger, less common wetlands reach nearly 1,000 acres.

On average, each WPA contains one large, seasonal or semipermanent wetland. Wetlands under district ownership total 11,117 acres. The ratio of wetland to upland is 1:1. The wetlands occupy about 16% of their watershed, with the wetland receiving its runoff water from agricultural lands. Waterfowl production area watersheds range in size from 153 to 15,852 acres, with an average size of 3,024 acres.



**Figure 23. Groundwater-level changes in Nebraska—predevelopment to spring 2006.**  
 (Source: University of Nebraska–Lincoln [UNL] 2006b.)




Figure 24. Historical wetlands in the Rainwater Basin, Nebraska.




Figure 25. Current wetlands in the Rainwater Basin, Nebraska.

Dikes, ditches, and water control structures exist at the larger WPAs. Divided ownership of a large wetland has caused the district to try to restore its portion of the wetland to the maximum level it can. For example, surface drains occur on 13 WPA wetlands and affect the hydrology to some degree. Most of the drains only function when the wetlands are more than 80% full.

In the early 1900s, counties authorized drainage districts to facilitate wetland drainage. Each district was specific to one large wetland, with drainage being done with subsurface tile drains. Five WPAs—Miller's Pond, Nelson, Springer, Troester, and Wilkins—have active tile drains. Tile drains as deep as 50 feet were tunneled through hills (Farrar 1982, 1996b). The age and structural integrity of the drains cause concern for their future. The district's desire to abandon drains is in conflict with other landowners owning a portion of the wetlands.



This water control structure at Funk WPA (Phelps County) helps direct water to three different units.

### Water Rights

The district does not have any surface water rights because no water from streams is diverted to wetlands. However, the district has water contracts with Central Nebraska Public Power and Irrigation District for the delivery of 762 acre-feet annually to Funk WPA and Victor Lakes WPA. The water contracts transferred with the property when it was bought. The district has filed to cancel the contracts because the timing and amount of water delivered does not benefit waterfowl. These contracts are scheduled to end within 5 years.

There are no well-defined legal rights protecting surface water in wetlands. For example, where the Service owns half of a wetland, the neighboring landowner might dig a pit for irrigation re-use or for a more permanent source of livestock water or for hunting. The neighbor captures the first runoff, and the Service-owned wetland will only receive runoff water after the neighbor's pit is filled. If there is not enough runoff to overflow the pit, the Service-owned wetland will get little or no water, especially when most of the runoff comes from the neighboring land.

Nebraska's water law requires irrigators to control irrigation runoff. Infractions are difficult to enforce and the irrigator often views the runoff flowing onto public lands as a benefit to wildlife. However, the law does allow natural runoff to be diverted or used by anyone within a closed watershed. A closed watershed does not drain into a perennial or permanent stream. In an open watershed, runoff from a normal rain event runs off the land into a stream. In the future, water may become valuable enough to cause landowners to capture and use runoff before it can get to the WPAs.

Groundwater rights differ from surface water rights. Nebraska's water law only requires that irrigation wells have a permit before being drilled. After well installation, the owner is allowed "reasonable and beneficial use of the groundwater." High water demands and drought conditions are expected to cause natural resource districts to place more restrictions on groundwater use. Currently, the natural resource districts have a moratorium on new wells.

### AIR QUALITY

The National Ambient Air Quality Standards include maximum allowable pollution levels for particulate matter, ozone, sulfur dioxide, nitrogen dioxide, lead, and carbon dioxide.

Particulate matter (PM) is a generic term for a broad class of chemically and physically diverse substances that exist as discrete particles (liquid droplets or solids) over a wide range of sizes. The smaller the size of the particle, the deeper it can penetrate into lung tissue. There are no standards for PM size in relation to prescribed fire because prescribed fires release many sizes of PM. However, there are regulations for the amount of different sizes of particles. The standards for PM<sub>2.5</sub> (published in the *Federal Register* on October 17, 2006) set a 24-hour standard of 35 microns/cubic meter, with an annual standard of 15 microns/cubic meter.

The Rainwater Basin in general has good air quality, with no nearby manufacturing sites or major air pollution sources. Feedlots near the WPAs produce odors and can produce particulate matter. The state of Nebraska has minimal smoke management guidelines and regulations that pertain to agricultural burning and the prevention of smoke lingering on roadways and other sensitive areas. Windblown dirt from cropland and carbon from automobiles and diesel engines contributes to particulate matter.

## 3.2 BIOLOGICAL RESOURCES

This section describes vegetation, wildlife, and their associated communities at the district. Species lists are found in appendix H (plants), appendix I (amphibians, reptiles, and mammals), and appendix J (birds).

## WETLANDS AND ASSOCIATED VEGETATIVE COMMUNITIES

The district manages 11,117 acres of hydrophytes (wetland plant species). During drawdown or drought, emergent vegetation dominates the wetlands. In wet years, submerged vegetation can occur. The 2004 vegetation mapping established community names and determined the acreage for each vegetative zone (see table 4). These names or categories have been cross-referenced with the National Vegetation Classification System (NVCS). Appendix K contains more explanation.

Native wetland plants occur on 9,148 acres. Currently, cattail dominates 657 acres and river bulrush dominates 1,145 acres. Moist-soil plants (5,014 acres) and wet meadow (1,718 acres) are the more prevalent associations. Reed canarygrass and reeds add an additional 1,965 acres of wetland habitat but are considered introduced species and are targeted for reduction (see table 4). Other invasive plants that occur in wetlands at the WPAs include Canada thistle, purple loosestrife, and “volunteer” deciduous trees.

**Table 4. Habitat types at the district in 2004.**

<i>Habitat Type</i>	<i>Acres</i>	<i>Percent</i>
buildings	2.7	0.01
bull thistle	0.4	0.00
Canada thistle	67.1	0.29
cattail	656.6	2.83
leafy spurge	0.1	0.00
moist-soil plants	5,014.5	21.62
musk thistle	10.6	0.05
parking lot	0.3	0.00
reed species ( <i>Phragmites</i> spp.)	1.1	0.00
reed canarygrass	1,963.8	8.47
road	3.2	0.01
bulrush species	1,145.0	4.94
trees	554.3	2.39
water or mud flat	585.0	2.52
wet meadow	1,717.8	7.41
annual weeds	99.5	0.43
cropland	478.1	2.06
introduced forbs	4.4	0.02
invasive cool-season plants	3,192.9	13.77
native grassland	232.4	1.00
newly seeded	370.7	1.60
planted native cool-season plants	409.2	1.76
planted native warm-season plants	6,668.9	28.75
prairie dog town	14.9	0.06
Total	23,193.5	100.00

## UPLAND AND ASSOCIATED VEGETATIVE COMMUNITIES

The district manages 12,044 acres of upland habitat at the WPAs. Vegetation includes native, cool- and warm-season grasses, forbs, planted and “volunteer” trees, and invasive plants. Native, warm-season grass stands are dominated by big bluestem, little bluestem, Indiangrass, switchgrass, Junegrass, needle and thread, and leadplant. Undesirable and invasive plant species include Canada thistle, musk thistle, leafy spurge, smooth brome, intermediate wheatgrass, and Kentucky bluegrass.

During the 1960s and ‘70s, the district planted native warm-season grass and dense nesting cover on many areas to promote waterfowl production. These plantings lacked vegetative diversity and frequently became dominated by one or two species, which reduced the grassland’s attractiveness to a variety of wildlife species. Most of these stands have been “interseeded” with a high-diversity (more than 80 species) grassland mix.

Between 400 and 700 acres are farmed, usually with soybeans. Farming is done for 2–4 years to reduce the seed source of weed seeds in the soil and to prepare the land to reseed with native grasses and forbs.

The WPAs have undesirable woody species—green ash, cottonwood, willow, eastern red cedar, and elm. These species are continually controlled or removed throughout the district. Native tree species such as buffaloberry, mulberry, chokecherry, and American plum occur and are not targeted for removal; however, some of these native trees may be inadvertently cleared during large-scale reduction treatment of undesirable trees.

### SHRUB AND TREE PLANTINGS (SHELTERBELTS)

Shelterbelts were planted along the boundaries of many of the district’s WPAs in the 1970s. The shelterbelts are mostly mature eastern red cedars that have shaded out native vegetation. Since that time, the district has realized that tree planting is not compatible with the purposes of the district.

### FIRE AND GRAZING HISTORY

Historically, grassland species in the northern Great Plains co-existed or adapted to various disturbance regimes such as fire and large-scale grazing. Settlement and the expansion of agriculture have removed fire and high-intensity, short-duration grazing from the basin.

#### Fire

Whether lightning-induced or deliberately set by Native Americans, fire has influenced the grassland ecosystem of the basin. A handful of fire-tolerant shrubs such as chokecherry, American plum, and leadplant are present; other woody species that are

killed by fire are now restricted to areas protected from fire.

It is estimated that presettlement (historical) fire frequency on mixed-grass prairie was 5–7 years. These fires sustained diverse and healthy grasslands. The district uses prescribed fire to simulate the historical influence wildland fires had on the plant communities. Most prescribed fire treatments are completed during late winter through green-up in spring. Spring presents opportunities to complete prescribed burns when temperatures are low, humidity is high, and the fire is easier to control. In addition, the timeframe coincides with other district management activities. During the last 10 years, district fire treatments have increased from about 6 prescribed burns to 25–35 burns each year, covering 4,000–5,500 acres. District experience has shown that prescribed burning of residual wetland vegetation helps promote annual plants during drier years.

### Grazing

Similar to fire, grazing greatly influenced the structure and composition of grassland communities. Herbivores such as the bison, elk, deer, pronghorn, and black-tailed prairie dog maintained grassland vigor and created patches of varying stand height and density. Grazing influenced the soil productivity, plant diversity, animal diversity, and other processes to produce unique successional patterns in the landscape at multiple scales.

Vegetation in grassland communities has developed growth strategies that allows for grazing at or near the ground without killing the plants. Some plant species contain bitter or toxic substances that cause animals to avoid grazing on them. Some species have spines that cause injury to a grazing animal's mouth. Small mammals and deer “naturally” graze at the WPAs. Additional grazing by large herbivores such as cattle is needed, at times, to maintain wetland and upland vegetation.

The district staff works with local cattle producers to provide grazing disturbance. Upland grazing is generally conducted during spring and early summer, and again in fall, to (1) stress exotic cool-season grasses or invasive plants, and (2) increase the vigor of native warm-season grasses and forbs. Wetland grazing occurs for much of the growing season to stress and physically injure invasive or aggressive wetland species such as cattail. This grazing strategy favors early successional species that provide wildlife food, create diverse habitats, and limit the expansion of invasive plants.

### WILDLIFE

This section describes the animals that are common at the WPAs, as well as those that are uncommon, threatened, or endangered.

### Amphibians and Reptiles

Amphibians that occur in the basin include the plains spadefoot toad, Woodhouse's toad, Great Plains toad, Blanchard's cricket frog, western chorus frog, bullfrog, and plains leopard frog. Appendix I contains a list of amphibians and reptiles that occur in the basin.

Semipermanent wetlands provide habitat for the painted turtle, chorus frog, common gray treefrog, and snapping turtle. The ornate box turtle lives in grassland areas. The western garter snake, red-sided garter snake, bullsnake, and eastern yellowbelly racer are common. The western hog-nosed snake occurs less frequently and prefers dry sandy prairies. The two rare snakes that can occur are smooth green snake and red-bellied snake.

The lesser earless lizard may occur in open sandy soil with little or sparse vegetation. The six-lined racerunner can be found in both lowland and upland sites.

### Mammals

The most common, larger mammals found at the WPAs are white-tailed deer, coyote, raccoon, striped skunk, eastern cottontail, American badger, and Virginia opossum. Mule deer have been seen at the western WPAs but are uncommon. During wet periods, muskrat and mink are common. The black-tailed prairie dog occurs at five WPAs: McMurtrey, Prairie Dog, Atlanta, Hultine, and Clark. Appendix I contains a list of mammals that occur in the basin.



*Muskrat*

Less common mammals are red fox, black-tailed jackrabbit, woodchuck, Franklin's ground squirrel, and eastern fox squirrel. Small mammals that are common include thirteen-lined ground squirrel, northern pocket gopher, plains pocket gopher, Ord's kangaroo rat, meadow jumping mouse, meadow vole, northern grasshopper mouse, and white-footed mouse.

## Birds

The Rainwater Basin is internationally known for its spectacular bird migrations—not only numbers of birds but also the large proportion of the continental population of prominent species. Thirty species of shorebirds use wetlands in the basin. Of the 329 species of birds that have been observed in the basin (Sharpe et al. 2001), 96 species nest (Mullhoff 2001). Appendix J contains a list of birds that occur in the basin.

Common grassland species include grasshopper sparrow, dickcissel, western meadowlark, bobolink, northern bobwhite, ring-necked pheasant, field sparrow, and northern harrier. The greater prairie-chicken is becoming more common and is found at these WPAs: Harvard, Hultine, Funk, Quadhamer, Prairie Dog, Lindau, and Jensen.

The spring migration chronology usually starts with a buildup of Canada geese on the Platte River until the basin's wetlands begin to thaw. Numbers of snow geese, white-fronted geese, and mallards begin to peak by mid- to late-February. In early March, northern pintail numbers peak, followed by Ross's geese and green-winged teal. The numbers of other divers and puddle ducks usually peak during mid- to late-March. Cinnamon teal may be seen as far east as Harvard WPA in late March, but is more common west of Hastings.

Most shorebirds pass through the basin between April 15 and May 15. According to Jorgensen (2004), the most common, spring, shorebird migrants are the following species:

- black-bellied plover
- American golden-plover
- semipalmated plover
- greater yellowlegs
- lesser yellowlegs
- willet
- upland sandpiper
- Hudsonian godwit
- dunlin
- short-billed dowitcher
- long-billed dowitcher
- Wilson's snipe
- Wilson's phalarope
- semipalmated sandpiper
- least sandpiper
- white-rumped sandpiper
- Baird's sandpiper
- stilt sandpiper
- buff-breasted sandpiper

Rainwater Basin has one of the largest concentrations of buff-breasted sandpiper during its spring migration. Common, late-summer migrants are the following: greater yellowlegs; lesser yellowlegs; Wilson's snipe; long-billed dowitcher; and sandpipers (solitary, upland, least, semipalmated, stilt, and pectoral).

The peregrine falcon frequents wetlands during peak, shorebird migration periods. In contrast, the prairie



Donna Dewhurst/USFWS

*Greater Yellowlegs*

falcon is most numerous in late January when horned lark and meadowlark are common. The merlin is primarily a winter visitor and a spring migrant (Johnsgard 1997). The burrowing owl nests at Prairie Dog WPA and Atlanta WPA. The short-eared owl is common at many of the larger WPAs during the winter months.

The Harris's sparrow can be seen at the eastern WPAs that have brushy growth or American plum thickets.

### Threatened and Endangered Species

In Nebraska, there are nine state- and federally listed endangered species, three state- and federally listed threatened species, five state-listed-only endangered species, and nine state-listed-only threatened species (see table 5).

The state- and federally endangered whooping crane and piping plover use wetlands at the basin. Forty-two percent of confirmed observations of whooping crane in Nebraska have been at the basin's wetlands (Richert 1999). Most of these sightings occur the first two weeks of April, or late October through mid-November. Piping plovers are rarely seen at the basin's wetlands due to their size and the number of other shorebirds that use mud flat habitats in late April through mid-May (Johnsgard 1997).

The Service removed the bald eagle from the endangered species list on August 8, 2007. However, the district and its partners will follow the bald eagle management guidelines recently published by the Service.

**Table 5. Nebraska’s state-listed and federally listed species.**

<i>Common Name</i>	<i>Scientific Name</i>	<i>State Status</i>	<i>Federal Status</i>
<b>PLANTS—7 Species</b>			
Hayden’s (blowout) penstemon	<i>Penstemon haydenii</i>	endangered	endangered
Colorado butterfly plant	<i>Gaura neomexicana</i> ssp. <i>coloradensis</i>	endangered	endangered
saltwort	<i>Salicornia rubra</i>	endangered	—
western prairie fringed orchid	<i>Platanthera praeclara</i>	threatened	threatened
Ute lady’s tresses	<i>Spiranthes diluvialis</i>	threatened	threatened
ginseng	<i>Panax quinquefolium</i>	threatened	—
small white lady’s slipper	<i>Cypripedium candidum</i>	threatened	—
<b>INSECTS—2 Species</b>			
American burying beetle	<i>Nicrophorus americanus</i>	endangered	endangered
Salt Creek tiger beetle	<i>Cincindela nevadica lincolniiana</i>	endangered	—
<b>REPTILES—1 Species</b>			
massasauga	<i>Sistrurus catenatus</i>	threatened	—
<b>FISH—7 Species</b>			
pallid sturgeon	<i>Scaphirhynchus albus</i>	endangered	endangered
Topeka shiner	<i>Notropis topeka</i>	endangered	endangered
sturgeon chub	<i>Macrhybopsis gelida</i>	endangered	—
blacknose shiner	<i>Notropis heterolepis</i>	endangered	—
lake sturgeon	<i>Acipenser fulvescens</i>	threatened	—
northern redbelly dace	<i>Phoxinus eos</i>	threatened	—
finescale dace	<i>Phoxinus neogaeus</i>	threatened	—
<b>BIRDS—5 Species</b>			
Eskimo curlew	<i>Numenius borealis</i>	endangered	endangered
whooping crane	<i>Grus americana</i>	endangered	endangered
interior least tern	<i>Sterna antillarum athalassos</i>	endangered	endangered
piping plover	<i>Charadrius melodus</i>	threatened	threatened
mountain plover	<i>Charadrius montanus</i>	threatened	—
<b>MAMMALS—4 Species</b>			
black-footed ferret	<i>Mustela nigripes</i>	endangered	endangered
swift fox	<i>Vulpes velox</i>	endangered	—
river otter	<i>Lutra canadensis</i>	threatened	—
southern flying squirrel	<i>Glaucomys volans</i>	threatened	—

### 3.3 CULTURAL RESOURCES

Archaeological and architectural remains representing 12,000 years of human occupation are potentially located in the district. There has been very limited, formal, cultural resource survey done in the area. However, sites in the surrounding areas span the time from the earliest Paleo-Indian occupation through the rural and agricultural development of the early twentieth century. Nearby sites are located in a variety of geographical settings and exhibit a wide range of artifacts and features, but definite trends in site types and changes through time are apparent.

Archaeological evidence indicates that the earliest humans, called the Paleo-Indians, migrated to the region near the close of the Ice Age approximately 12,000 years ago. These people had a highly mobile lifestyle that depended on the hunting of big game including mammoths and the huge, now-extinct bison. The hallmark of most Paleo-Indian sites are the beautiful but deadly spear points that are generally recovered from animal kill and butchering sites and small temporary camps.

There was a gradual but definite shift in the pattern of human use of the area beginning about 9,000 years ago. The changes were due to a combination of climatic fluctuations and an increasing population, coupled with tremendous social change and technological innovation. This stage is referred to as the Archaic and lasted until about 2,000 years ago. Although the Archaic stage is better represented in the archaeological record than the preceding Paleo-Indian stage, the interpretation of the remains is difficult. Evidence of a greater diversity of tools and increased use of native plants is found on many sites but the remains also suggest a more localized and less mobile population.

By 2,000 years ago, the populations of the area became increasingly influenced by the woodland cultures to the east. This period, referred to as the Plains Woodland (1,000–2,000 years ago), brought great changes and innovation including the advent of pottery, the bow and arrow, and semipermanent dwellings. Small villages began to be established and evidence of early agriculture is found along some of the waterways.

Beginning approximately 1,000 years ago until approximately 600 years ago, evidence of an increasingly sedentary population is found at many of the sites. This adaptation is referred to as the Central Plains village tradition and amplifies many of the trends began during the Plains Woodland period. Small villages of earthen structures with associated agricultural fields were more common. The increased use of pottery in conjunction with the construction of food storage pits reflect a population that was spending increasing amounts of time in one location.

Early postcontact occupation of the area (100–400 years ago) included the Pawnee and possibly the Arikara peoples. Their settlements tended to be large villages

with extensive agricultural fields—often located along the major waterways. Bison hunting, fishing, and Euro-American trade were also primary components of the economy. Beginning in the early 1700s, explorers began to make incursions into the area and by the mid-1800s there was a regular stream of emigrants passing through on their way west. Many of these travelers chose to stay and settle in the area referred today as the Rainwater Basin.

### 3.4 SPECIAL MANAGEMENT AREAS

McMurtrey WPA, which lies adjacent to the Meat Animal Research Center, is closed to public use because no public access exists. The WPA contains 1,067 acres (513 wetland acres). Land and vegetation management is the same as for other areas managed by the district. The WPA has two wells and water is pumped during dry years to provide a refuge area for migratory birds.

The district manages two additional areas that are located outside the basin. Schwisow WPA, in Saline County (within the district's boundary), is 61 acres. Haseman WPA is located on the border of Cuming and Dodge counties, outside of the district's boundary. The Farmers Home Administration transferred both properties to the district. Schwisow WPA is primarily riparian habitat along the Little Blue River with some oak-timbered upland. Haseman WPA covers 229.11 acres, with 160 acres of riparian wetland along State Highway 275. Both areas are open to the public uses allowed at the district's other WPAs. The Service is considering these WPAs for property exchange or divestiture for the following reasons: (1) the lands do not meet the purpose of the district, (2) the distance of the WPAs is so far from the district's facilities that staff visits require overnight lodging, and (3) another agency may be able to more efficiently manage the riparian habitats.

### 3.5 VISITOR SERVICES

The district has 60 of the 61 WPAs it manages open to the public; McMurtrey is the only WPA closed to all public use. The Improvement Act outlines wildlife-dependent and wildlife-compatible uses. These uses include hunting, trapping, fishing, wildlife observation, photography, environmental education, and interpretation. Hunting and wildlife observation are the most popular activities, accounting for 80% of annual visitation. (BBC Research and Consulting 2006.)

There are no official visitation data available for the district. All visitation figures reflect best available estimates by district staff. Visitation levels fluctuate between 60,000 and 80,000 visitor days per year, depending on the water level and habitat quality. A typical breakdown of annual visitation by use is about 60% hunting and 40% wildlife viewing. (BBC Research and Consulting 2006.)

All WPAs are marked with boundary signs. Proper signs identify four of the most prominent WPAs in the district. The other WPAs are unmarked and the only locator for these areas is a district brochure map that shows their locations within the district.

### **HUNTING, TRAPPING, AND FISHING**

Visitors generally hunt various waterfowl, pheasant, and deer at the district. The major hunting season for all species is October through December, although there is a light-geese-hunting season in the spring. (BBC Research and Consulting 2006.)

Hunting, trapping, and fishing are allowed in accordance with state regulations. There is a viewing blind at Funk WPA, which is accessible to hunters with disabilities.

The WPAs, other than McMurtrey, are open to fishing. However, the physical characteristics and hydrologic conditions of the wetlands in the WPAs do not provide for a viable fishery.

### **WILDLIFE OBSERVATION AND PHOTOGRAPHY**

The most popular wildlife-viewing season is during spring migration, which lasts from mid-February through April (BBC Research and Consulting 2006).

Every spring and fall, thousands of tourists and locals visit the district to observe and enjoy the semiannual migration of waterfowl and other birds from the southern United States and Mexico to Canada and vice-versa. As a result, ornithologists, amateur birders, hunters, and the general population are witness to a remarkable natural phenomenon. (BBC Consulting 2006.)

There are two viewing blinds, one each at Funk WPA and Massie WPA. Funk WPA has approximately 2 miles of developed trail (constructed in 2006).



*Visitors can view flocks of snow geese (seen in the distance) from this blind at Massie WPA (Clay County). An interpretive sign provides information.*

### **ENVIRONMENTAL EDUCATION**

The district does not provide any public environmental education programs. It is only on rare occasion that district staff is available to conduct a wetland tour.

One reason for that is the long distance between the office and the WPAs. The district's office is located in an industrial park in Kearney, approximately 20 miles from its closest WPA.

### **INTERPRETATION**

There are information kiosks at four WPAs: Mallard Haven, Massie, Funk, and Harvard. Plans for 2007 include additional interpretive signs.

## **3.6 PARTNERSHIPS**

The district is engaged in various partnerships that allow the district to pursue the purposes for the district. While these partnerships help the district to conserve habitats, the district will benefit from expansion in the number and variety of partnerships. Current formal and informal partnerships include those with groups such as the RWBJV, Ducks Unlimited, Pheasants Forever, and The Nature Conservancy.

## **3.7 SOCIOECONOMIC ENVIRONMENT**

A socioeconomic study prepared by BBC Consulting (2006) is the source for information in this section.

The Rainwater Basin supports hunting, trapping, photography, wildlife observation, and environmental education for the public. These recreational activities attract both visitors and dollars from outside the 17-county basin that benefits the local communities. Ancillary visitor activity—such as spending on food, gasoline, and overnight lodging in the local area—provides local businesses with supplemental income and increases the local tax base. District management decisions regarding public access, expansion of services, and other district-related operations may affect recreational traffic and thus visitor expenditure levels.

### **POPULATION AND DEMOGRAPHICS**

Nebraska's population in 2006 was estimated at 1,768,331, a 3.4% increase from 2000. The reported population for the basin in 1999 was 201,245. Rural emigration is a concern with 89% of Nebraska's cities having fewer than 3,000 people. Fifty-three of the 93 counties have reported declining populations between 1990 and 2000. The three major cities of Grand Island, Kearney, and Hastings contribute to 48% of basin's 2004 population (U.S. Census Bureau 2000–2007). These cities' populations totaled 157,662 in 2004 (U.S. Census Bureau 2000–2007).

The 17 counties encompassing the basin are currently neither losing nor gaining population. In comparison, the population of Nebraska has increased at a steady pace since 1980 (see figure 26). While the population retention of the 17 counties is better than many rural Midwestern counties, the gradual loss of residents undoubtedly affects the socioeconomic conditions of the area.

Figure 27 illustrates the aging population within the 17 counties. In 1980, about 26% of the population was between 18 and 34 years old, while that same demographic constituted only 22% of the population in 2004. The median age for the 17 counties has increased by about 6 years since 1980, a trend that is found throughout the state.

### EMPLOYMENT

The civilian workforce for the 17 counties within the basin has increased by about 250 workers per year since 2000, and it is predicted to increase by another 1,000 workers to 107,980 by the year 2009. Estimates from the year 2004 calculated the unemployment rate for the 17 counties within the basin at 3.4%. This is on par with the state of Nebraska, whose 2004 estimate calculated the statewide unemployment rate at 3.5%.

Agricultural row cropping is the main economic enterprise throughout the basin. Eighty percent of the land base is cropland. Major crops are corn and soybeans. Other crops that may occur include wheat, milo, sorghum, and alfalfa. Ranching is a secondary enterprise with 10% of the area being grassland. The number of farms per county in the basin range from 345 in Gosper County to 974 in Seward County. The average-sized farm in the basin in 1987 was 506 acres.

While agriculture has been the traditional mainstay of the state’s economy, new industries such as health care and services now employ an increasing portion of the state’s residents. In the 17 counties within the basin, the total number of farms has decreased at a notable rate. In 1964, the 17 counties supported over

16,500 farms. That number of farms declined to just over 9,200 in 2002 (see figure 28).

### BUSINESS CONCENTRATION


The 17 counties within the basin employ a large number of white-collar workers, with over 13% of its total workforce in the office and administrative support occupations. However, the basin also contains a large manufacturing sector, with 12% of its total workforce employed in production occupations. Other high-ranking occupations include sales and related occupations, education training and library occupations, and food preparation. It should be noted that only 6% of the entire workforce is employed in the agricultural industry.

### RAINWATER BASIN WETLAND MANAGEMENT DISTRICT CURRENT CONDITIONS


About 60% of district visitors live in the 17 counties within the basin. Most destination visitors come for the weekend and stay 1–2 nights, usually in Grand Island, Kearney, Hastings, or York.

### Visitor Spending


Off-site spending by visitors helps support local lodging and retail establishments in surrounding towns. Approximately 40% of the district’s visitor days, or about 30,000 visitor days, are from nonlocal visitors. If 50% of these guests spend the night locally in commercial lodging or campgrounds—and, on average, nonlocal visitors spend \$60 per day for lodging, food,



**Figure 26. Populations of Nebraska and the 17 counties in the Rainwater Basin.**  
 (Note: 2004, 2009, and 2014 populations are estimates. Source: U.S. Census Bureau, PCensus, 2006.)



**Figure 27. Age composition within the 17 counties in the Rainwater Basin.**  
 (Source: U.S. Census Bureau, PCensus, 2006.)



**Figure 28. Farm count for the 17 counties in the Rainwater Basin.**  
 (Source: U.S. Department of Agriculture.)

and supplies—the district’s activity spurs about \$900,000 of new annual spending in the regional economy.

### **Other Economic Considerations**

The purchase of additional farmland for conversion into wildlife habitat by the district is sometimes problematic for local governments. National legislation requires the district to receive permission from the governor before purchasing additional lands. The governor, in turn, often seeks the advice of local county commissioners. In the past, county commissioners have expressed concerns about the conversion of local farmland to wildlife habitat because the land is retired from agricultural production and local purchases for fertilizer, grain, and farming supplies are reduced. Conversion from agricultural designation to public ownership reduces tax revenue.

While county commissioners may not always wholeheartedly support public land acquisition, there has never been a formal denial of land acquisition by the Service on record. Local authorities generally balance the modest loss of business activity against the prospect of additional visitors and the business activity associated with wildlife viewing and hunting. A citizen’s broader right to sell his or her personal property to whomever they choose is also a consideration.

To address the issue of lost tax revenues, Congress passed the Refuge Revenue Sharing Act (revised in 1964), which mandated  $\frac{3}{4}$  of 1% of the current appraised value of land bought for use by the Service be paid to the counties in which the land was bought, a form of payments in lieu of taxes (PILT). These payments are designed to help the counties recoup lost tax revenue. The Service finances these reimbursements with revenues generated from permits sold and services rendered on Service-controlled land nationwide (such as a permit sold for the right to drill for oil on a refuge in Texas). Although land values of refuges have risen over time, the annual revenues used to reimburse the counties have not kept pace. As a result, the participants currently only receive approximately 40% of the money they are owed annually by the Service. Increased tourist spending can mitigate some of this shortfall.

## **3.8 OPERATIONS**

The Service achieves the purposes of the district through effective use of available funding, staff, partners, and facilities.

## **STAFF AND FUNDING**

Spring migration is the primary season of aggressive habitat management, which includes extensive water-pumping operations and shoreline habitat enhancement. Off-season management activities include (1) wetland restoration efforts on public and private agricultural lands, and (2) vegetation management using a combination of prescribed fire, grazing, haying, resting, and grassland reseeded. (BBC Research and Consulting 2006.)

Current federal employment at the district is 12 permanent full-time employees and 3 seasonal employees. (BBC Research and Consulting 2006.)

The district had a \$1.8 million budget in fiscal year 2005. The district does not collect any fees for recreational use of its facilities and does not directly generate any basic local revenue. (BBC Research and Consulting 2006.)

## **LAW ENFORCEMENT**

Refuge law enforcement officers are charged with resource protection and provision of visitor and employee safety on Service lands. Resource protection includes protection of wildlife, fish, and plants, as well as cultural resources. Common activities include (1) investigating illegal hunting, arson, theft, vandalism, dumping, and drug-related activities, and (2) answering questions and assisting visitors in the field.

The district has two dual-function refuge officers who spend 25% of their time dealing primarily with resource protection activities.

## **FACILITIES AND EQUIPMENT**

The district’s facilities include the leased headquarters/shop building with an adjacent, fenced, equipment storage area. These facilities do not meet safety standards for personnel and do not provide adequate space to welcome visitors. The headquarters facility is difficult for visitors to find because it is located in an industrial area in the city of Kearney, which is outside the district’s boundary.

The district owns and maintains two old farm sheds that serve as storage facilities. Two, surplus, mobile trailer homes provide volunteer housing. The district maintains 135 parking lots at the WPAs. The parking areas are fenced (0.2 acre) and contain native vegetation. Each fall, these parking areas are mowed to reduce the risk of wildland fire.

# 4 Management Direction

The management direction for the Rainwater Basin Wetland Management District is to manage the district's resources in a holistic manner. There will be a focus on cooperation, coordination, and better exchange of information. An expanded district staff will work with partners to improve the waterfowl production areas across the landscape of the Rainwater Basin. The emphasis will be on adaptive management—as more information is known, management will be changed to improve effects on the environment.


Through integrated restoration, the district will work to restore ecological processes where appropriate and achieve habitat conditions that require reduced management over time. This will be accomplished while recognizing (1) the role of the district in the overall landscape, and (2) the capabilities of its staff and resources to complete the proposed management actions during the next 15 years. A high priority will be to monitor the effects of habitat management practices and to use research results to direct restoration and management. Another priority will be to increase opportunities for wildlife-dependent, compatible public use and visitor services.

## 4.1 GOALS, OBJECTIVES, STRATEGIES, AND RATIONALE

The goals and objectives in this section are the guidance that will enable the district to provide for its purposes and reach its vision. Strategies to carry out the objectives, to reach the district's goals, will provide for resource needs and public use.

- A goal is a descriptive, broad statement of desired future conditions that conveys a purpose, but does not define measurable units.
- An objective is a concise statement of what is to be achieved, how much is to be achieved, when and where it is to be achieved, and who is responsible to achieve it.
- Strategies are ways to achieve an objective.
- Rationale for the objectives includes background information, assumptions, and technical details used to formulate the objective. The rationale provides context to enhance comprehension and facilitate future evaluations.

NOTE: The overall guidance for use of prescribed fire and management of wildland fire is in the description of the fire management program (appendix L).



*Mallard Pair.*

USFWS

### WETLAND GOAL

Restore, enhance, and maintain the hydrology and early successional vegetation conditions essential to the conservation of migratory birds.

### Wetland Objective A

Within 15 years, restore, enhance, and manage the wetland plant composition to achieve a high level of preferred moist-soil and wet-meadow vegetation. Table 6 shows the shift in plant associations.

### Wetland Objective B

During the next 15 years, manage wetland vegetation to create a vegetative mix of species of various heights, with spring (February–April) habitat conditions having 20–50% of the wetland vegetation between 6 and 12 inches tall.

**Table 6. Current, preferred, and achievable plant composition at WPA wetlands.\***

<i>Associations</i>	<i>Current Composition %</i>	<i>Preferred Composition %</i>	<i>Achievable Composition %</i>
Nonnative undesirable vegetation	19	<1	5–10
Native undesirable vegetation	18	10	10–20
Moist-soil community	44	65	55–65
Wet meadow	17	25	20
Trees	2	0	<1% in wetlands
Total	100		

(Source: Drahota et al. 2004.)

\*Although the percentages are the collective total for all the federal wetlands, it is the district's intent that each wetland has a percent composition that falls close to the preferred composition. For example, no one WPA would have most of the trees that exist on district lands. During wet years, open-water areas reduce the acreage of moist soil, native, and nonnative communities; but the percentages should stay within the achievable composition for most communities during normal years.

### Wetland Objective C

During the next 15 years, improve water-pumping capabilities on wetlands that currently have water-pumping facilities; develop water-pumping facilities for 800–1,000 additional wetland acres; and increase pumping potential to allow more opportunities to supplement water during the fall and spring migration periods.

### Wetland Objective D

Within 15 years, acquire from willing sellers fee-title ownership on 10 adjoining portions of wetlands to complete ownership to allow for better management of individual wetlands.

### Wetland Objective E

Within 15 years, acquire permanent protection from willing sellers on 15 additional wetlands within the basin.

### Wetland Objective F

During the next 15 years, develop baseline information on water quality and quantity of inflowing water into WPAs for use in developing desired conditions or standards.

### Wetland Objective G

Through the duration of the CCP, apply prescribed grazing at a rate, timing, and intensity that is appropriate for management needs.

### Wetland Strategies

1. Annually apply grazing, fire, disking, haying and shredding on 35% of the wetland acres to create a vegetative mix of various heights.
2. Develop annual grazing plans that identify the objective and grazing method that will be used at each WPA.
3. Seasonally monitor and review the effects of grazing to determine if the objective is being met or if modifications need to be made.

4. Construct and maintain adequate boundary fences at 80% of the WPAs.
5. Develop livestock watering facilities for at least 10 WPAs to allow intense grazing treatments to reduce invasive plants and establish native plants.
6. Conduct 1,000–3,500 acres of prescribed burning in wetland habitats each year to encourage and promote the plant composition described in table 6.
7. Continue using IPM strategies to reduce noxious weeds and other invasive plants. Besides burning and grazing, use other management practices including disking, haying, flooding, and herbicide application.
8. Work with partners to increase supplemental water-pumping capacity at WPA wetlands, with a desired water depth of 2–18 inches during migration (October–April).
9. Replace antiquated water-pumping facilities with modern, energy-efficient systems.



Gleason WPA (Kearney County).

10. Coordinate with partners to install additional water-pumping facilities and to improve water delivery in a manner that optimizes water distribution at WPAs within existing high-use areas, wetland complexes, and areas currently without water-pumping capability.
11. Delineate all WPA watersheds to determine actual hydrologic effects on each wetland, and assess the cost and feasibility of restoring each wetland.
12. Implement hydrologic improvements on 10 WPAs using one or more of the following practices: install sediment control structures; replace culverts; install water control structures; remove trees; fill water concentration pits; fill drainage ditches and drains; remove sediment and fill material; and construct dikes or berms.
13. Work with partners and private landowners within WPA watersheds to increase water quantity and quality received by implementing 120 (8 per year) of the following practices (RWBJV 1994):
  - Fill water concentration pits
  - Replace culverts
  - Install buffer areas
  - Restore grassland
  - Install sediment control structures
  - Install “Variable Flow Tailwater Recovery Systems”
  - Remove restrictions to natural runoff
  - Remove sediment and fill material
  - Fill drainage ditches and drains
14. Continue to encourage the drainage districts and county governments to abandon existing wetland drainage tiles associated with WPAs.
15. Work with partners to develop a monitoring program to document quality of water entering the wetlands after storm events.
16. Close a WPA to hunting when threatened or endangered species (such as whooping crane) occur at the WPA.

### Wetland Rationale

Each spring, for a short period, a significant portion of the North American waterfowl population uses habitat of the Rainwater Basin. However, compared with historical conditions, the extent, distribution, and quality of remaining wetlands in the basin is reduced. These changes in habitat conditions likely have various adverse effects related to the life cycle requirements of waterfowl. Wetland vegetation and watershed management need to be done at optimal levels to meet the needs of all types of waterbirds. To increase the benefit of the basin’s wetlands for spring migration, it is critical that a diversity of flooded and moist-soil wetland habitat is made available throughout the basin to provide more, natural wetland

foods and reduce the risk of disease (Samuel et al. 2005, Smith et al. 1989, RWBJV 1993). The basin has a history of large mortality associated with avian cholera during spring migration (Gordon 1989, Samuel 1995, Samuel et al. 2005, Smith and Higgins 1990).

It is the historical nature of the basin’s wetlands to provide resting and feeding habitats for prenesting survival and overall annual waterfowl production (Gersib et al. 1989a, LaGrange and Dinsmore 1988). Moist soil plants such as smartweed and barnyard grass are typical early successional plants that respond quickly to disturbed areas, especially on bare soil (Fredrickson and Taylor 1982). Baldassarre and Bolen (2006) stated that the feeding ecology of waterfowl is a complex interaction of nutritional needs, resource availability, habitat quality, and waterfowl behavior. Feeding ecology is further complicated during winter (November–April) when waterfowl are migrating, preparing for production, and facing increased energy demands due to environmental stresses (Kendeigh et al. 1977, Dubovsky and Kaminski 1994, Ballard et al. 2004). Although Nebraska has an abundance of agricultural fields, waste grains lack many nutrients found in natural foods found in wetlands (Baldassarre et al. 1983, Loesch and Kaminski 1989, Krapu et al. 2004, Baldassarre and Bolen 2006).

Reid et al. (1989) found that native or naturally occurring wetland plant seeds are necessary in a duck’s diet to offset the protein and mineral deficiencies in waste grain. Ankney and MacInnes (1978), Krapu (1981), and Ankney and Afton (1988) showed a positive relationship between lipid reserves and clutch size for various waterfowl species. Failure to meet the nutritional need of waterfowl during winter and spring migration may result in reduced recruitment (the addition of members to the overall population). This is often called the “lipid limitation hypothesis” (Ankney and Afton 1988) and is supported by Ankney and Alisauskas (1991) as a limiting factor. Lipids are an efficient form of energy storage and are more efficiently catabolized than protein, causing Petrie and Rogers (2004) to conclude that these advantages alone explain why most studies conclude that ducks rely heavily on stored lipids during reproduction.

Heitmeyer and Fredrickson (1981), later confirmed by Kaminski and Gluesing (1987), first suggested a relationship between winter habitat conditions and duck recruitment in the following breeding season. Raveling and Heitmeyer (1989) linked increases in northern pintail populations to winter habitat conditions. LaGrange and Dinsmore (1988) went further to say those stopover areas close to breeding areas were crucial habitats for female mallards to acquire adequate nutrients. Many other authors have suggested the correlation between wintering and spring migration energetics and their implications during nesting (Krapu 1981, Rohwer 1984, Dubovsky and Kaminski 1994). This suggests that the basin’s wetland habitat is important for prenesting survival and overall annual waterfowl production (Gersib et al. 1989a).

Food production in wetlands can be very impressive in terms of the number of seeds produced and the varieties. In a 1951 study by J.R. Singleton of the east Texas gulf coast, researchers found that salt marsh bulrush produced an average of 300 pounds (dry weight) of seeds per acre per year, and each acre produced about 5 tons of plant corms. Reinecke and others (1989) concluded that seeds provide the greatest biomass of food in moist-soil habitats, but tubers (Taylor and Smith 2005), roots, rhizomes, stems, leaves, and invertebrates can be important (Jorde 1981, Krapu 1981, Heitmeyer 1988, LaGrange 1985, Ballard et al. 2004, Bowyer et al. 2005). Anderson and Smith (1999) found managed moist soil wetlands had four to five times more ducks than unmanaged wetlands.

Reinecke et al. (1989) and Laubhan and Fredrickson (1992) synthesized seed production into metabolizable energy associated with moist soil plants, as well as daily energy requirements, allowing an estimate of duck use-days (DUDs) based on a wetland's vegetative community and seed production potential. Metabolized energy (ME) is described as a measure of available energy to waterfowl from their diet (Miller and Reinecke 1984). Kendeigh et al. (1977) describes ME as the total daily energy intake compared to the total food biomass required to supply energy needed for any individual or population. Wetland bioenergetics can be described as the relationship between seed biomass and gross energy value. The more energy a wetland can provide, the more bioenergetically efficient it is.

The average energy of moist-soil seeds is 2.5 kilocalories per gram (kcal/g). Ducks do not exploit moist-soil seeds in flooded environments when the seed mass is less than 45.1 pounds per acre (50 kilograms per hectare [kg/ha]). The minimum threshold for energy requirements of a 2.4-pound (1.1-kg) duck is 292 kcal/day (Reinecke et al. 1989). Prince (1979) and Reinecke and others (1989) proposed the calculation of DUDs as a desirable means for evaluating waterfowl habitat management. Haukos and Smith (1993) described DUDs as the number of ducks that could survive on a wetland for 1 day based on native seed availability. Cox and Davis (2002) and Fredrickson and Reid (1988) suggested that it takes larger ducks 2–3 days to replenish endogenous (produced within the body) fat reserves at 480 kcal/day in good habitat. For example, a mallard weighing 2.4 pounds (1.1 kg) would need 3 days of feeding, at a rate of 480 kcal/day, to replenish fat reserves following an 8-hour flight (figure 29, duck-use days).

Using Reinecke's energetics figures, a minimally suitable hectare of moist soil would yield 50,600 kcal/acre (equation 1) or 173 DUDs/acre (equation 2).

$$\text{Equation 1: } 45.1 \text{ lbs/acre (50kg/ha)} \times 2.5 \text{ kcal/g} \\ = 50,600 \text{ kcal/acre (125,000 kcal/ha)}$$

$$\text{Equation 2: } \frac{125,000 \text{ kcal/ha}}{292 \text{ kcal/day}} = 173 \text{ DUDs/acre} \\ \quad \quad \quad (428 \text{ DUDs/ha})$$

As energy requirements go up (such as with weather, disturbance, and stress), the number of DUDs a wetland can provide would go down (Fredrickson and Reid 1988). The DUDs would also vary within vegetative stands since metabolizable energy can vary from 1.0 to 3.5 kcal/g, depending on what plant species are present (Sherfy and Kirkpatrick 1998). During periods when ducks need 480 kcal/day (see figure 30), only 260 DUDs/ha would be provided in the above example.

Based on recent findings from J. Drahota (personal communication) and Rabbe et al. (2004), the basin's wetlands that are dominated by moist-soil plant communities support about 1,779 DUDs/ha and wet meadow communities support 575 DUDs/ha (see figure 29). Conversely, undesirable stands such as cattail (115 DUDs/ha), reed canarygrass (102 DUDs/ha), and bulrush (*Scirpus* spp.) (23 DUDs/ha) plant associations cannot support endogenous or exogenous (produced outside the body) nutrient storage during periods of high energy demands. Drahota (2006) found that wetland management during a dry year increased seed production and the moist-soil plant association provided 8,739 DUDs/ha, compared to 3,250 DUDs/ha in a mixed moist-soil/river bulrush stand.


Estimation of duck use-days will improve when estimates of metabolized energy are available for all moist-soil plants that occur in the basin and when average seed production per stand can be estimated for a variety of environmental and management influences.

Native, undesirable plants such as cattail and river bulrush replace the high food-producing plants if a wetland is left undisturbed for a period of years (Reid et al. 1989). The result is a decline in seed production but an increasing amount of shelter and visual barriers for birds. Woody plants such as elm, cottonwood, and green ash can quickly invade a drying mud flat to convert a grassed wetland into a wooded wetland.

Nonnative undesirables such as reed canarygrass and Canada thistle spread quickly and can dominate or quickly turn a wetland into a monotypic stand of vegetation that is less beneficial and unattractive to waterfowl or other waterbirds (Lavergne and Molofsky 2004). Wet meadow species provide a food source but at a lesser degree (Reinecke et al. 1989, Rabbe et al. 2004).


The shallowness of the wetlands and their frequent flooding and drying make the basin's wetlands suitable for moist-soil plants. Moist-soil plants such as smartweed and barnyard grass are the typical early successional plants that respond quickly to disturbance, especially after a disturbance leaves bare mineral soil (Fredrickson and Taylor 1982).

Experience at the WPAs has shown that grazing, fire, haying, disking, and shredding create an interspersed of plant species. Annual district reports beginning in 1964 document the change in plant communities with various management practices used at the WPAs over



**Figure 29. Rainwater Basin duck use-days.**

*(Calculated using metabolizable energy numbers found in Sherfy and Kirkpatrick [1998] and Checkett et al. [2002]; frequency of occurrence results were then used for each vegetation association as described in Drahota et al. [2004] to extrapolate DUDs for each community based on a daily energy requirement of 292 kcal/day.)*



**Figure 30. Time required for replenishment of endogenous fat reserves following an 8-hour migratory move.**

*(For a duck weighing 2.5 pounds. Taken from Fredrickson and Reid [1988].)*

the years. Grazing and fire were absent from the WPAs between 1966 and the late 1970s. The reports describe increasing problems with (1) smooth brome on uplands, (2) reed canarygrass and trees in wetlands, (3) vegetation-choked wetlands, and (4) noxious weed (primarily musk thistle) spread. In the late 1970s, the herbicide Rodeo® was aerially applied to cattail-choked wetlands to create open water. After about 3 years of use, it was discontinued because the cost per acre was too high. In the early 1980s, fire and grazing were used but at a conservative rate, with haying being the primary management practice. The annual reports indicate that wetland and upland improvement occurred but not enough to change plant composition. In the mid-1990s, the district increased grazing on wetlands dominated with reed canarygrass, river bulrush, and cattail. The district found that a combination of fire followed by intense grazing was the most effective management tool for changing monotypic stands of vegetation into a diverse stand of seed-producing, moist-soil plants.

Grazing has been an integral part of the prairie wetland ecosystem. Most techniques of rangeland management were developed with the idea of increasing and sustaining livestock production by decreasing the inherent variability associated with grazing (Fuhlendorf and Engle 2001). However, this is not the approach used at the WPAs. Grazing treatments are used as a vegetation management tool. Wetland grazing can reduce perennial vegetation, increase diversity, and decrease stand density to result in more wildlife use, especially by migratory waterfowl.

Animal stocking rates of two to four animals per acre in river bulrush and cattail marshes reduced the vegetative cover by 25% (USFWS 1983–93). During drought and low-water periods, livestock trampling compacts the soil, which improves the pooling of water, and tills the surface to allow seed germination. High-intensity grazing also stimulates regrowth (Ermacoff 1968). Cattle should be removed from wetlands before August 10 to allow annual plants to produce seed heads (USFWS 1983–93) if moist soil plant communities are desired. Later grazing and multiyear grazing may be needed to reduce the frequency of occurrence of undesirable species before moist-soil plants can grow.

Livestock grazing generates revenue for use to offset the costs of fencing and control of invasive plants at the WPAs. In addition, grazing provides economic benefits to the local community.

Anderson and Smith (1998) suggest intense moist-soil management for existing wetlands should increase overall nutrition available to waterfowl and other wildlife. Noted moist-soil expert, Leigh Fredrickson (St. Louis, MO), confirmed this (personal communication). Fredrickson stated that increasing disturbance will increase seed production, reduce perennials, and reduce the woody component. The number of years after a disturbance occurs will also

decrease the amount of seed produced. Millet seed production postdisturbance was about 50% less after 3 years and an additional 25% less after 4 years. Undesirable species like cattail and reed (*Phragmites* spp.) have a tremendous amount of belowground biomass—cattail is about 2.5 times more than aboveground biomass and reed (*Phragmites* spp.) is about 6.5 times more. The belowground biomass has to be reduced or eliminated to allow desirable species that are in the seed bank an opportunity to grow.

Effective migratory bird habitat must include a complex of habitat types that provide important food and cover resources (Reid et al. 1989). Historically within this area, wildland fire and grazing by free-roaming bison and elk herds kept wetland vegetation in an early successional stage. Today, natural disturbances have to be replaced with planned management including fire, grazing, haying, flooding, and disking. The frequency and intensity of their use depends on management objectives and various factors including vegetative composition, saturation of soils, and hydrologic patterns within the wetland. Weather events usually determine the timing of treatments. However, if production of moist-soil plant seed is desirable during the same growing season, treatments should be completed before August 10. Kantrud (2006) noted that further studies in wetland management need to occur due to the unknown effects of (1) fire suppression, (2) differential grazing regimes, (3) cultivation, (4) mowing, (5) changes in hydrology, (6) siltation, and (7) pesticides.

Kantrud (2006), after reviewing numerous waterfowl studies, surmised that reductions in height and density of tall emergent plants generally increases breeding duck use. In the basin, migratory habitat has been the focus after researchers found that the basin produces relatively few waterfowl annually (Evans et al. 1967). Kantrud also stated that most waterfowl worldwide favor feeding in shallow water areas where tall emergent plants do not block sunlight.

A variety of bird species depends on plants of various heights. Height variations create structure within habitats that can accommodate greater diversity and higher use by wildlife. Research done by Brennan (2006) et al. have shown that ideal waterfowl habitat, especially for ducks, is an interspersed of tall and short vegetation to create a hemi-marsh condition when spring runoff pools in wetlands. Reinecke et al. (1989) found moist-soil impoundments provide an interspersed of open water and vegetation; a diversity of water depths was attractive to various waterfowl species.

Solid stands of tall vegetation that are greater than 12 inches above the water make areas less attractive to waterfowl (Reid et al. 1989). They may provide an abundance of food, but much of it remains unused. Scattered areas of shorter wetland plants or bare, open water increases bird use of an area. Pederson et al. (1989) stated that, during winter, freedom from



Jeff Drahota/USFWS

*White-rumped sandpipers at Johnson WPA (PHELPS County).*

harassment by predators, availability of food, and thermally protected habitats may be critical. This suggests that this type of habitat in early spring, migratory, staging areas could be similarly critical (Jorde et al. 1984). Brennan (2006) found the percent of emergent vegetation to be a positive indicator for waterbirds—intermediate levels of vegetation in wetlands have the highest species richness. The lower end of the range (6 inches) is provided as a guideline; however, most researchers have found that short vegetation increases the attractiveness to a wider range of species and provides broader benefits for these species (such as feeding, loafing, and breeding).

Wetlands that contain the mix of tall and short vegetation require less runoff or pumped water before the wetland becomes suitable for waterfowl use. Gersib et al. (1989a) noted the temporary and seasonal wetlands provided the highest feeding values to waterfowl due to seasonal fluctuations in hydrology that directly affect vegetative growth and seed production. In drier years, this becomes a critical factor in determining how much migration habitat will be available. The open-water areas will attract the birds and provide them better access to the higher food plants in the flooded, emergent-plant areas of these wetlands.

Although the intent is to have most of the wetlands fall within the category described in the objectives, a few wetlands (<20%) need to be managed for the two extremes: (1) those that have taller, denser stands of

vegetation (not attractive to geese), and (2) those that are open or sparsely vegetated (attractive to geese and shorebirds). White and James (1978) found water depth and the presence of emergent vegetation to be important factors associated with niche partitioning for wintering waterfowl species. The open-water wetlands will benefit snow geese use (Traylor 2000, Drahota 2000) and provide hunting opportunities. Past use of this strategy in the basin has shown separation of snow geese populations from other species of waterfowl. This strategy is expected to reduce the potential or extent of avian cholera. Conversely, wetlands with dense stands of emergent vegetation provide winter habitat for resident species such as pheasant (Baxter and Wolfe 1973, Gabbert et al. 1999, Bakker 2003).

Although vegetation management is critical, water management is equally so. The Great Plains GIS office assessed the Rainwater Basin's wetland conditions in spring 2004 (a year drier than normal) and found that only 14% of the original, historical basins provided any habitat. Of that amount, 91% were in private ownership but only provided 55% of the available waterfowl habitat. Those in public ownership represented only 9% and provided 45% of the available waterfowl habitat. The low number of wetlands within the basin makes it critical that as many as possible contain optimal habitat for all types of waterbirds, and it is important that these water areas be distributed throughout the basin. Rainfall is not consistent throughout the area, therefore, multiple

complexes throughout the basin guarantee that there will be some water areas where water pools when scattered rainfall occurs. As birds concentrate, they quickly deplete their food supply and expose the entire population to disease outbreaks.

Pumping water in the fall can provide habitat for early waterfowl migrants and increase invertebrate abundance in the fall. In addition, maintenance of this water through winter into spring will substantially increase invertebrate abundance (Anderson and Smith 2000). Increased invertebrate numbers will benefit female ducks that molt the first week of March (Jorde 1981) during feather replacement. During dry years, northern pintails with higher body mass survived better in wintering areas (Fleskes et al. 2002, Moon and Haukos 2006), suggesting that quality wetland habitats along the fall migration route play a role in winter survival. Overall, increases in suitable habitat in the breeding, migration, and wintering areas potentially correspond with mallard populations (Heitmeyer and Fredrickson 1981, Bergan and Smith 1993). Fall water pumping can benefit other species such as whooping crane (Richert 1999).



*New submersible pump at Harvard WPA (Clay County).*

Pumping water can be a major expense especially when precipitation is limited or when needed to accomplish moist-soil management objectives (Anderson and Smith 1999). Water pumping should deliver 2–18 inches of water to accommodate foraging needs of shorebirds, waterfowl, and other waterbirds (Laubhan et al. 2006). Since 2000, water-pumping costs in the district have averaged about \$14 per acre. Moist-soil management practices that use fall water pumping can swell clay soil, reduce soil cracking, and slow infiltration rates (Anderson and Smith 1999).

The district has 71 wells scattered over 36 WPAs. Only 35 of these have water-pumping capability and occur at 23 WPAs. Those 35 wells have the ability to deliver water to 2,230 wetland acres (approximately 20% of the total). Each year, WPA and WMA wetlands with pumped water provide core, migrational habitat on a

consistent basis. In drier years, these wetland areas need to have water pumped to them to keep the birds from concentrating on a few small reservoirs and the Platte River. Jorde et al. (1983) found that mallards in Nebraska would move to riverine habitat during winter or cold periods. Therefore, pumping water into wetlands can contribute to the overall distribution of waterfowl within south-central Nebraska.

To increase the benefit of wetlands in the basin for spring migration, it is critical that water be available throughout the basin (RWBJV 1993). Since a significant portion of the entire North American waterfowl population passes through the region, it is even more critical that those populations are not placed at risk. Krapu et al. (1995) advises that waterfowl managers in the basin provide favorable conditions by maintaining well-distributed, wetland-roosting habitat. Supplementing water will increase available habitat, provide more natural foods, and reduce risk associated with crowding (Samuel et al. 2005, Smith et al. 1989).

Each water-pumping facility has a different level of efficiency (for example, the cost per acre-foot of water, gallons per minute, and level of maintenance). Less efficient wells can only be used when wet conditions exist and only when a little supplemental water is needed to reach a desirable habitat condition. Submersible, electrical wells require minimal maintenance and can increase the flexibility of the district's water management. For example, they can be turned on during the harshest part of the winter without fear of freezing or damage, allowing more basins to be ready in the event of an early spring migration. Submersibles also reduce the use of high-cost diesel and natural gas.

Pumping water to wetlands is dependent on various factors (RWBJV 1993). The cost of water pumping and the limited funds makes it necessary to focus water pumping to areas where it will have the most effect for the least cost. For some areas, the capability of the well is less than needed to flood the entire wetland. After a portion of the wetland is flooded, percolation and evaporation begins to equal the pumping capacity of the well.

Thirty-eight WPAs need further acquisition to complete ownership of the wetlands. Partially owned wetlands are not being managed to their full potential. Management treatments such as pumping water, prescribed fire, and grazing are limited or absent. Often the adjoining landowner has different uses or interests in their portion of a wetland. Acquisition will greatly expand the number of fully functional wetlands in Service ownership.

The RWBJV has used GIS technology to identify wetlands having the highest migratory bird value (Bishop 2005). That value is based on biological needs of waterfowl and shorebirds and the geophysical condition of the wetlands. This information will help

the district and its partners target the right wetlands for the right conservation strategy—be it acquisition or enhancement of privately owned wetlands (Bishop 2005).

Natural wetland function occurs when there is a balance in the hydrology between the size of the wetland and the watershed. Alterations within the wetland and watershed change that hydrologic balance. This, in turn, changes the plant composition—often to a less preferred habitat community (Gersib et al. 1989a). In fact, Smith (1990) stated that hydrology is the most likely factor influencing plant community composition.

Much of what occurs in a WPA wetland is dependent on what happens hydrologically within its watershed. Working with private landowners not only addresses the district's hydrology objectives, but also assists landowners in meeting their needs. Pits within a watershed capture water before it reaches the wetland. That water is confined to a deep, smaller, artificial wetland that has little value for migratory waterfowl (Gersib et al. 1989a). The Great Plains GIS office inventoried water concentration pits in the district. There were 11,859 pits found, with 627 pits existing within WPA watersheds. Roads and culverts restrict or impound the natural runoff so one portion of the wetland becomes flooded while other portions receive a smaller portion—again affecting the vegetation and amount of surface water in the wetland. Restoration of watershed hydrology should increase the frequency, size, and duration of pooled water.

The Service's Partners for Fish and Wildlife Program assists the district by working with private landowners within the watersheds. Their work adheres to the practices outlined in the "Rainwater Basin Joint Venture Private Lands Program" criteria (appendix M). Using funds and expertise from various partners since 2000, 89 pits have been filled on privately owned property.

It is unknown what the full extent is and effects are of agricultural runoff in the basin's wetlands. Agricultural runoff can carry fertilizers, pesticides, and heavy silt loads to wetlands at the WPAs. Agricultural chemical exposure at WPAs can have two types of effects:

- direct effects—for example, toxic pesticides that affect nontarget species
- indirect effects—for example, habitat quality that is degraded from nutrient enrichment (Dewey 1986)

Gordon et al. (1997) reported some district wetlands had concentrations of mercury, copper, lead, iron, and zinc that exceeded water quality criteria developed by the U.S. Environmental Protection Agency (USEPA) and concluded that high pesticide and fertilizer use in the area was a likely cause. In addition, herbicides (atrazine for corn and glyphosate for soybean acres) and insecticides (organophosphates and pyrethroids) were applied to 93% and 36% of all corn acreages,

respectively (NASS 2004). Many of these chemicals appear to cause serious degeneration and malformation of organs in wildlife, and could interfere with the normal function of hormonal systems in humans.

In aquatic systems, atrazine exposure can adversely affect periphyton (organisms attached to submerged plants that provide food for invertebrates) (Nelson et al. 1999), invertebrates (Dewey 1986, Dodson et al. 1999), and amphibians (Larson et al. 1998, Hayes et al. 2002). For example, Dewey (1986) found atrazine concentrations of 20 micrograms per liter decreased adult insect emergence by 90% and insect diversity by 60%. Leopard frogs in York County, Nebraska, had gonadal dysgenesis (degeneration of the reproductive organs) in 28% of the sample and testicular oocytes (abnormally occurring female eggs in a male) were found in a single male (Hayes et al. 2002). Insecticides frequently recommended for corn pests in Nebraska include the third-generation pyrethroid permethrin and fourth-generation pyrethroids including bifenthrin, tefluthrin, gamma cyhalothrin, lambda cyhalothrin, cyfluthrin, zeta cypermethrin, and esfenvalerate (UNL 2004). Recent studies by Go et al. (1999) and Kim et al. (2004) indicate that certain pyrethroid insecticides, including permethrin, may function as endocrine modulators in both wildlife and humans.

The Natural Agricultural Statistics Service (NASS) reported 95% and 76% of the district planted to corn receive applications of nitrogen and phosphorus, respectively (NASS 2004). In addition, nutrient-rich runoff can cause wetland eutrophication (overenrichment of a waterbody with nutrients, resulting in excessive growth of organisms and depletion of oxygen) and may result in decreased foraging potential for waterfowl (Gaiser and Lang 1998).

Soil erosion and sedimentation, especially during high flows, reduces the storage capacity of wetlands and forces some of the surface water to flood nonwetland areas (increasing percolation rates) (Stutheit et al. 2004). The buildup of sediment also allows less-preferred perennial plants to survive during the hotter, drier summer period (Reid et al. 1989). Sedimentation of only 0.2 inch (0.5 cm) caused a 92% reduction in seedling emergence of hydrophytic plants and a 99.7% reduction in total invertebrate emergence in northern prairie wetlands (Gleason et al. 2003). Roads, terraces, culverts, and tile drains also reduce, delay, or redirect runoff from wetlands.

Livestock runoff is of particular concern for at least 22 WPAs that have concentrated animal-feeding operations (CAFO) within their watersheds (USFWS 2006). Mindy Meade-Vohland (personal communication) suggested more confinement might exist as enclosures that were not detectable using remote sensing techniques. Of these CAFOs, five are major operations (larger than 40 acres in size) and are within the watersheds of Theesen, Jones, Cottonwood, McMurtrey, Funk, Sinninger, and Prairie Dog waterfowl production areas.

Runoff into the WPAs from CAFOs may directly affect beneficial uses (as defined by the Nebraska Department of Environmental Quality [NDEQ]) such as aquatic life, wildlife, agricultural and municipal water supply, and aesthetics. A recent court case involved feedlot discharge into Cottonwood WPA. In that case, the owner was charged with illegal discharge that violated aesthetic standards and exceeded ammonia standards. Nebraska's water quality standard (NDEQ 2006) states the following:

*To be aesthetically acceptable, wetlands shall be free from human-induced pollution which causes: (1) noxious odors; (2) floating, suspended, colloidal, or settleable materials that produce objectionable films, colors, turbidity, or deposits; and (3) the occurrence of undesirable or nuisance aquatic life (e.g., algal blooms). Wetlands shall also be free of junk, refuse, and discarded dead animals.*

In recent years, aesthetic violations are suspected to occur annually in association with four of the five WPAs with CAFOs (Funk, Theesen, Cottonwood, and McMurtrey WPAs). The biological integrity of wetlands is assumed to diminish when aesthetic violations occur.

Pollutants associated with CAFOs include pesticides, trace elements, salts, nutrients, cyanobacteria toxins, bacterial pathogens, hormones, and antibiotics (USEPA 2003). Studies indicate that heavy metals associated with livestock medicine may be above acceptable levels for waterfowl and other waterbirds. Schwarz et al. (2004) found water and sediments from a swine CAFO served as a source for high concentrations of nutrients, antibiotics, hormones, bacterial pathogens, and elemental contaminants (such as selenium, nickel, copper, and zinc). During large storms, contaminant flows can come from flooded wastewater treatment ponds and from runoff on fields fertilized with animal waste (Sharpley et al. 1999).

The district is working with Service contaminant specialists by providing staff time to collect data as outlined in a research proposal entitled "FY07 Environmental Contaminants Program On-refuge Investigations." This work will provide baseline information about water quality coming into and at the WPAs. Results from this study will define what kind of water quality tests should be conducted by the district.

The "Wetland Management District Ditch and Tile Maintenance Policy" in appendix N will apply to existing ditches or tiles that come onto the WPAs where there is no reservation of a drainage easement in the deed.

## UPLAND GOAL

Reestablish and maintain native grassland communities of the Rainwater Basin.

### Upland Objective A

Within 15 years, restore, enhance, and manage the native grassland plant composition to achieve a high level of diversity. The accumulated, current, and desired percentages for Service lands are shown in table 7.

### Upland Objective B

Through the duration of the CCP, apply prescribed grazing at a rate, timing, and intensity that is appropriate for management needs.

### Upland Strategies

1. Use the "Rapid Assessment Vegetation Monitoring System" to document existing vegetation occurring with each plant community or association.
2. Harvest seed from the WPAs or other local, privately owned grasslands. Collect more than 80 species of grasses, forbs, and sedges from seed harvesting.
3. Use the seed mixes to reestablish native grassland at WPAs containing cropland or areas dominated by nonnative grasses.
4. Conduct 200–3,000 acres of prescribed burning in upland habitats each year to encourage and promote the plant composition shown in table 7.
5. Continue use of IPM strategies to reduce noxious weeds and other invasive plants.
6. Continue to remove trees to create an open grassland.
7. Develop annual grazing plans that identify the objective and grazing method that will be used at each WPA.
8. Seasonally monitor and review the effects of grazing and prescribed fires to determine if the objective is being met or if modifications need to be made.
9. Construct and maintain adequate boundary fences at 80% of the WPAs.
10. Develop livestock watering facilities for at least 10 WPAs to allow intense grazing treatments to reduce invasive plants and establish native plants.

### Upland Rationale

Grassland plays a vital role in buffering runoff and in providing feeding, nesting, and shelter habitat for migrating and residential wildlife. Within the Rainwater Basin, agriculture and roads have replaced the tall-grass prairie in the eastern portion and the mid-grass prairie in the western portion of the basin. Steinauer and Rolfsmeier (2003) reported that more than 97% of tall-grass prairie that once covered the

**Table 7. Current, preferred, and achievable plant composition at WPA uplands.**

<i>Associations</i>	<i>Current Composition %</i>	<i>Preferred Composition %</i>	<i>Achievable Composition %</i>
Native grassland	64	>95	>80
Invasive grassland	27	<5	<19
Cropland	4	0	0
Trees*	5	<1	<1
Total	100		

(Source: Drahota et al. 2004.)

\*Although the percentages are the collective total for all the WPA uplands, it is the district's intent that each upland unit has a percent composition that falls close to the preferred composition. For example, no single WPA would have more than 1% of the trees that exist on district lands. The native grassland association includes all grasses and forbs that historically occurred within the basin (refer to table 4 and appendix H for a list of plant associations that are recorded during transect sampling). Transect data collection determines the frequency of occurrence for all vegetative associations that occur in the stand. Dominant plant communities are determined by data analysis. Plant associations that have the most occurrences within the sample area are considered dominant.

eastern one-third of Nebraska has been lost. Over the entire basin, less than 10% of the original grassland remains (personal communication with Ryan Reker, RWBJV, Grand Island, NE). Nearly all of the remaining grassland has been significantly altered by land use that promotes invasive cool-season plants such as smooth brome and Kentucky bluegrass.



USFWS



USFWS

District staff use prescribed fire as a tool to manage uplands at the waterfowl production areas.

Grassland birds have experienced dramatic declines because of the loss of grasslands. The North American Breeding Bird Survey reports that 70% of the 29 species characteristic of North American prairies have experienced a decline in population. A portion of that decline is attributed to the small acreage of remaining grassland parcels and the increasing number of trees found within the grasslands (Bakker 2003). Cowbird parasitism is especially concerning (Bakker 2003) in the district due to planted shelterbelts and scattered volunteer trees that are numerous at the WPAs.

It is within the directive of the Service to manage areas according to their historical conditions for the benefit of multiple species of plants, animals, and insects (see habitat requirements in table 8). Because of the small amount of remaining native grassland, it is important to manage the uplands in this manner.

For most of the grassland bird species, the percentage of woody cover should be less than 5% of the plant community (McKee et al. 1998). Prairie chickens prefer less than 1% woody vegetation for lek sites (Merrill et al. 1999). McCarthy et al. (1997) found woody cover encroachment directly decreased adequate nesting cover for prairie chickens.

Burger et al. (1994) found prairie fragmentation directly affected predation rates on bobwhite quail, noting that nests found more than 60 meters from woody cover were three times more successful than those found less than 60 meters from woody cover. Therefore, tree and shrub removal is critically important for those WPAs that have a high percentage of trees or shrubs. Bakker (2000) recommends removing woodland habitat within or adjacent to grassland and acquiring or preserving grassland patches large enough (300–600 acres) to attract the majority of grassland-dependent species.

Native grassland responds better to natural ecological processes (including drought), which provides a more stable habitat to meet wildlife needs. It provides for a greater diversity of plants and animals.

Since the district reseeds areas with a high-diversity (80+ species), grassland seed mix, it is assumed that the species diversity of the established grassland will be greater than 50 species. High-diversity grassland is important for most of the native-grassland bird species (Bakker 2000). Native grassland management will benefit all of the species listed in table 9 (Bakker 2003); however, each has specific habitat requirements that are not the same. Grassland management treatments should provide a variety of grassland

conditions that include short-vegetation grazed or hayed areas, medium-height vegetation with low thatch, tall vegetation with low thatch, and grass stands with heavy thatch (Bakker 2003, Sporrang 2001). High-diversity grassland can also be an important line of defense against invasive plant species. Kennedy et al. (2002) found that restoration and revegetation practices that included high-diversity seeding proved effective for exclusion of undesirable invaders.

**Table 8. Habitat requirements for selected grassland birds.**

<i>Species</i>	<i>Vegetation Height (inches)</i>	<i>Litter (inches)</i>	<i>Patch Size (acres)</i>	<i>Distance from Trees (feet)</i>
bobolink	10–18	1.3–3.6	100	150
burrowing owl	<5	minimal	10	>328
dickeissel	8–40	0.6	25	prevent woody encroachment
long-billed curlew	<12	minimal	104	avoids areas with high-density trees and shrubs
grasshopper sparrow	8–24	not available	20	164
sharp-tailed grouse	6–16	use areas that are idle for several years	150	>164
short-eared owl	12–24	2–8 years of residual cover	183	not available
upland sandpiper	1–24	1.0	250	328

(Sources: Grant 1965, Wiens 1973, Clark 1975, Duebber and Lokemeon 1977, Redmond et al. 1981, Johnsgard 1983, Prose 1987, Renken and Dinsmore 1987, Messmer 1990, Haug et al. 1993, Herkert et al. 1993, Pampush and Anthony 1993, Helzer 1996, Hughes 1996, Madden 1996, Connelly et al. 1998, Clayton and Schmutz 1999, Helzer and Jelinski 1999, Dugger and Dugger 2002, and Laubhan et al. 2005.)

**Table 9. Key grassland species found in the Rainwater Basin.**

<i>Grassland-nesting Species</i>	<i>Partners in Flight (score is greater than 20 points)</i>	<i>USFWS Focal Species (2005)</i>	<i>BCC Bird Conservation Region 19* (2002)</i>	<i>Nebraska Tier 1 Species</i>
northern harrier	✓	—	✓	—
Swainson's hawk	✓	—	✓	—
upland sandpiper	✓	✓	—	✓
burrowing owl	—	✓	—	✓
short-eared owl	—	✓	—	✓
sedge wren	—	✓	—	—
lark bunting	✓	—	—	—
grasshopper sparrow	✓	✓	—	—
dickeissel	✓	—	—	—
bobolink	—	✓	—	—
eastern meadowlark	—	✓	—	—


(Source: Sharpe et al. 2001.)

\*Designated by Birds of Conservation Concern, 2002; region 19 comprises central Nebraska, Kansas, Oklahoma, and Texas.



District staff mix 31,000 pounds of seed harvested during the summer. More than 100 species were collected.

Game birds also benefit from managed native grassland stands. Bakker (2003) noted that pheasant researchers found nesting success to increase with increasing distance from trees. Nesting success was lower in and near shelterbelts in South Dakota and Colorado (Trautman et al. 1959, Olson and Flake 1975, Snyder 1984). Heavy predation rates near shelterbelts, road ditches, and fencerows prevented successful nesting (Trautman et al. 1959). Gabbert et al. (1999) found that predation was significantly higher than severe winter mortality—suggesting that thermal cover provided by native grass stands will provide the highest winter survival rates for pheasants. Managing natural areas for grassland bird species involves providing the nesting habitat requirements and food resources essential for production and survival. These



Greater Prairie-chicken.

requirements include large, treeless patches that contain diverse vegetative structure (Renken and Dinsmore 1987, Johnson and Temple 1990, Volkert 1992, Helzer and Jelinski 1999, DeJong 2001, Herkert et al. 2003, Davis 2004, Fritcher et al. 2004). Management practices that favor grassland-nesting birds will benefit the grassland species of management concern shown in table 9 (Sharpe et al. 2001).

Grazing has been an integral part of the prairie ecosystem. Most techniques of rangeland management were developed with the idea of increasing and sustaining livestock production by decreasing the inherent variability associated with grazing (Fuhlendorf and Engle 2001). However, this is not the approach used at the WPAs. Grazing treatments are used as a vegetation management tool. Proper upland grazing can reduce undesirable species, maintain healthy grasslands, and promote heterogeneity (Fuhlendorf and Engle 2001).

Livestock grazing generates revenue for use to offset the costs of fencing and control of invasive plants at the WPAs. In addition, grazing provides economic benefits to the local community.

The district has mapped the uplands at the WPAs but not to the level of detail or accuracy needed to assess the status or future changes in uplands over a period of years. Currently, the district is not staffed to accurately measure the effect management tools, such as fire and grazing, is having on grassland communities. Measurements from vegetation transects will provide that information. Measurements taken during vegetation transect sampling will provide enough information to adequately address upland-monitoring needs. Without this data, it will be difficult to determine whether upland objectives have been met.

## **WATER RIGHTS GOAL**

Develop partnerships to protect the natural hydrology of WPA watersheds and ensure the necessary water rights are in place to protect future use of both ground and surface water.

### **Water Rights Objective A**

Within 2 years of CCP approval, complete an inventory of existing district-owned water rights and monitor changing natural resource district regulations associated with groundwater use.

### **Water Rights Objective B**

Through the duration of the CCP, work closely with partners to obtain all necessary water rights and to protect the integrity and hydrology of district wetlands.

### **Water Rights Strategies**

1. Work with water rights experts at the regional office to develop and perform a formal review and determination of the legal status of existing water rights.

2. Work with partners to identify threats and possible solutions to the loss of surface water runoff to district wetlands.

### **Water Rights Rationale**

State law entitles groundwater users to “reasonable and beneficial use of the groundwater.” The increase of groundwater irrigation and drought conditions has caused a decline in groundwater throughout the district. The Nebraska Ground Water Management and Protection Act, amended in 2004, requires the Nebraska Department of Natural Resources to annually determine which river basins, subbasins, or reaches are considered fully appropriated. Fully appropriated basins will cause the respective natural resource district to (1) place a moratorium on new surface and groundwater uses, and (2) develop an integrated surface water and groundwater management plan. Only WPAs located in the Tri-Basin Natural Resource District are in fully or overappropriated basins. The Upper Big Blue Natural Resource District established the goal of holding its average groundwater level above the 1978 level. If the average groundwater level drops below a level 3 feet above the 1978 average level, groundwater users will be required to report annual groundwater use.

It is uncertain what future actions will be taken by the natural resource districts or what effect those actions may have on future use of groundwater to supplement wetlands. With water becoming scarcer, there is concern that neighboring lands will capture natural runoff before it reaches the district’s wetlands. It is not clear if any legislation or regulations are in place to protect public wetlands from being dried up by diversion of surface water runoff.

### **INVASIVE PLANT SPECIES GOAL**

Reduce and control the spread of nondesirable, nonnative plant species within wetland and upland habitats for the benefit of native plant and wildlife communities.

#### **Invasive Plant Species Objective A**

Throughout the life of the CCP, continue to monitor and control invasive plant species in wetland and upland habitats by using the appropriate treatment for each situation.

#### **Invasive Plant Species Strategies**

1. Annually map and document treatment of nonpreferred plant communities throughout the district.
2. Develop an integrated pest management approach that will include chemical, biological (such as insects), mechanical, and physical (such as fire, grazing) treatments.
3. Treat known stands of state-identified noxious weeds. Other species that degrade wetland and upland habitats will be second priority.
4. Establish healthy stands of preferred, native plants that can compete with invasive plants.
5. Develop partnerships that will find new ways to efficiently control invasive plant species by combining resources of all partners.

### **Invasive Plant Species Rationale**

Invasive, nonnative wetland and upland plants are a serious problem affecting thousands of acres of wildlife habitat along the Platte River and within the basin. Nonnative plants such as Canada thistle and reed canarygrass can outcompete with native flora—creating a monotypic stands if left undisturbed. Native species, such as cattail and river bulrush can do the same if left unmanaged.

In 2004 7,596 acres of undesirable plant communities (includes noxious weeds, cattail, reed canarygrass, bulrush, and invasive cool-season grasses) were mapped at the WPAs.

Vegetation management is key to providing optimal wetland and upland habitat for both migratory and resident wildlife. Healthy native plant communities are better equipped to withstand weed invasions (Kennedy et al. 2002). Long-term control requires the cooperation of public and private land managers throughout the basin. A joint effort by all partners is needed to conduct research on finding the best management practices to control or eliminate individual species.

### **WILDLIFE DISEASES GOAL**

Work with partners to prevent or control the outbreak and spread of wildlife-borne diseases to protect human and migratory bird populations.

#### **Wildlife Diseases Objective A**

Through the duration of the CCP, continue to monitor WPAs for outbreaks of various wildlife diseases, especially avian cholera and influenza.

#### **Wildlife Diseases Objective B**

Respond in an appropriate manner to contain any disease outbreak that occurs.

#### **Wildlife Diseases Strategies**

1. Work closely with the NGPC and other state and federal specialists to monitor and control all wildlife diseases at the WPAs.
2. Maintain an up-to-date disease contingency plan.
3. Follow federal and state guidelines for monitoring and control of wildlife diseases.
4. Use partnerships to increase awareness and preparedness for the monitoring, detection, and control of wildlife diseases.
5. Where possible, use management practices such as supplemental water pumping to reduce the spread or effect of disease.

## Wildlife Diseases Rationale

Avian cholera was first documented in the district in 1975. It has occurred in the district every year since. The level of outbreak fluctuates from year to year. Cholera mortality in 1998 was estimated at more than 100,000 birds, primarily snow geese, while the mortality in the past 5 years has been in the low hundreds. It is not known for certainty what environmental or physiological factors trigger an outbreak, but it appears to be associated with physiologically stressed birds that are concentrated on a limited number of wetlands (Smith and Higgins 1990). Avian cholera epizootics (diseases affecting large numbers of animals) were found to be inversely related to densities of semipermanent wetland basins. Avian cholera is widely distributed and poses a constant threat to migratory bird populations, especially where dense concentrations of birds occur.

There is a growing concern that an avian influenza pandemic could occur within the next few years. What remains unknown is the possibility that other diseases could reach outbreak proportions while birds are concentrated in the district. The best approach to take is to be vigilant and prepared.

### RESEARCH AND SCIENCE GOAL

Encourage and support research that substantially contributes to the understanding and management of the Rainwater Basin wetland and grassland ecosystem.

#### Research Objective A

Through the duration of the CCP, support research that furthers the understanding of the ecology, wildlife populations, socioeconomics, and hydrology within the Rainwater Basin.

### Research and Science Strategies

1. Complete baseline research that determines the watershed boundaries for the WPAs and the hydrologic events that affect wetlands. Determine what practices should be done to restore wetland hydrology.
2. Conduct an in-depth inventory of invertebrates, amphibians, reptiles, birds, and small mammals within the district.
3. Determine the baseline composition for each upland plant community using the “Rapid Assessment Vegetation Monitoring System.”
4. Determine the socioeconomic effects associated with Service-owned lands: (1) property tax deficiencies, (2) increased local revenue generated from recreational opportunities provided by public access and uses, and (3) aesthetic values.
5. Conduct an invertebrate study to assess the effects of land and water management actions.
6. Identify the energetics contribution of the district to waterfowl and other migratory birds.



*A crew of district staff, Bureau of Reclamation employees, and summer interns map vegetation at all the waterfowl production areas.*

7. Conduct or support contaminant research associated with nonpoint and point source pollution.
8. Work with partners to continue to identify needed research, obtain funding, and support the research process.
9. Conduct district data collection to support ongoing research.

### Research and Science Rationale

Smith (1998) identified the current district research needs. The district is working with the RWBJV to accomplish those needs; however, district staff and resources are limited. Research done within the district can be divided into six categories: wetland inventory, bird inventory, habitat evaluation, bird biology, avian cholera, and hydrology.

Most of the early research involved wetland inventory. The first research assessed Clay and Fillmore counties (USFWS 1954). In the late 1960s, a more detailed inventory used soil data and field surveys to assess how many large wetlands had been drained or degraded (McMurtrey et al. 1972). Schildman and Hurt (1984) updated the McMurtrey data and found that 10% of the original wetlands and 22% of the original wetland acres remained. Raines et al. (1990) included small wetlands in their review; they noted the declining number of basins and that the frequency of degraded wetlands was increasing. An assessment in 2004 of land use in the district reported 14% of the historical wetlands have some wetland function (pooled water). Of that amount, 49% are classified as cropped wetlands (Bishop and Reker 2006).

Bird inventories started in the late 1950s with the Service counting numbers of greater white-fronted geese. Those counts were used through 1992 as a population index for the midcontinental population (Benning 1987, Solberg 1992). The district conducted aerial waterfowl surveys during spring migration (USFWS 1983–1993). The NGPC inventoried the basin for species of concern to assess the potential

conflicts that may occur with the “Conservation Order for Mid-continent Light Geese” (COMLG) (NGPC 1997–1999). The Canadian Wildlife Service conducted two goose counts in 1999 to compare population numbers using the district with previous population estimates (Warner and Nieman 1999). In addition, the NGPC inventoried spring waterfowl numbers during a 4-year study that looked at effects on nontarget species during the COMLG (NGPC 2000–2003). Austin and Richert (2001) mapped and evaluated whooping crane habitat selection. Although less is known about populations of nonwaterfowl species, the wetlands provide habitat for a minimum of 200,000–300,000 shorebirds represented by over 30 species (LaGrange 2005). Jorgensen (2004) was able to summarize shorebird use in the district. Jorgensen (personal communication) is currently completing buff-breasted sandpiper research. Records of birds occasionally observed in the basin are found in “The Birds of Nebraska and Adjacent Plains States” (Johnsgard 1997).

Habitat evaluations in the basin are limited. Taylor et al. (1978) correlated landscape changes with pheasant numbers. Gersib et al. (1989a) looked at waterfowl densities and activity time-budgets for the basin’s wetlands; they found that temporary and seasonal wetlands were the most valuable for spring-staging waterfowl. Gilbert (1989) sampled 47 wetlands in the basin and organized its plant communities in accordance with the dominant hydric soils that were present. Gersib et al. (1989b) also completed a functional assessment concluding that the basin’s wetlands have a high probability of providing wildlife habitat, food chain support, long-term and seasonal nutrient retention, flood storage, and sediment trapping. The NGPC (1977–1999), the Service (USFWS 1977–1999), and Richert (1999) have documented whooping crane roost sites and habitat preferences for central Nebraska. Stutheit et al. (2004) provided a comprehensive review of wetland hydrology and function in the basin. Drahota et al. (2004) mapped and sampled vegetation communities at public areas. Brennan (2006) assessed local and landscape factors influencing migratory bird use.

Duck production research by Evans and Wolfe (1967) found the district’s wetlands to be intermittent, producing 10,000 birds to flight stage annually. Their findings led managers to focus more on managing for spring migration habitat rather than nesting habitat. Krapu et al. (1995) looked at habitat, food, and nutrients in white-fronted geese and concluded that fat and protein contents increased for females during spring staging in the district. Cox (1998) looked at weight gain, nutrient reserves, and habitat use by white-fronted goose, snow goose, and northern pintail. Cox and Davis (2002) used telemetry techniques to assess northern pintail habitat use, movements, and survival during spring migration and found the smaller wetlands in the basin to be the most important habitat. Farmer and Parent (1997) found that the distance

between wetlands influenced use by the pectoral sandpiper. Thus, higher wetland densities provided greater variability in food resources to maximize foraging opportunities and minimize energy expenditures (Farmer and Wiens 1999). Max Post van der Burg (2005) looked at factors affecting songbird nest survival and brood size.

Numerous avian cholera studies have been completed in the district. The first avian cholera outbreak in the district was reported in 1975 (Zinkl et al. 1977). Research conducted in the 1980s and ‘90s had little success pinpointing the cause of outbreaks and could not develop any strategies that minimized bird loss during outbreaks (Windingstad et al. 1984, Smith et al. 1989, Smith and Higgins 1990, Smith et al. 1990). However, it was determined that the bacteria can remain in the environment for several days after an outbreak occurs (Price and Brand 1984), which potentially jeopardizes the next migrants that stop at the wetland. Samuel (1995) listed those factors that can affect cholera outbreaks and survival of the bacterium in the environment. Cox (1999) found no correlation between a bird’s body condition or size and its susceptibility to cholera.

Wetland hydrology appears to be the newest frontier for research in the basin. It is clear that the cumulative hydrologic impact within the hydric footprint and within the watershed can affect pooling duration, frequency, and flooded acres. Wetlands in the basin have the potential to recharge depleted groundwater resources—soil profiles support this due to the lack of visible chloride deposits in dry wetlands (personal communication with Warren Wood, Michigan State University).

## **CULTURAL RESOURCES GOAL**

Identify and evaluate the cultural resources in the district and protect those that are determined to be significant.

### **Cultural Resources Objective A**

Within 10 years of CCP approval, complete a comprehensive cultural resources survey and overview that identifies sensitive areas and helps to preserve historic records and information within the district to ensure protection of cultural resources and compliance with state and federal cultural resources protection laws.

### **Cultural Resources Rationale**

Protecting significant sites, buildings, structures, and objects is the primary goal of the cultural resource work in the district. Cultural resources include the material evidence of past human activities: prehistoric, historic, and architectural in addition to any traditional cultural properties that may or may not have material evidence. A resource is considered significant if it is listed, eligible, or potentially eligible for the National Register of Historic Places.

Protection of significant cultural resources is primarily accomplished through compliance with section 106 of the National Historic Preservation Act. Any project that has the potential to affect structures older than 50 years or cause ground disturbance should be evaluated for its potential to impact cultural resources. Cultural resource personnel should be notified early in the planning process so that alterations of plans can be made if necessary and delays can be avoided.

Surveys are the best tool available to determine the location of cultural resources on the district. Through surveys, both historic and prehistoric sites are identified and key information is gathered that helps for planning, research, and educational outreach. Although small surveys have been done, usually as a part of the section 106 process, large-scale surveys are needed to better understand the distribution and nature of the resources.

A cultural resource overview is needed for the district. This comprehensive study will describe the nature and extent of past cultural resource investigations, the types of resources known on the district, and the interpretive context for these resources. The document will outline specific threats to the resources and the ability of future work to address regional research questions. It will also serve as a planning tool to help encourage consideration of cultural resources during project planning.

Long-term and past employees, in addition to local residents and members of regional historical societies, can be a wealth of information concerning the history of the district and the location of specific resources. District staffs, especially maintenance personnel, often remember alterations to historic structures and know the location of unrecorded archaeological resources.

The extent and condition of historical records, maps, artifacts, and photographs at the district is unknown. This type of historical documentation can provide valuable insight into the development and changes at the district through time. A comprehensive inventory of these items is needed.

### **Cultural Resources Strategies**

1. Notify state and federal cultural resources personnel (for example, the State Historic Preservation Office) early in planning processes so alterations of plans can be made if necessary and delays can be avoided.
2. Notify all district staff of known cultural resource locations to facilitate management and protect the resource. Identify district areas that have not been surveyed but have a high potential for cultural resources.
3. Notify cultural resource staffs (state and federal) when previously unrecorded cultural resources are found in the district.
4. Develop partnerships and work with state and federal cultural resource staffs to develop a

comprehensive inventory and compilation of the cultural resources within the district.

5. Conduct cultural resource-related interviews with district staff and local residents.
6. Locate individuals with knowledge on the general history, location of sites, or alterations to various buildings and structures within the district and document this information to preserve historic records.
7. Obtain assistance from state and federal cultural resource staffs, as well as from state universities and private organizations, to carry out an analysis as to how to best stabilize and store cultural resource items for future reference and educational purposes.

### **VISITOR SERVICES GOAL**

Provide quality wildlife-dependent recreation and educational opportunities by instilling an understanding of basic ecological processes, purpose of the Rainwater Basin Wetland Management District, and mission of the Service for persons of all abilities and cultural backgrounds.

#### **Visitor Services Objective A**

Over the next 15 years, continue to provide safe and quality hunting and trapping opportunities at WPAs.

#### **Visitor Services Objective B**

Within 5 years, fund and hire a full-time outdoor recreation planner (ORP) to develop demographic, attitude, and expectation profiles of wildlife-dependent recreational users, which will determine a long-term plan for providing quality public use opportunities.

#### **Visitor Services Objective C**

Through the duration of the CCP, work with partners to provide demonstrations, written information, and other methods of communication that inform the public about the benefits of management actions and increase and improve education, outreach, and recreational opportunities within the district. Development will be guided by the arrival of the proposed ORP and the creation of a future visitor services plan.

### **Visitor Services Strategies**

1. Construct and maintain at least one additional handicap-accessible blind at a WPA in the eastern portion of the district.
2. Maintain parking areas and access points to meet visitor needs.
3. Continue to construct and maintain adequate parking facilities on at least 95% of the WPAs.
4. Increase the amount of signage (such as boundary, regulation, and directional) at the WPAs.
5. Post 50% of the WPAs with entrance signs that include the WPA's name.

6. Provide adequate law enforcement coverage of all hunting and trapping seasons to ensure compliance with laws and regulations while providing for public safety and welfare.
7. Establish mechanisms to work collaboratively with USGS–BRD economists, state universities' departments of agriculture and resource economics, other agencies, national and worldwide travel agencies, outfitter groups, and nongovernmental organizations to obtain the necessary data to determine travel trends to the district.
8. Work with USGS–BRD economists and state universities' departments of agriculture and resource economics to better understand the values and needs of local, national, and international visitors to the district.
9. Within 5 years of designation and funding of a full-time ORP, expand the quality and quantity of wildlife- and habitat-oriented events and programs within the district.
10. Develop a visitor services plan.
11. Designate five WPAs that will become focus areas, representing other WPAs in the district. An ORP position will facilitate integration of environmental education and interpretation for counties in the district.
12. Within 5 years of designation and funding of a full-time ORP, install kiosks that have interchangeable interpretive panels at focus WPAs.
13. Develop a watchable wildlife brochure that identifies the district's WPAs and state areas, as well as seasons that offer exceptional wildlife observation and photography opportunities.
14. Develop a partnership with Nebraska's visitors bureau and other similar agencies to develop and include in their existing publication and websites information about the district and wildlife observation and photography opportunities.
15. Within 3 years of developing an outreach partnership, create an outreach plan that increases the awareness of the district's assets both within and outside the Service.

### Visitor Services Rationale

The WPAs are open to the public for hunting, fishing, and trapping during legal seasons. Photography, environmental education, and interpretation are allowed year-round during daylight hours.

The COMLG, approved in 1999, allows for light-geese-hunting during the spring migration. To provide refuge areas during the migration, some public areas are closed to hunting and the entire district is on a day-closure schedule (hunting allowed Saturday, Sunday, Wednesday, and Thursday). Closed WPAs include Bluestem, Clark, Eckhardt, Funk, Hultine, Lindau, Massie, Nelson, Prairie Dog, Springer, Verona, and



*A district intern assists a young bird watcher.*

USFWS

Wilkins. Closure provides safe haven for waterfowl during periods of significant public use (Delnicki and Reinecke 1986). Hunting pressure and disturbance often discourage waterfowl use on preferred wetlands (Jessen 1970, Raveling et al. 1972, Koerner et al. 1974, Raveling 1978).

The district staff does not have training or responsibilities directed toward education, outreach, or public use in general. The district has always hired personnel with expertise in managing wetlands and grassland. Nearly all the work done by the district to encourage public use has focused on hunting. That work has mostly been limited to placement of boundary signs and maintenance of parking lots. Although spring and fall water pumping is done for the health of waterfowl populations, it also increases hunting opportunities.

In the last decade, informational kiosks have been constructed at 4 of the 59 WPAs. Funk WPA has a viewing and hunting blind that is accessible to people with disabilities, and there are interpretive signs. At Massie WPA, the neighboring community of Clay Center has collaborated with the district to construct and maintain an observation blind.

Although the district periodically gets requests from schools and Scout groups to provide environmental education programs, the district often declines because of lack of staff, expertise, and materials.

## **PARTNERSHIP GOAL**

Promote and develop partnerships with adjacent landowners, public and private organizations, Native American tribes, and other interested individuals to protect, restore, enhance, and maintain a diverse and productive ecosystem.

### **Partnership Objective A**

Continue to provide strong support and active participation in the RWBJV partnerships to promote activities and projects that are mutually beneficial.

### **Partnership Objective B**

Develop more community-based partnerships that involve local individuals, groups, or organizations in the protection, management, enhancement, and enjoyment of the basin's wetlands.

### **Partnership Strategies**

1. Provide representation on joint venture work groups and committees such as the private lands work group, acquisition work group, public lands work group, and technical committee.
2. Set priorities for Service funding and support for projects (land protection, staff, and equipment) that accomplish district objectives and use partner contributions.
3. Work with NGPC to more efficiently manage public lands that are near each other through coordinated exchange of staff, cooperators, equipment, and facilities.
4. Pursue partnerships to develop a field bird guide that is specific to the basin.
5. Pursue partnership funding for an ORP.
6. Develop a list of high-priority and innovative projects that overlap between district and other joint venture partner needs.
7. Foster a working relationship with individual producers to enhance and maintain habitat conditions at the WPAs.
8. Develop, coordinate, and maintain working relationships with state and local law enforcement authorities and fire departments to protect district properties and trust species.
9. Develop, coordinate, and maintain working relationships with cooperating agencies and joint venture partners who conduct prescribed burns.
10. Through the Partners for Fish and Wildlife Program and other partners, develop, coordinate, and maintain working relationships with joint venture partners who also deliver private lands projects.

### **Partnership Rationale**

The basin has a mix of wetland types. Because of their large size and water permanence, some wetlands

are best for public ownership. The other wetlands will remain in private ownership and will require a partnership approach to restore or enhance them. The value of each WPA for waterfowl is dependent on its proximity to a complex of other wetlands. A WPA that is adjacent to other wetlands will be of more value to waterfowl than one that is isolated from other wetlands. It is important that the district work with its neighbors and conservation partners to improve the basin-wide landscape for the benefit of migratory birds, other wildlife, and the human environment.

Working together has been characteristic for the neighbors, agencies, and organizations in the Rainwater Basin. Partnerships have flourished because each group recognizes that it cannot “do it” alone. The RWBJV (described in chapter 1, section 1.4) is a partnership organization made up of government and conservation organizations, as well as landowners.

The RWBJV's goal is to

*Restore and maintain sufficient wetland habitat in the Rainwater Basin area of Nebraska to assist in meeting population objectives identified in the North American Waterfowl Management Plan.*

The joint venture's objectives are in line with those of the district. The RWBJV commonly joins with nontraditional partners to (1) restore and protect additional wetlands, (2) provide reliable water to at least one-third of the protected wetlands, and (3) enhance existing wetlands.

Ducks Unlimited has identified the Rainwater Basin as a conservation priority for their organization. They are actively involved in wetland restoration and acquisition. Much of the restoration work done by Ducks Unlimited has occurred on NGPC and district lands. In recent



*This combine used to harvest native grasses was purchased through a three-way partnership—the district, NGPC, and Pheasants Forever.*

years, they have acquired lands that are being restored and planned to be transferred to NGPC or district ownership. The U.S. Department of Agriculture's (USDA) Wetland Reserve Program has restored and protected wetlands throughout the basin. The Nature Conservancy has helped protect 1,765 acres that are currently under the Service's management. Another 466 acres are under the management of the NGPC.

## **SOCIOECONOMICS GOAL**

Obtain a better understanding of the social and economic contribution WPAs make to the people and communities within the Rainwater Basin.

### **Socioeconomics Objective A**

Develop an economic impact analysis within 5 years of CCP approval to determine how the district's existence and management activities affect the local and state economies.

### **Socioeconomics Objective B**

Within 5 years of CCP approval, evaluate the aesthetic and environmental benefits of the district's existence and management activities to the state and local communities.

### **Socioeconomics Strategies**

1. Through joint venture partnerships, work collaboratively with USGS–BRD economists or state universities to develop an economic impact analysis of district management actions and the recreation that the WPAs provide.
2. Work with RWBJV partners, university and USGS–BRD ecologists, sociologists, and landscape architects to develop an environmental and aesthetical impact analysis of WPAs in the district.

### **Socioeconomics Rationale**

The WPAs provide a service to communities and to those who visit the areas. Wetlands improve water quality, recharge groundwater, control erosion, and provide flood control. Wetlands provide habitat for many species of wildlife and offer recreational opportunities. However, it is not known to what extent the district's wetlands provide these services and benefits.

The basin is intensively farmed and many of the local citizens see wetlands as a detriment to farming operations. The loss of cropland due to seasonal flooding has caused many landowners to drain or fill the wetlands, which removes the benefits they may provide to the community. An accurate assessment of the wetlands' economic and social worth will help increase public understanding of the value of protecting wetlands.

## **OPERATIONS GOAL**

Safely and efficiently use funding, staffing, infrastructure, and partnerships to achieve the purpose and objectives of the Rainwater Basin Wetland Management District.

### **Operations Objective A**

Within 10 years of CCP approval, build and maintain Service-owned facilities that serve as an office, visitor contact center, maintenance shop, equipment storage, and housing for researchers, volunteers, and seasonal employees.

### **Operations Objective B**

Within 2 years of CCP approval, construct adequate storage facilities for heavy and farm equipment currently stored at McMurtrey and Cottonwood WPAs.

### **Operations Objective C**

Through the duration of the CCP, continue to maintain adequate housing facilities for researchers and volunteers.

### **Operations Objective D**

Through the duration of the CCP, continue to maintain existing roads and dikes at the WPAs.

### **Operations Objective E**

Through the duration of the CCP, continue to maintain equipment and vehicles at or above Service standards.

### **Operations Objective F**

Within 5 years of CCP approval, strive to obtain additional funding for necessary staffing to address the needs of the district.

### **Operations Strategies**

1. Work with partners and the regional office to obtain funding and secure a suitable site for the construction of a Service-owned facility.
2. Prioritize the building and maintenance schedule based on funding projects in the Service Asset Maintenance Management System (SAMMS). Identify an office/visitor center as the top priority construction project.
3. Schedule equipment and vehicle replacements to achieve industry standards when normal life expectancy is reached.
4. Seek mutual agreements to maintain roads that provide access to the WPAs.
5. Work with partners and the regional office to obtain funding to fill four additional positions: outdoor recreation planner, law enforcement officer (park ranger), maintenance worker, and refuge operations specialist.
6. Seek to close minimum-maintenance roads that dissect four WPAs.

## Operations Rationale

The district is operating out of the same building it leased in the late 1970s. At that time, the number of properties managed and the amount of equipment and staff were significantly less than what exists today. The building was constructed as a metal warehouse. Before leasing it, a portion of the warehouse was converted to office space. In the mid-1990s, more office space was needed and the shop area was reduced to allow for three additional offices.

Since the property was first leased, the work accomplished by the district has drastically changed. Staff has changed with the addition of a wildlife biologist, two Partners for Fish and Wildlife biologists, two permanent fire specialists, and three seasonal firefighters. With more staff has come more office and field equipment. The current building is not adequate to store the additional fire engines and equipment. Heavy equipment and some of the vehicles have to be stored at two other locations: McMurtrey WPA and Cottonwood WPA. Both areas are more than 40 miles from the office/shop.

The office portion of the building has desks crowded together with limited space for filing cabinets, computers, and books. The conference room is a small room, partitioned off from the storage room, located above the offices. Because the office is contained within the warehouse, the ventilation system draws shop fumes (welding, vehicle exhaust, and chemicals) into the office area.

The facility is located within an older industrial park. The immediate neighbors include an older trailer park, auto repair shops, grain elevator, and outdoor storage yard. The location of the facility and its appearance are not inviting to the public. Visitors are primarily delivery persons and a few cooperating landowners. The visitor contact portion of the office also serves as the mailroom and photocopying room. Staff vehicles fill the small parking lot and some parking occurs on the street. Although a security fence protects the storage yard, theft and vandalism still occur.

Storage facilities for vehicles and heavy equipment are lacking. Nearly all the equipment remains exposed to extreme weather conditions.

Temporary quarters for researchers and volunteers are old, surplus mobile homes. Mice infest the mobile homes, which are located on a site that does not have potable water. In addition, the temporary quarters are located 40 miles from the office, making it difficult to arrange work schedules between volunteers and staff. In recent years, arrangements with The Nature Conservancy have allowed their rural office/home to house fire crew and volunteers. Numerous opportunities to have research studies and prescribed fires conducted in the district have not happened because the district lacks housing.

Two dual-function officers provide law enforcement operations. However, their primary responsibility is land management. The time allocated toward law enforcement is not adequate to address game violations and vandalism.

## 4.2 STAFF AND FUNDING

The district has a staff of 12 full-time employees. Table 10 lists these positions along with four new positions that are needed for full implementation of the CCP. Projects required to carry out the CCP are funded through two separate systems, as follows:

- The Refuge Operations Needs System (RONS) is used to document requests to Congress for funding and staffing needed to carry out projects above the existing base budget.
- The SAMMS is used to document the equipment, buildings, and other existing properties that require repair or replacement.

Lists of the RONS and SAMMS projects required to carry out this CCP (including maintenance of structures and equipment to a safe and productive standard for the 15 years of the CCP) are in appendix O.

## 4.3 STEP-DOWN MANAGEMENT PLANS

The final CCP for the Rainwater Basin Wetland Management District will be a broad umbrella plan that (1) outlines general concepts and objectives for habitat, wildlife, visitor services, cultural resources, partnerships, and operations; and (2) guides district management for the next 15 years. Step-down management plans provide greater detail for carrying out specific actions authorized by the CCP. Table 11 presents step-down management plans that are anticipated to be needed, along with their current status and next revision date.

## 4.4 MONITORING AND EVALUATION

Adaptive management is a flexible approach to long-term management of biotic resources. Adaptive management is directed, over time, by the results of ongoing monitoring activities and other information. More specifically, adaptive management is a process by which projects are carried out within a framework of scientifically driven experiments to test the predictions and assumptions outlined within a CCP.

To apply adaptive management, specific survey, inventory, and monitoring protocols will be adopted for the district's WPAs. The habitat management strategies will be systematically evaluated to determine management effects on wildlife populations. This information will be used to refine approaches and determine how effectively the objectives are being accomplished. Evaluations will include participation by appropriate partners. If monitoring and evaluation indicate undesirable effects for target and nontarget

**Table 10. Current and proposed staff for Rainwater Basin Wetland Management District, Nebraska.**

<i>Staff Group</i>	<i>Current Positions</i>	<i>Additional Proposed Positions (Unfunded Staff)</i>
management	refuge project leader, GS-13 deputy project leader, GS-12 refuge operations specialist, GS-9	refuge operations specialist, GS-9
biological	wildlife biologist, GS-11 biological technician (wildlife), GS-7	no additional positions
visitor service	none	outdoor recreation planner, GS-11
administrative	administrative support assistant, GS-8	no additional positions
maintenance	maintenance worker, WG-8 biological technician (wildlife), GS-6	maintenance worker, WG-7
fire management	prescribed fire specialist, GS-9 supervisory range technician, GS-7	no additional positions
law enforcement	none	park ranger, GS-9
Partners for Fish and Wildlife	fish and wildlife biologist, GS-9 fish and wildlife biologist, GS-9	no additional positions

GS =General schedule position.

WG=Wage grade position.

**Table 11. Step-down management plans for Rainwater Basin Wetland Management District, Nebraska.**

<i>Step-down Management Plan</i>	<i>Completed Plan (year approved)</i>	<i>New or Revised Plan, (completion year)</i>
disease contingency plan	2006	2013
fire management plan	1998	2009
habitat management plan	—	2010
habitat management plan (annual)	2007	2008
integrated pest management plan	2003	2008
law enforcement plan	—	2010
prairie dog management plan	2003	2011
safety plan	2004	2009
visitor services plan	—	2012
water management plan	2007	2008

species or communities, alterations to the management projects will be made. Subsequently, the CCP will be revised.

## 4.5 PLAN AMENDMENT AND REVISION

This CCP will be reviewed annually to determine the need for revision. The Service will revise this CCP if and when significant information becomes available. This CCP will be supported by detailed step-down management plans to address the completion of specific strategies in support of the district's goals and objectives. Revisions to this CCP and the step-down management plans will be subject to public review and compliance with the NEPA. At a minimum, this CCP will be evaluated every 5 years and revised after 15 years.

# Glossary

**accessible**—Pertaining to physical access to areas and activities for people of different abilities, especially those with physical impairments.

**adaptive resource management**—Rigorous application of management, research, and monitoring to gain information and experience necessary to assess and modify management activities; a process that uses feedback from research, monitoring, and evaluation of management actions to support or modify objectives and strategies at all planning levels; a process in which policy decisions are carried out within a framework of scientifically driven experiments to test predictions and assumptions inherent in a management plan. Analysis of results helps managers determine whether current management should continue as is or whether it should be modified to achieve desired conditions.

**Administration Act**—National Wildlife Refuge System Administration Act of 1966.

**alternative**—Reasonable way to solve an identified problem or satisfy the stated need (40 CFR 1500.2); one of several different means of accomplishing refuge and district purposes and goals and contributing to the Refuge System mission (The Fish and Wildlife Service Manual, 602 FW 1.5).

**amphibian**—Class of cold-blooded vertebrates including frogs, toads, or salamanders.

**annual**—Plant that flowers and dies within 1 year of germination.

**ATV**—All-terrain vehicle.

**acre-foot**—Amount of water it takes to cover a level acre of land (43,560 square feet) to a depth of 1 foot; about 43,560 cubic feet of water or 325,851 gallons.

**avian**—Relating to or characteristic of birds.

**baseline**—Set of critical observations, data, or information used for comparison or a control.

**basin**—Referring to the landform that acts as a water catchment; here used generically to refer to the hydric footprint that pools water.

**the basin**—*See* Rainwater Basin.

**bioenergetics**—Study of energy transformation in living systems.

**biological control**—Use of organisms or viruses to control invasive plants or other pests.

**biological diversity, also biodiversity**—Variety of life and its processes, including the variety of living organisms, the genetic differences among them, and the communities and ecosystems in which they occur (The Fish and Wildlife Service Manual, 052 FW 1.12B). The National Wildlife Refuge System's focus is on indigenous species, biotic communities, and ecological processes.

**biotic**—Pertaining to life or living organisms; caused, produced by, or comprising living organisms.

**CAFO**—Concentrated animal-feeding operation.

**canopy**—Layer of foliage, generally the uppermost layer, in a vegetative stand; midlevel or understory vegetation in multilayered stands; canopy closure (*also canopy cover*) is an estimate of the amount of overhead vegetative cover.

**catabolized (catabolism)**—Breakdown of more complex substances into simpler ones, with the release of energy.

**CCP**—*See* comprehensive conservation plan.

**CFR**—*See* Code of Federal Regulations.

**Code of Federal Regulations (CFR)**—Codification of the general and permanent rules published in the "Federal Register" by the executive departments and agencies of the federal government. Each volume of the CFR is updated once each calendar year.

**COMLG**—Conservation Order for Mid-continent Light Geese.

**compatibility determination**—*See* compatible use.

**compatible use**—Wildlife-dependent recreational use or any other use of a refuge or district that, in the sound professional judgment of the director of the U.S. Fish and Wildlife Service, will not materially interfere with or detract from the fulfillment of the mission of the Refuge System or the purposes of the refuge or district (The Fish and Wildlife Service Manual, 603 FW 3.6). A compatibility determination supports the selection of compatible uses and identified stipulations or limits necessary to ensure compatibility.

**comprehensive conservation plan (CCP)**—Document that describes the desired future conditions of the

refuge or district and provides long-range guidance and management direction for the refuge or wetland district manager to accomplish the purposes of the refuge or district, contribute to the mission of the Refuge System, and to meet other relevant mandates (The Fish and Wildlife Service Manual, 602 FW 1.5).

**concern**—*See* issue.

**cool-season grasses**—Grasses that begin growth earlier in the season and often become dormant in the summer. These grasses will germinate at lower temperatures.

**coteau**—Hilly upland including the divide between two valleys; a divide; the side of a valley.

**cover, also cover type, canopy cover**—Present vegetation of an area.

**critical habitat**—Area essential to the survival of a species; includes all air, land, and water that a species requires to carry out its normal living patterns, as well as other living things used by the species for food, shelter, or other necessary activities.

**cultural resources**—Remains of sites, structures, or objects used by people in the past.

**CWCS**—Comprehensive wildlife conservation strategy.

**cyanobacteria**—Blue-green algae; widely distributed group of predominantly photosynthetic prokaryotic organisms of the subkingdom Cyanophyta, resembling phototrophic bacteria, occurring singly or in colonies in diverse habitats: some species can fix atmospheric nitrogen.

**dense nesting cover**—Composition of grasses and forbs that allows for a dense stand of vegetation that protects nesting birds from the view of predators, usually consisting of one to two species of wheatgrass, alfalfa, and sweetclover.

**the district**—*See* wetland management district.

**drawdown**—Act of manipulating water levels in an impoundment to allow for the natural drying-out cycle of a wetland.

**DUD**—Duck use-day; number of days that an area can support a duck's energetics' needs.

**EA**—*See* environmental assessment.

**ecosystem**—Dynamic and interrelating complex of plant and animal communities and their associated nonliving environment; a biological community, together with its environment, functioning as a unit. For administrative purposes, the Service has designated 53 ecosystems covering the United States and its possessions. These ecosystems generally correspond with watershed boundaries and their sizes and ecological complexity vary.

**emergent**—Plant rooted in shallow water and having most of the vegetative growth above water such as cattail and hardstem bulrush.

**endangered species, federal**—Plant or animal species listed under the Endangered Species Act of 1973, as amended, that is in danger of extinction throughout all or a significant portion of its range.

**endangered species, state**—Plant or animal species in danger of becoming extinct or extirpated in a particular state within the near future if factors contributing to its decline continue. Populations of these species are at critically low levels or their habitats have been degraded or depleted to a significant degree.

**endogenous**—Stored reserves of lipids, proteins, and other nutrients; energy generated from mobilizing these reserves.

**environmental assessment (EA)**—Concise public document, prepared in compliance with the National Environmental Policy Act, that briefly discusses the purpose and need for an action and alternatives to such action, and provides sufficient evidence and analysis of impacts to determine whether to prepare an environmental impact statement or finding of no significant impact (40 CFR 1508.9).

**epizootic**—Pertaining to a disease that affects large numbers of animals throughout a large area and spreads with great speed.

**erosion**—Wearing away of the land surface by various natural processes such as wind and moving water in the form of rivers, streams, rain, and melting snow.

**eutrophication**—Overenrichment of a waterbody with nutrients, resulting in the excessive growth of organisms and the depletion of oxygen.

**evaporation**—Physical process by which a liquid or solid is transformed to a gas.

**exogenous**—Pertaining to a nutrient that is not stored and is readily available.

**exotic**—Nonnative species of plants or animals often brought into an area by human activity.

**extinction**—Complete disappearance of a species from the earth; no longer existing.

**extirpation**—Extinction of a population; complete eradication of a species within a specified area.

**fauna**—All the vertebrate and invertebrate animals of an area.

**federal trust species**—Species where the federal government has primary jurisdiction including federally endangered or threatened species, migratory birds, anadromous fish, and certain marine mammals.

**FGDC**—Federal Geographic Data Committee.

**flood**—Unusual accumulation of water above the ground caused by heavy rain, melting snow, or rapid runoff; a temporary condition of partial or complete inundation of lands that normally do not pool water throughout the entire year.

**floodplain**—Low-lying, nearly level area along a river or stream that is periodically subject to being flooded by water from any source.

**flora**—Plants, specifically plants within a particular set of boundaries that may be geographical, temporal, or biological.

**flyway**—Pathway taken by migrating birds to or from their nesting grounds in northern North America to their overwintering grounds in southern parts of the United States; a broad, well-defined grouping of migration corridors.

**FMP**—Fire management plan.

**forb**—Broad-leaved, herbaceous plant; a seed-producing annual, biennial, or perennial plant that does not develop persistent woody tissue but dies down at the end of the growing season.

**fragmentation**—Alteration of a large block of habitat that creates isolated patches of the original habitat that are interspersed with a variety of other habitat types; the process of reducing the size and connectivity of habitat patches, making movement of individuals or genetic information between parcels difficult or impossible.

**geographic information system (GIS)**—Computer system capable of storing and manipulating spatial data; a set of computer hardware and software for analyzing and displaying spatially referenced features (such as points, lines and polygons) with nongeographic attributes such as species and age.

**GIS**—*See* geographic information system.

**goal**—Descriptive, open-ended, and often broad statement of desired future conditions that conveys a purpose but does not define measurable units (The Fish and Wildlife Service Manual, 620 FW 1.5).

**groundwater recharge**—Act of adding water to the aquifer by surface water movement downward through the soil.

**GS**—General schedule (pay rate schedule for certain federal positions).

**habitat**—Suite of existing environmental conditions required by an organism for survival and reproduction; the place where an organism typically lives and grows.

**habitat type, also vegetation type, cover type**—Land classification system based on the concept of distinct plant associations.

**hemi-marsh**—Wetland with a 50–50 interspersed of open-water and emergent vegetation.

**hydric soil**—Soil that is saturated, flooded, or pooled long enough during the growing season to develop conditions that do not require oxygen and that favor the growth and regeneration of hydrophytic vegetation; hydric soils suggest the presence of wetlands.

**hydrophyte**—Plant that is adapted to grow in water; a wetland plant species.

**hydrophytic vegetation**—Visible plants growing in water or on a substrate and periodically deficient in oxygen as a result of excessive water content.

**impoundment**—Body of water created by collection and confinement within a series of levees or dikes, creating separate management units although not always independent of one another.

**Improvement Act**—National Wildlife Refuge System Improvement Act of 1997.

**indigenous**—Originating or occurring naturally in a particular place.

**integrated pest management (IPM)**—Methods of managing undesirable species such as invasive plants; education, prevention, physical or mechanical methods of control, biological control, responsible chemical use, and cultural methods.

**“interseed”**—Mechanical seeding of one or several plant species into existing stands of established vegetation.

**interspersed**—Diversity of vegetation spread throughout a stand.

**interstitial flow**—Movement of surface water into the ground by flowing through individual soil grains and eventually adding to the groundwater levels.

**introduced species**—Species present in an area due to intentional or unintentional escape, release, dissemination, or placement into an ecosystem as a result of human activity.

**invasive plant, also noxious weed**—Species that is nonnative to the ecosystem under consideration and whose introduction causes, or is likely to cause, economic or environmental harm or harm to human health.

**invertebrate**—Animal without an backbone.

**involute sanctuary**—Place of refuge or protection where animals and birds may not be hunted.

**IPM**—*See* integrated pest management.

**issue**—Unsettled matter that requires a management decision; for example, a Service initiative, opportunity, resource management problem, a threat to the resources of the unit, conflict in uses, public concern,

or the presence of an undesirable resource condition (The Fish and Wildlife Service Manual, 602 FW 1.5).

**limiting factor**—Chemical or physical factor that limits the growth, abundance, or distribution of the population of a species in an ecosystem and determines whether or not an organism can survive.

**lipids**—Group of organic compounds including the fats, oils, waxes, sterols, and triglycerides that are insoluble in water but soluble in nonpolar organic solvents, are oily to the touch, and together with carbohydrates and proteins constitute the principal structural material of living cells.

**macropore flow**—Movement of surface water through large holes in the ground such as animal burrows, desiccation cracks, root tubes and solution pipes, eventually adding to the groundwater levels.

**management alternative**—*See* alternative.

**ME**—*See* metabolized energy.

**metabolized energy (ME)**—Energy produced by digestion.

**migration**—Regular extensive, seasonal movements of birds between their breeding regions and their wintering regions; to pass usually periodically from one region or climate to another for feeding or breeding.

**migratory birds**—Birds that follow a seasonal movement from their breeding grounds to their wintering grounds; includes waterfowl, shorebirds, raptors, and songbirds.

**mission**—Succinct statement of purpose or reason for being.

**mitigation**—Measure designed to counteract an environmental impact or to make an impact less severe.

**mixed-grass prairie**—Transition zone between the tall-grass prairie and the short-grass prairie dominated by grasses of medium height that are approximately 2–4 feet tall. Soils are not as rich as the tall-grass prairie and moisture levels are less.

**monitoring**—Process of collecting information to track changes of selected parameters over time.

**NASS**—Natural Agricultural Statistics Service.

**national wildlife refuge (NWR)**—Designated area of land, water, or an interest in land or water within the National Wildlife Refuge System, but does not include coordination areas; a complete listing of all units of the Refuge System is in the current “Annual Report of Lands Under Control of the U.S. Fish and Wildlife Service.”

**National Wildlife Refuge System (Refuge System)**—Various categories of areas administered by the Secretary of the Interior for the conservation of fish

and wildlife including species threatened with extinction, all lands, waters, and interests therein administered by the Secretary as wildlife refuges, areas for the protection and conservation of fish and wildlife that are threatened with extinction, wildlife ranges, game ranges, wildlife management areas, and waterfowl production areas.

**National Wildlife Refuge System Improvement Act of 1997 (Improvement Act)**—Sets the mission and the administrative policy for all units of the National Wildlife Refuge System; defines a unifying mission for the Refuge System; establishes the legitimacy and appropriateness of the six priority public uses (hunting, fishing, wildlife observation, photography, environmental education, and interpretation); establishes a formal process for determining appropriateness and compatibility; establish the responsibilities of the Secretary of the Interior for managing and protecting the Refuge System; requires a comprehensive conservation plan for each unit of the Refuge System by the year 2012. This act amended portions of the Refuge Recreation Act and National Wildlife Refuge System Administration Act of 1966.

**native species**—Species that, other than as a result of an introduction, historically occurred or currently occurs in that ecosystem.

**NAWMP**—North American Waterfowl Management Plan.

**NDEQ**—Nebraska Department of Environmental Quality.

**NEPA**—National Environmental Policy Act.

**nest success**—Percentage of nests that successfully hatch one or more eggs of the total number of nests initiated in an area.

**NGPC**—Nebraska Game and Parks Commission.

**niche partitioning**—Distribution of species by preference of habitat conditions and needs.

**NOI**—Notice of intent; published in the “Federal Register.”

**nongovernmental organization**—Any group that is not comprised of federal, state, tribal, county, city, town, local, or other governmental entities.

**noxious weed, also invasive plant**—Any living stage (including seeds and reproductive parts) of a parasitic or other plant of a kind that is of foreign origin (new to or not widely prevalent in the U.S.) and can directly or indirectly injure crops, other useful plants, livestock, poultry, other interests of agriculture, including irrigation, navigation, fish and wildlife resources, or public health. According to the Federal Noxious Weed Act (PL 93-639), a noxious weed (such as an invasive plant) is one that causes disease or has adverse effects on humans or the human environment and, therefore,

is detrimental to the agriculture and commerce of the United States and to public health, and is listed as such on Nebraska Noxious Weed List.

**NVCS**—National Vegetation Classification System.

**NWR**—*See* national wildlife refuge.

**objective**—Concise target statement of what will be achieved, how much will be achieved, when and where it will be achieved, and who is responsible for the work; derived from goals and provide the basis for determining management strategies. Objectives should be attainable and time-specific and should be stated quantitatively to the extent possible. If objectives cannot be stated quantitatively, they may be stated qualitatively (The Fish and Wildlife Service Manual, 602 FW 1.5).

**ORP**—Outdoor recreation planner.

**palustrine**—Pertaining to nonflowing wetlands that can be dominated by emergent or submergent plant wetlands with less than 0.5% salinity.

**patch**—Area distinct from that around it; an area distinguished from its surroundings by environmental conditions.

**pathogen**—Agent that causes disease, especially a living microorganism such as a bacterium or fungus.

**pelagic waters**—The surface waters and middle depths of the open ocean, extending from the edge of the continental shelf.

**perennial**—Lasting or active through the year or through many years; a plant species that has a life span of more than 2 years.

**periphyton**—Complex matrix of organisms such as algae and microbes that are attached to submerged plants; serves as an important food source for invertebrates and some fish; important indicator of water quality.

**PILT**—Payments in lieu of taxes.

**PL**—Public law.

**plant community**—Assemblage of plant species unique in its composition; occurs in particular locations under particular influences; a reflection or integration of the environmental influences on the site such as soil, temperature, elevation, solar radiation, slope, aspect, and rainfall; denotes a general kind of climax plant community, such as ponderosa pine or bunchgrass.

**playa**—Nearly level, flat area that is temporarily covered with water, at the bottom of an undrained basin.

**PM**—Particulate matter.

**prescribed fire**—Skillful application of fire to natural fuels under conditions such as weather, fuel moisture, and soil moisture that allow confinement of the fire to

a predetermined area and produces the intensity of heat and rate of spread to accomplish planned benefits to one or more objectives of habitat management, wildlife management, or hazard reduction.

**priority public use**—One of six uses authorized by the National Wildlife Refuge System Improvement Act of 1997 to have priority if found to be compatible with the purposes for a refuge or district. This includes hunting, fishing, wildlife observation, photography, environmental education, and interpretation.

**proposed action**—Alternative proposed to best achieve the purpose, vision, and goals of a refuge or district (contributes to the Refuge System mission, addresses the significant issues, and is consistent with principles of sound fish and wildlife management).

**public**—Individuals, organizations, and groups; officials of federal, state, and local government agencies; Indian tribes; and foreign nations. It may include anyone outside the core planning team. It includes those who may or may not have indicated an interest in Service issues and those who do or do not realize that Service decisions may affect them.

**public involvement**—Process that offers affected and interested individuals and organizations an opportunity to become informed about, and to express their opinions on, Service actions and policies. In the process, these views are studied thoroughly and thoughtful consideration of public views is given in shaping decisions for refuge and district management.

**purpose of the district**—Reason for and role of a district that is specified in or derived from the law, proclamation, executive order, agreement, public land order, donation document, or administrative memorandum establishing authorization or expanding a district or district subunit such as a waterfowl production area (The Fish and Wildlife Service Manual, 602 FW 1.5).

**Rainwater Basin (the basin)**—Geographic area in south-central Nebraska that contains closed-basin wetlands with characteristic clay soils that have very low transmissivity.

**raptor**—Carnivorous bird such as a hawk, a falcon, or a vulture that feeds wholly or chiefly on meat taken by hunting or on carrion (dead carcasses).

**RWBJV**—Rainwater Basin Joint Venture.

**refuge**—*See* national wildlife refuge.

**Refuge Operating Needs System (RONS)**—National database that contains the unfunded operational needs of each refuge and district. Projects included are those required to carry out approved plans and meet goals, objectives, and legal mandates.

**Refuge System**—*See* National Wildlife Refuge System.

**resident species**—Species inhabiting a given locality throughout the year; nonmigratory species.

**rest**—Free from biological, mechanical, or chemical manipulation, in reference to refuge and district lands.

**restoration**—Management emphasis designed to move ecosystems to desired conditions and processes, such as healthy upland habitats and aquatic systems.

**riparian area or riparian zone**—Area or habitat that is transitional from terrestrial to aquatic ecosystems including streams, lakes, wet areas, and adjacent plant communities and their associated soils that have free water at or near the surface; an area whose components are directly or indirectly attributed to the influence of water; of or relating to a river; specifically applied to ecology, “riparian” describes the land immediately adjoining and directly influenced by streams. For example, riparian vegetation includes all plant life growing on the land adjoining a stream and directly influenced by the stream.

**RONs**—*See* Refuge Operating Needs System.

**SAMMS**—*See* Service Asset Maintenance Management System.

**scoping**—Process of obtaining information from the public for input into the planning process.

**scouring**—Removal of earth or rock by the action of running water or wind eroding material.

**seasonally flooded**—Surface water is present for extended periods in the growing season, but is absent by the end of the season in most years.

**sediment**—Material deposited by water, wind, and glaciers.

**sedimentation**—Deposit of waterborne particles, resulting from a decrease in water’s transport capacity.

**seeping**—Movement of water into or through porous material.

**Service**—*See* U.S. Fish and Wildlife Service.

**Service Asset Maintenance Management System (SAMMS)**—National database which contains the unfunded maintenance needs of each refuge and district; projects include those required to maintain existing equipment and buildings, correct safety deficiencies for the implementation of approved plans, and meet goals, objectives, and legal mandates.

**shelterbelt**—Single to multiple rows of trees and shrubs planted around cropland or buildings to block or slow down the wind.

**shorebird**—Any of a suborder (Charadrii) of birds such as plover or snipe that frequent the seashore or mud flats.

**spatial**—Relating to, occupying, or having the character of space.

**special status species**—Plants or animals that have been identified through federal law, state law, or agency policy as requiring special protection of monitoring. Examples include federally listed endangered, threatened, proposed, or candidate species; state-listed endangered, threatened, candidate, or monitor species; Service’s species of management concern; species identified by the Partners in Flight Program as being of extreme or moderately high conservation concern.

**special use permit**—Special authorization from the refuge or wetland district manager that is required for any refuge and district service, facility, privilege, or product of the soil provided at refuge or district expense and not usually available to the general public through authorizations in Title 50 CFR or other public regulations (Refuge Manual, 5 RM 17.6).

**species of concern**—Plant and animal species, while not falling under the definition of special status species, that are of management interest by virtue of being federal trust species such as migratory birds, important game species, or significant keystone species; species that have documented or apparent population declines, small or restricted populations, or dependence on restricted or vulnerable habitats.

**step-down management plan**—Plan that provides the details necessary to carry out management strategies identified in the comprehensive conservation plan (The Fish and Wildlife Service Manual, 602 FW 1.5).

**strategy**—Specific action, tool, or technique or combination of actions, tools, and techniques used to meet unit objectives (The Fish and Wildlife Service Manual, 602 FW 1.5).

**submergent**—Vascular or nonvascular hydrophyte, either rooted or nonrooted, that lies entirely beneath the water surface, except for flowering parts in some species.

**succession**—Natural replacement of one biotic community by another; a slow but continuous process, beginning with the invasion of a patch of open ground or newly created body of water by pioneer species. Succession continues through a series of recognizable stages known as seres, ending with the formation of a climax community in which the mix of species forming the community no longer changes with successive generations; late successional communities can also be climax communities.

**SWG**—State wildlife grant.

**threatened species, federal**—Species listed under the Endangered Species Act of 1973, as amended, that are likely to become endangered within the foreseeable future throughout all or a significant portion of their range.

**threatened species, state**—Plant or animal species likely to become endangered in a particular state within the near future if factors contributing to population decline or habitat degradation or loss continue.

**tree harvest**—Commercial or private removal of trees for economic or personal benefit.

**trust species**—*See* federal trust species.

**UNL**—University of Nebraska–Lincoln.

**USC**—United States Code.

**USDA**—U.S. Department of Agriculture.

**USDOE**—U.S. Department of Energy.

**USEPA**—U.S. Environmental Protection Agency.

**U.S. Fish and Wildlife Service (Service, USFWS)**—Federal agency of the U.S. Department of the Interior that is responsible for conserving, protecting, and enhancing fish and wildlife and their habitats for the continuing benefit of the American people. The Service manages the 93-million-acre National Wildlife Refuge System comprised of more than 530 national wildlife refuges and wetland management districts and thousands of waterfowl production areas. It also operates 65 national fish hatcheries and 78 ecological service field stations, the agency enforces federal wildlife laws, manages migratory bird populations, restores national significant fisheries, conserves and restores wildlife habitat such as wetlands, administers the Endangered Species Act, and helps foreign governments with their conservation efforts. It also oversees the federal aid program that distributes millions of dollars in excise taxes on fishing and hunting equipment to state wildlife agencies.

**USFWS**—*See* U.S. Fish and Wildlife Service.

**U.S. Geological Survey (USGS)**—Federal agency of the U.S. Department of the Interior whose mission is to provide reliable scientific information to describe and understand the earth; minimize loss of life and property from natural disasters; manage water, biological, energy, and mineral resources; and enhance and protect our quality of life.

**USGS**—*See* U.S. Geological Survey.

**USGS–BRD**—Biological Research Division of the U.S. Geological Survey.

**vision statement**—Concise statement of the desired future condition of the planning unit, based primarily on the Refuge System mission, specific refuge or district purposes, and other relevant mandates (The Fish and Wildlife Service Manual, 602 FW 1.5).

**waterfowl**—Category of birds that includes ducks, geese, and swans.

**watershed**—Area that drains into a river, a river system, or a body of water.

**wetland**—Area of land that pools water long enough annually to support hydrophytic vegetation.

**wetland management district (the district, WMD)**—Land that the Refuge System acquires with Federal Duck Stamp funds for restoration and management primarily as prairie wetland habitat critical to waterfowl and other wetland birds.

**wet meadow**—Area where wetland and upland plants can occur, usually along the edge of a wetland; may have pooled water for a short period of time, usually in the spring or after a heavy rain event.

**WG**—Wage grade schedule (pay rate schedule for certain federal positions).

**wildland fire**—Free-burning fire requiring a suppression response; all fire other than prescribed fire that occurs on wildlands (Draft, The Fish and Wildlife Service Manual 621 FW 1.7).

**wildlife-dependent recreational use**—Use of a refuge or district involving hunting, fishing, wildlife observation, photography, environmental education, or interpretation. The National Wildlife Refuge System Improvement Act of 1997 specifies that these are the six priority general public uses of the Refuge System.

**WMA**—Wildlife management area.

**WMD**—*See* wetland management district.

**woodland**—Open stands of trees with crowns not usually touching, generally forming 25–60% cover.

**WPA**—Waterfowl production area.



# Appendix A

## *Key Legislation and Policies*

Administration of units of the National Wildlife Refuge System is governed by (1) bills passed by the U.S. Congress and signed into law by the president of the United States, and (2) by regulations developed by the various branches of the government. Following are brief descriptions of some of the most pertinent laws and statutes establishing legal parameters and policy direction for the Refuge System.

*In alphabetical order of the name of the act, order, or regulation.*

**American Indian Religious Freedom Act:** Affirms the right of Native Americans to have access to their sacred places. If a place of religious importance to Native Americans may be affected by an undertaking, the act promotes consultation with Native American religious practitioners, which may be coordinated with section 106 consultation.

**Americans with Disabilities Act (1992):** Prohibits discrimination in public accommodations and services.

**Antiquities Act (June 8, 1906; 16 USC 431–433; 34 Stat. 225):** Authorizes the president to designate as national monuments objects or areas of historic or scientific interest on lands owned or controlled by the United States. Requires that a permit be obtained for examination of ruins, excavation of archaeological sites, and the gathering of objects of antiquity on lands under the jurisdiction of the Secretaries of Interior, Agriculture, and Army, and provided penalties for violations.

**Archaeological Resources Protection Act (Public Law [PL] 96-95; October 31, 1979; 16 USC 470aa–470ll; 93 Stat. 721):** Protects archaeological resources and sites on public (federal) lands and Native American lands. The act calls for the preservation of objects and associated records in a suitable repository once recovered from a site. The act sets guidelines for proper procedures to obtain permission and permits to excavate archaeological sites on public lands by qualified individuals.

**Architectural Barriers Act (1968):** Requires federally owned, leased, or funded buildings and facilities to be accessible to persons with disabilities.

**Archeological and Historic Preservation Act (PL 86-523; June 27, 1960; 16 USC 469–469c; 74 Stat. 220 [as amended by PL 93-291; May 24, 1974; 88 Stat. 174]):** Carries out the policy established by the Historic Sites,

Buildings and Antiquities Act; directs federal agencies to notify the Secretary of the Interior whenever they find a federal or federally assisted, licensed, or permitted project may cause loss or destruction of significant scientific, prehistoric, or archaeological data. Authorizes use of appropriated, donated, and transferred funds for the recovery, protection, and preservation of such data.

**Clean Water Act (1977):** Requires consultation with the U.S. Army Corps of Engineers for major wetland modifications.

**Criminal Code of Provisions of 1940 (as amended, 18 USC 41):** States the intent of Congress to protect all wildlife within federal sanctuaries, refuges, fish hatcheries, and breeding grounds. Provides that anyone (except in compliance with rules and regulations promulgated by authority of law) who hunts, traps, or willfully disturbs any such wildlife, or willfully injures, molests, or destroys any property of the United States on such land or water, shall be fined up to \$500 or imprisoned for not more than 6 months or both.

**Emergency Wetland Resources Act of 1986:** Authorizes the purchase of wetlands from Land and Water Conservation Fund monies, removing a prior prohibition on such acquisitions. Requires the Secretary to establish a national wetlands priority conservation plan, requires the states to include wetlands in their comprehensive outdoor recreation plans, and transfers to the Migratory Bird Conservation Fund amount equal to import duties on arms and ammunition.

**Endangered Species Act of 1973 and recent amendments (16 USC 1531–1543, 87 Stat. 884; as amended):** Provides for conservation of threatened and endangered species of fish, wildlife, and plants by federal action and by encouraging state programs. Specific provisions include the listing and determination of critical habitat for endangered and threatened species and consultation with the Service on any federally funded or licensed project that could affect any of these agencies; prohibition of unauthorized taking, possession, sale, transport, etc., of endangered species; an expanded program of habitat acquisition; establishment of cooperative agreements and grants-in-aid to states that establish and maintain an active, adequate program for endangered and threatened species; assessment of civil and criminal penalties for violating the act or regulations.

**Environmental Education Act of 1990 (PL 101-619; November 16, 1990; 20 USC 5501–5510; 104 Stat. 3325):**

Establishes the Office of Environmental Education within the USEPA to develop and administer a federal environmental education program. Responsibilities of the office include developing and supporting programs to improve understanding of the natural and developed environment and the relationships between humans and their environment; supporting the dissemination of educational materials; developing and supporting training programs and environmental education seminars; managing a federal grant program; and administering an environmental internship and fellowship program. Requires the office to develop and support environmental programs in consultation with other federal natural resource management agencies including the Service.

**EO 11644—Use of Off-road Vehicles on Public Lands (1972):** Provides policy and procedures for regulating off-road vehicles.

**EO 11988—Floodplain Management (May 24, 1977):** Prevents federal agencies from contributing to the “adverse impacts associated with occupancy and modification of floodplains” and the “direct or indirect support of floodplain development.” In the course of fulfilling their respective authorities, federal agencies “shall take action to reduce the risk of flood loss, to minimize the impact of floods on human safety, health, and welfare, and to restore and preserve the natural and beneficial values served by floodplains.”

**EO 11990—Protection of Wetlands.**

**EO 12996—Management and General Public Use of the National Wildlife Refuge System (1996):** Defines the mission, purpose, and priority public uses of the Refuge System; presents four principles to guide management of the system.

**EO 13007—Indian Sacred Sites (1996):** Directs federal land management agencies to accommodate access to and ceremonial use of Indian sacred sites by Indian religious practitioners, avoid adversely affecting the physical integrity of such sacred sites, and where appropriate, maintain the confidentiality of sacred sites.

**Federal Noxious Weed Act (1990):** Requires the use of integrated management systems to control or contain undesirable plant species, and an interdisciplinary approach with the cooperation of other federal and state agencies.

**Federal Records Act (1950):** Requires the preservation of evidence of the government’s organization, functions, policies, decisions, operations, and activities, as well as basic historical and other information.

**Federal Water Pollution Control Act of 1972, Section 401 (PL 92-500, USC 1411, 86 Stat. 816.33):** Requires any applicant for a federal license or permit to conduct any

activity which may result in a discharge into navigable waters to obtain a certification from the state in which the discharge originates or will originate, or, if appropriate, from the interstate water pollution control agency having jurisdiction over navigable waters at the point where the discharge originates or will originate, that the discharge will comply with applicable effluent limitations and water quality standards. Requires that a certification obtained for construction of any facility must also pertain to subsequent operation of the facility.

**Federal Water Pollution Control Act of 1972, Section 404 (PL 92-500, 86 Stat. 816):** Authorizes the Secretary of the Army, acting through the Chief of Engineers, to issue permits, after notice and opportunity for public hearing, for discharge of dredged or fill material into navigable waters of the United States, including wetlands, at specified disposal sites. Requires that selection of disposal sites be in accordance with guidelines developed by the Administrator of the USEPA in conjunction with the Secretary of the Army. States that the Administrator can prohibit or restrict use of any defined area as a disposal site whenever she/he determines, after notice and opportunity for public hearings, that discharge of such materials into such areas will have an unacceptable adverse effect on municipal water supplies, shellfish beds, fishery areas, wildlife, or recreational areas.

**Fish and Wildlife Act of 1956 (16 USC 742a–742j, 70 Stat. 1119; as amended):** Establishes a comprehensive fish and wildlife policy and directs the Secretary of the Interior to provide continuing research and extension and conservation of fish and wildlife resources.

**Fish and Wildlife Conservation Act of 1980 (PL 96366; September 29, 1980; 16 USC 2901–2911; as amended 1986, 1988, 1990, and 1992):** Creates a mechanism for federal matching funding of the development of state conservation plans for nongame fish and wildlife. States that subsequent amendments to this law require that the Secretary monitor and assess migratory nongame birds, determine the effects of environmental changes and human activities, identify birds likely to be candidates for endangered species listing, and identify conservation actions that would prevent this from being necessary. In 1989, Congress also directed the Secretary to identify lands and waters in the Western Hemisphere, the protection, management, or acquisition of which would foster conservation of migratory nongame birds. All of these activities are intended to assist the Secretary in fulfilling the Secretary’s responsibilities under the Migratory Bird Treaty Act and the Migratory Bird Conservation Act, and provisions of the Endangered Species Act implementing the Convention on Nature Protection and Wildlife Preservation in the Western Hemisphere.

**Fish and Wildlife Coordination Act (1958):** Allows the U.S. Fish and Wildlife Service to enter into agreements with private landowners for wildlife management purposes.

**Fish and Wildlife Improvement Act of 1978:** Improves the administration of fish and wildlife programs and amends several earlier laws including the Refuge Recreation Act, the National Wildlife Refuge System Administration Act, and the Fish and Wildlife Act of 1956. Authorizes the Secretary to accept gifts and bequests of real and personal property on behalf of the United States. Authorizes the use of volunteers for Service projects and appropriations to carry out volunteer programs.

**Historic Sites, Buildings and Antiquities Act (August 21, 1935; 16 USC 461–462, 464–467; 49 Stat. 666; known as the “Historic Sites Act” [as amended by PL 89-249; October 9, 1965; 79 Stat. 971]):** Declares it a national policy to preserve historic sites and objects of national significance, including those located at refuges and districts. Provides procedures for designation, acquisition, administration, and protection of such sites. Provides for designation of National Historic and Natural Landmarks.

**Land and Water Conservation Fund Act of 1965:** Provides funds from leasing bonuses, production royalties, and rental revenues for offshore oil, gas, and sulphur extraction to the Bureau of Land Management, the USDA Forest Service, the U.S. Fish and Wildlife Service, and state and local agencies for purchase of lands for parks, open space, and outdoor recreation.

**Migratory Bird Conservation Act of 1929 (16 USC 715–715d, 715e, 715f–715r):** Establishes the Migratory Bird Conservation Commission, which consists of the Secretaries of the Interior (chair), Agriculture, and Transportation; two members from the House of Representatives; and an ex-officio member from the state in which a project is located. States that the commission approves acquisition of land and water, or interests therein, and sets the priorities for acquisition of lands by the Secretary of the Interior for sanctuaries or for other management purposes. Requires that, to acquire lands or interests therein, the state concerned must consent to such acquisition by legislation. Such legislation has been enacted by most states.

**Migratory Bird Conservation Act of 1929 (16 USC 715s, 45 Stat. 1222, as amended):** Authorizes acquisition, development, and maintenance of migratory bird refuges; cooperation with other agencies in conservation; and investigations and publications on North American birds. Authorizes payment of 25% of net receipts from administration of national wildlife refuges to the country or counties in which such refuges are located.

**Migratory Bird Hunting and Conservation Stamp Act of 1934 (March 16, 1934; 16 USC 718–718h; 48 Stat. 51; known as The “Duck Stamp Act”;** as amended): Requires each waterfowl hunter 16 years of age or older to possess a valid federal hunting stamp. Authorizes the requirement of an annual stamp for the hunting of waterfowl; proceeds go toward the purchase

of habitat for waterfowl and other wildlife. Duck stamps are also purchased (1) for entry into some refuges, (2) by conservationists, and (3) for stamp collections. Receipts from the sale of the stamp are deposited in a special Treasury account known as the Migratory Bird Conservation Fund and are not subject to appropriations.

**Migratory Bird Treaty Act of 1918 (16 USC 703–711; 50 CFR, subchapter B; as amended):** Implements treaties with Great Britain (for Canada) and Mexico for protection of migratory birds whose welfare is a federal responsibility. Provides for regulations to control taking, possession, selling, transporting, and importing of migratory birds and provides penalties for violations. Enables the setting of seasons and other regulations (including the closing of areas, federal or nonfederal) related to the hunting of migratory birds.

**National and Community Service Act of 1990 (PL 101-610; November 16, 1990; 42 USC 12401; 104 Stat. 3127):** Authorizes several programs to engage citizens of the United States in full and part-time projects designed to combat illiteracy and poverty, provide job skills, enhance educational skills, and fulfill environmental needs. Provides for grants to states for the creation of programs for citizens over 17 years of age. Programs must be designed to fill unmet educational, human, environmental, and public safety needs. Initially, participants will receive postemployment benefits of up to \$1,000 per year for part-time participants and \$2,500 for full-time participants.

Several provisions are of particular interest to the Service:

*American Conservation and Youth Service Corps:* As a federal grant program established under subtitle C of the law, the corps offers an opportunity for young adults between the ages of 16 and 25, or in the case of summer programs, between 15 and 21, to engage in approved human and natural resources projects that benefit the public or are carried out on federal or Indian lands. To be eligible for assistance, natural resources programs will focus on improvement of wildlife habitat and recreational areas, fish culture, fishery assistance, erosion, wetlands protection, pollution control, and similar projects. A stipend of not more than 100% of the poverty level will be paid to participants. A commission established to administer the Youth Service Corps will make grants to states, the Secretaries of Agriculture and Interior, and the Director of ACTION to carry out these responsibilities.

*Thousand Points of Light:* Creates a nonprofit Points of Light Foundation to administer programs to encourage citizens and institutions to volunteer to solve critical social issues, discover new leaders, and develop institutions committed to serving others.

**National Environmental Policy Act of 1969 (PL 91-190; January 1, 1970; 42 USC 4321–4347; 83 Stat. 852 [as amended by PL 94-52; July 3, 1975; 89 Stat. 258] [as amended by PL 94-83; August 9, 1975; 89 Stat. 424]):**

Requires all agencies, including the Service, to examine the environmental impacts of their actions, incorporate environmental information, and use public participation in the planning and the implementation of all actions, federal agencies must integrate the act with other planning requirements, and to prepare appropriate documents to facilitate better environmental decision making (40 CFR 1500). Declares national policy to encourage a productive and enjoyable harmony between humans and their environment.

Section 102 of that act directs that “to the fullest extent possible the policies, regulations, and public laws of the United States shall be interpreted and administered in accordance with the policies set forth in this act, and all agencies of the Federal Government shall ... insure that presently unquantified environmental amenities and values may be given appropriate consideration in decision making along with economic technical considerations.”

Section 102(2)c of the NEPA requires all federal agencies, with respect to major federal actions significantly affecting the quality the quality of the human environment, to submit to the Council on Environmental Quality a detailed statement of the environmental impact of the proposed action; any adverse environmental effect that cannot be avoided should the proposal be implemented; alternatives to the proposed action; the relationship between local short-term uses of the environment and the maintenance and enhancement of long-term productivity; any irreversible and irretrievable commitments of resources that would be involved in the proposed action, should it be implemented.

**National Historic Preservation Act of 1966 (PL 89-665; October 15, 1966; 16 USC 470–470b, 470c–470n; 80 Stat. 915; and repeatedly amended):**

Instructs federal agencies to consider the effect their undertakings have on cultural resources. The act is comprehensive legislation with many components, with the most important aspect to management being section 106. The purpose of section 106 is stated in 36 CFR 800.1: “The section 106 process seeks to accommodate historic preservation concerns with the needs of Federal undertakings through consultation among the agency official and other parties with an interest in the effects of the undertaking on historic properties.” Historic property is defined in 36 CFR 800.16(l) as “any prehistoric or historic district, site, building, structure, or object in, or eligible for inclusion in, the National Register of Historic Places maintained by the Secretary of the Interior.” Another important section, section 110, directs federal agencies to inventory cultural resources on public lands—not necessarily in relationship to a project or undertaking—so cultural resources can be evaluated and managed.

**National Wildlife Refuge System Administration Act of 1966 (PL 89-669; 16 USC 668dd–668ee; 80 Stat. 929; as amended):**

Defines the Refuge System as including wildlife refuges, areas for protection and conservation of fish and wildlife that are threatened with extinction, wildlife ranges, game ranges, wildlife management areas, and waterfowl production areas. Authorizes the Secretary to permit any use of an area provided such use is compatible with the major purposes for which such area was established. States that purchase considerations for rights-of-way go into the Migratory Bird Conservation Fund for the acquisition of lands. By regulation, up to 40% of an area acquired for a migratory bird sanctuary may be opened to migratory bird hunting unless the Secretary finds that the taking of any species of migratory game birds in more than 40% of such area would be beneficial to the species. Requires an act of Congress for the divestiture of lands in the system, except for (1) lands acquired with Migratory Bird Conservation Commission funds, and (2) lands that can be removed from the System by land exchange, or if brought into the System by a cooperative agreement, then pursuant to the terms of the agreement.

**National Wildlife Refuge System Improvement Act of 1997 (PL 105-57; October 9, 1997; Amendment to the National Wildlife Refuge System Administration Act of 1966):**

Sets the mission and the administrative policy for all units in the Refuge System. Clearly defines a unifying mission for the Refuge System; establishes the legitimacy and appropriateness of the six priority public uses (hunting, fishing, wildlife observation, photography, environmental education, and interpretation); establishes a formal process for determining appropriateness and compatibility; establishes the responsibilities of the Secretary of the Interior for managing and protecting the Refuge System; and requires a CCP for each refuge by the year 2012. Also amended portions of the Refuge Recreation Act and the National Wildlife Refuge System Administration Act of 1966.

Key provisions include the following:

- a requirement that the Secretary of the Interior ensures maintenance of the biological integrity, diversity, and environmental health of the Refuge System;
- the definition of compatible wildlife-dependent recreation as “legitimate and appropriate general public use of the [National Wildlife Refuge] System”;
- the establishment of hunting, fishing, wildlife observation, photography, environmental education, and interpretation as “priority public uses” where compatible with the mission and purpose of individual national wildlife refuges;
- the refuge managers’ authority to use sound professional judgment in determining which public uses are compatible at national wildlife refuges and whether or not they will be allowed

(a formal process for determining “compatible use” is currently being developed);

- the requirement of open public involvement in decisions to allow new uses of national wildlife refuges and renew existing ones, as well as in the development of CCPs for national wildlife refuges.

**National Wildlife Refuge Regulations (50 CFR 25-35, 43 CFR 3103.2 and 3120.3–3):** Provides regulations for administration and management of national wildlife refuges including mineral leasing, exploration, and development.

**Rights-of-way General Regulations (50 CFR 29.21; 34 CFR 19907, December 19, 1969):** Provides for procedures for filing applications. Provides terms and conditions under which rights-of-way over, above, and across lands administered by the Service may be granted.

**National Wildlife Refuge System Volunteer and Community Partnership Enhancement Act of 1998 (PL 105-242, 112 Stat. 1575):** Encourages the use of volunteers to assist the Service in the management of refuges within the Refuge System. Facilitates partnerships between the Refuge System and nonfederal entities to promote public awareness of the resources of the Refuge System and public participation in the conservation of those resources. Encourages donations and other contributions by persons and organizations to the Refuge System.

**Native American Graves Protection and Repatriation Act:** Provides a process for museums and federal agencies to return certain Native American cultural items—human remains, funerary objects, sacred objects, and objects of cultural patrimony—to lineal descendants, culturally affiliated Indian tribes, and Native Hawaiian organizations.

**North American Wetlands Conservation Act (PL 101-233; December 13, 1989; 16 USC 4401–4412; 103 Stat. 1968):** Provides for the conservation of North American wetland ecosystems, waterfowl and other migratory birds, fish, and wildlife that depend on such habitats. Establishes a council to review project proposals and provided funding for the projects. Provides funding and administrative direction for implementation of the North American Waterfowl Management Plan and the Tripartite Agreement on wetlands between Canada, United States, and Mexico. Converts the Pittman-Robertson account into a trust fund, with the interest available without appropriation through the year 2006 to carry out the programs authorized by the act, along with an authorization for annual appropriation of \$15 million plus an amount equal to the fines and forfeitures collected under the Migratory Bird Treaty Act. Available funds may be expended, upon approval of the Migratory Bird Conservation Commission, for payment of not to exceed 50% of the United States share of the cost of wetlands conservation projects in Canada, Mexico, or the United States (or 100% of the cost of projects on

federal lands). At least 50% and no more than 70% of the funds received are to go to Canada and Mexico each year.

**Refuge Recreation Act of 1962:** Authorizes the Secretary of the Interior to administer refuges, hatcheries, and other conservation areas for recreational use, when such uses do not interfere with the areas’ primary purposes. Authorizes construction and maintenance of recreational facilities and the acquisition of land for incidental fish- and wildlife-oriented recreational development or protection of natural resources. Authorizes the charging of fees for public uses.

**Refuge Recreation Act of 1966 (PL 87-714, 16 USC 460k et seq., 76 Stat. 653–654):** Authorizes appropriate, incidental, or secondary recreational use at conservation areas administered by the Secretary of the Interior for fish and wildlife purposes.

**Refuge Recreation Act of 1969 [16 USC 460k–460k4], as amended.**

**Refuge Revenue Sharing Act, Section 401 (June 15, 1935; 16 USC 715s; 49 Stat. 383):** Provides for payments to counties in lieu of taxes, using revenues derived from the sale of products from refuges. Related legislation follows:

*PL 88-523 (August 30, 1964; 78 Stat. 701):* Makes major revisions by requiring that all revenues received from refuge products such as animals, timber and minerals, or from leases or other privileges, be deposited in a special Treasury account and net receipts distributed to counties for public schools and roads.

*PL 93-509 (December 3, 1974; 88 Stat. 1603):* Requires that monies remaining in the fund after payments be transferred to the Migratory Bird Conservation Fund for land acquisition under provisions of the Migratory Bird Conservation Act.

*PL 95-469 (October 17, 1978; 92 Stat. 1319):* Expands the revenue-sharing system to include national fish hatcheries and Service research stations. Includes in the Refuge Revenue Sharing Fund receipts from the sale of salmonid carcasses. Establishes payments to counties as follows:

On acquired land, the greatest amount calculated on the basis of 75 cents per acre,  $\frac{3}{4}$  of 1% of the appraised value, or 25% of the net receipts produced from the land.

- On land withdrawn from the public domain, 25% of net receipts and basic payments under PL 94-565 (31 USC 1601–1607, 90 Stat. 2662), payment in lieu of taxes on public lands.
- This amendment also authorizes appropriations to make up any difference between the amount in the fund and the amount scheduled for payment in any year. The stipulation that payments be used for schools and roads was

removed, but counties were required to pass payments along to other units of local government within the county that suffer losses in revenues due to the establishment of Service areas.

**Refuge Revenue Sharing Act of 1978 (PL 95-469; October 17, 1978; amended 16 USC 715s; 50 CFR, part 34):** Changes the provisions for sharing revenues with counties in a number of ways. Makes revenue sharing applicable to all lands administered by the Service, whereas previously it was applicable only to areas in the Refuge System. Makes payments available for any governmental purpose, whereas the old law restricted the use of payments to roads and schools. For lands acquired in fee simple, provides a payment of 75 cents per acre,  $\frac{3}{4}$  of 1% of fair market value or 25% of net receipts, whichever is greatest, whereas the old law provided a payment of  $\frac{3}{4}$  of 1% adjustment cost or 25% of net receipts, whichever was greater. Makes reserve (public domain) lands entitlement lands under PL 94-565 (16 USC 1601-1607) and provides for a payment of 25% of net receipts. Authorizes appropriations to make up any shortfall in net receipts, to make payments in the full amount for which counties are eligible. The old law provided that if net receipts were insufficient to make full payment, payment to each county would be reduced proportionality.

**Refuge Trespass Act of June 28, 1906 (18 USC 41, 43 Stat. 98; 18 USC 145):** Provides the first federal protection for wildlife at national wildlife refuges. Makes it unlawful to hunt, trap, capture, willfully disturb, or kill any bird or wild animal, or take or destroy the eggs of any such birds, on any lands of the United States set apart or reserved as refuges or breeding grounds for such birds or animals by any law, proclamation, or executive order, except under rules and regulations of the Secretary. The act also protects government property on such lands.

**Refuge Trespass Act of June 25, 1948 (18 USC 41, Stat. 686; Section 41 of the Criminal Code, Title 18):** Consolidates the penalty provisions of various acts from January 24, 1905 (16 USC 684-687, 33 Stat. 614)

through March 10, 1934 (16 USC 694-694b, 48 Stat. 400) and restates the intent of Congress to protect all wildlife within federal sanctuaries, refuges, fish hatcheries, and breeding grounds. Provides that anyone (except in compliance with rules and regulations promulgated by authority of law) who hunts, traps, or willfully disturbs any wildlife on such areas, or willfully injures, molests, or destroys any property of the United States on such lands or waters, shall be fined, imprisoned, or both.

**Rehabilitation Act of 1973 (October 1, 1973; 29 USC 794 [as amended by PL 93-112, Title 5; 87 Stat. 355]):** Prohibits discrimination on the basis of handicap under any program or activity receiving federal financial assistance.

**Transfer of Certain Real Property for Wildlife Conservation Purposes Act of 1948:** Provides that, upon determination by the Administrator of the General Services Administration, real property no longer needed by a federal agency can be transferred without reimbursement to the Secretary of the Interior if the land has particular value for migratory birds, or to a state agency for other wildlife conservation purposes.

**U.S. Department of the Interior Order No. 3226 (January 19, 2001):** Directs bureaus and offices of the Department to analyze the potential effects on climate change when undertaking long-range planning, when setting priorities for scientific research, and when making major decisions about use of resources.

**Wilderness Act of 1964 (PL 88-577; September 3, 1964):** Directs the Secretary of the Interior, within 10 years, to review every roadless area of 5,000 or more acres and every roadless island (regardless of size) within the Refuge System and National Park Service for inclusion in the National Wilderness Preservation System.

**Wilderness Preservation and Management (50 CFR 35; 16 USC 1131-1136; 43 USC 1201; 78 Stat. 890):** Provides procedures for establishing wilderness units under the Wilderness Act of 1964 at units of the Refuge System.

# Appendix B

## Compatibility Determinations

### DISTRICT NAME

Rainwater Basin Wetland Management District

### ESTABLISHING AND ACQUISITION AUTHORITIES

Migratory Bird Hunting and Conservation Stamp Act (16 USC 718 [c])

Migratory Bird Conservation Act (16 USC 715d[2], 715i[a])

Consolidated Farm and Rural Development Act (7 USC 2002 [a])

Emergency Wetlands Resources Act (16 USC 3901 [b])

Public Land Order 7206 (June 24, 1996)

Public Land Order 6979 (May 25, 1993)

### DISTRICT PURPOSES

*“Small areas, to be designated as ‘Waterfowl Production Areas’ may be acquired without regard to the limitations and requirements of the Migratory Bird Conservation Act, but all of the provisions of such Act which govern the administration and protection of lands acquired thereunder, except the inviolate sanctuary provisions of such Act, shall be applicable to areas acquired pursuant to this subsection.”* Migratory Bird Hunting and Conservation Stamp Act (16 USC 718 [c])

*“For use as an inviolate sanctuary, or for any other management purpose, for migratory birds.”* Migratory Bird Conservation Act (16 USC 715d[2])

*“Areas of lands, waters, or interests therein acquired or reserved pursuant to this subchapter shall ... be administered ... to conserve and protect migratory birds in accordance with treaty obligations with Mexico, Canada, Japan and the Union of Soviet Socialist Republics, and other species of wildlife found thereon, including species that are listed ... as endangered or threatened species, and to restore and develop adequate wildlife habitat.”* Migratory Bird Conservation Act (16 USC 715i[a])

*“For conservation purposes any real property, or interest therein ... that has marginal value for*

*agricultural production; is environmentally sensitive; or has special management importance.”*

Consolidated Farm and Rural Development Act (7 USC 2002 [a])

*“It is the purpose of this chapter to promote ... the conservation of the wetlands of the Nation in order to maintain the public benefits they provide and to help fulfill international obligations in various migratory bird treaties and conventions with Canada, Mexico, Japan, the Union of Soviet Socialist Republics, and with various countries in the Western Hemisphere.”* Emergency Wetlands Resources Act (16 USC 3901 [b])

*“The following described public lands are hereby withdrawn from settlement, sale, location or entry under the general land laws, including the U.S. mining law, but not from leasing under the mineral leasing laws, to protect waterfowl production areas. This withdrawal will expire 50 years from the effective date of this order unless ... the Secretary determines that the withdrawal shall be extended.”* Public Land Order 7206 (June 24, 1996)

*“To protect waterfowl production areas.”* Public Land Order 6979 (May 25, 1993)

### NATIONAL WILDLIFE REFUGE SYSTEM MISSION

The mission of the System is to administer a national network of lands and waters for the conservation, management, and where appropriate, restoration of the fish, wildlife, and plant resources and their habitats within the United States for the benefit of present and future generations of Americans.

#### 1. DESCRIPTION OF PROPOSED USE: FARMING, GRAZING, AND HAYING

The district will continue upland management activities such as farming, grazing, and haying that are conducted under cooperative farming or special use permit by private individuals. Currently these economic uses are used as tools to manage habitat for wildlife.

Between 400 and 700 acres are farmed each year. Farming is conducted for the sole purpose of grassland restoration. Grazing with livestock is used as a tool for grassland and wetland management.

About two-thirds of the WPAs receive some type of grazing treatment. Grazing does not occur over the entire WPA but involves rotational grazing over portions of a WPA. Wetlands are the most common habitat grazed. Area grazed ranges between 7,000 and 9,000 acres annually over the past 5 years.

Approximately 450 acres are hayed annually. Haying is sporadically used as a grassland management tool to control invasive plants and prepare areas for upland restoration and prescribed burns.

The CCP will continue grassland restoration activities within the district. The amount of farming done will be in direct relation to the amount of land acquired and the availability of native seed. Over time, farming is expected to decrease as areas are restored to grassland. Cooperative farming activities will be compatible only at areas that do not have established stands of native grasses and forbs. Farming will allow the district to establish seedbeds relatively free of noxious plants, maximizing the likelihood that grassland restoration will be successful. Soybean is the crop generally used during farming.

The CCP will use grazing as a management tool for wetland and upland habitats. Specific acreages have not been identified in the CCP because habitat conditions within wetland and upland areas can change dramatically on a yearly basis due to precipitation and temperatures. An adaptive approach will be used to prescribe grazing treatments for habitats.

### Availability of Resources

The resources necessary to administer haying, grazing, and farming programs are sufficient at current staffing and funding levels. Haying, grazing, and farming programs are conducted through special use permits or cooperative farming agreements, which minimizes the need for staff time and district assets to complete work.

### Anticipated Impacts of the Use

Over a 5-year period, grazing has been conducted on 7,000–9,000 acres annually. While annual acreages have not been specified in the CCP, it is expected that future refuge grazing will fall into this range. Farmed acres will remain in the range of 400–700 acres per year, but will slowly decrease as uplands are restored to grass. Haying is anticipated to increase as a management tool to prepare for prescribed burns and control of invasive plants. Haying is expected to increase to about 600 acres.

Without management, wetland and upland habitat conditions will deteriorate due to long periods of rest. Cool-season invasive plants will likely increase and infest additional areas without the use of spring grazing. While all these activities disturb habitat and wildlife in the short term, long-term habitat and wildlife benefits outweigh these disturbances. Farming

will cause decreases in wildlife habitat availability; however, habitat conditions will improve following grassland restoration activities.

No cultural resources will be impacted. No impact to endangered species should occur.

### Determination

The use of haying, grazing, and farming as habitat management tools is compatible.

### Stipulations Necessary to Ensure Compatibility

- Monitor vegetation and wildlife to assess the effects of the management tools.
- Require general and special conditions for each permit to ensure consistency with management objectives.
- Restrict the use of vehicles and motorized equipment to the minimum necessary to conduct haying, grazing, and farming.
- Restrict farming permittees to a list of approved chemicals that are less detrimental to wildlife and the environment.
- Restrict haying to after August 1 to avoid disturbance to nesting birds unless the refuge manager deems it necessary to hay earlier to control invasive plants or restore grasslands.

### Justification

To maintain and enhance the habitat for migratory birds and other wildlife, some habitat manipulation needs to occur. Upland and wetland habitat conditions will deteriorate without the use of a full range of management tools. Migratory bird habitat and ecological diversity will decrease as habitat suitability declines. Exotic and invasive plant species will increase and habitat diversity will decrease if grazing practices did not continue at the WPAs. Farming will provide a means to restore degraded grasslands for the benefit of grassland-dependent species.

### Mandatory 15-year Reevaluation Date: 2022

## 2. DESCRIPTION OF PROPOSED USE: ENVIRONMENTAL EDUCATION AND INTERPRETATION

All of the district's WPAs are open to environmental education and interpretation in accordance with the Migratory Bird Hunting and Conservation Stamp Act. The district will provide opportunities for environmental education and interpretation. Environmental education will consist of activities conducted by district staff and volunteers. Interpretation will occur in less formal activities with refuge staff and volunteers or through exhibits, educational trunks, signs, and brochures. The lack of an outdoor recreation planner and the scattering of the WPAs across 14 counties will not allow the

district to adequately address education and interpretation opportunities. Based on the CCP, the district will hire an outdoor recreation planner and increase education and interpretation for all visitors through the following actions:

- Construct an additional accessible blind for people with disabilities and an interpretive trail in the eastern portion of the district.
- Construct entrance signs at 50% of the WPAs.
- Develop five WPAs as focus areas for education and interpretation.
- Develop watchable wildlife brochures.
- Construct additional interpretive panels to be placed along trails and at parking lots.

### Availability of Resources

Implementing the new facilities outlined in the CCP is closely tied to funding requests for RONS and SAMMS projects (see appendix O). Existing programs such as district signs and brochures will be updated with available resources.

### Anticipated Impacts of Use

All of the WPAs are open to environmental education and interpretation in accordance with the Migratory Bird Hunting and Conservation Stamp Act. Minimal disturbances to wildlife and wildlife habitat will result from these uses at the current and proposed levels. Some disturbance to wildlife will occur in areas frequented by visitors. There will be littering, minor damage to vegetation, and increased maintenance.

No cultural resources will be impacted. No impact to endangered species should occur.

### Determination

Environmental education and interpretation are compatible.

### Stipulations Necessary to Ensure Compatibility

- Prohibit vehicle access beyond parking lots.
- Prohibit permanent and overnight blinds.
- Develop trails and viewing areas that have minimal impact on wildlife and their habitats.
- Annually review environmental education and interpretation activities to ensure these activities are compatible.

### Justification

Based on biological effects described previously and in the EA and draft CCP, it is determined that environmental education and interpretation within the Rainwater Basin Wetland Management District will not interfere with or detract from the purposes for which the district was established.

Environmental education and interpretation are priority public uses listed in the National Wildlife Refuge System Improvement Act of 1997. By facilitating environmental education, district visitors will gain knowledge and an appreciation of fish, wildlife, and their habitats, which will lead to increased public awareness and stewardship of natural resources. Increased appreciation for natural resources will support and complement the Service's actions in achieving the purposes of the district and the mission of the Refuge System.

**Mandatory 15-year Reevaluation Date: 2022**

## 3. DESCRIPTION OF PROPOSED USE: WILDLIFE OBSERVATION AND PHOTOGRAPHY

All of the district's WPAs are open to wildlife observation and photography in accordance with the Migratory Bird Hunting and Conservation Stamp Act. The district will provide opportunities that support wildlife-dependent recreation. Wildlife observation and photography are facilitated by one hiking trail and two wildlife observation blinds.

The CCP will continue the above uses and add the following to improve wildlife observation and photography:

- Construct an additional accessible blind for people with disabilities and a hiking trail in the eastern portion of the district.
- Construct entrance signs at 50% of the WPAs.
- Develop a wildlife brochure/bird guide that identifies the WPAs, as well as seasons that offer exceptional wildlife observation and photography opportunities.
- Update existing interpretive kiosks.
- Develop five WPAs as focus areas for wildlife observation and photography.

### Availability of Resources

Implementing the new facilities outlined in the CCP is closely tied to funding requests for RONS and SAMMS projects (see appendix O). Existing programs such as district signs and brochures will be updated with available resources.

### Anticipated Impacts of Use

All of the WPAs are open to wildlife observation and photography in accordance with the Migratory Bird Hunting and Conservation Stamp Act. Minimal disturbances to wildlife and wildlife habitat will result from these uses at the current and proposed levels. Some disturbance to wildlife will occur in areas frequented by visitors. There will be littering, minor damage to vegetation, and increased maintenance.

No cultural resources will be impacted. No impact to endangered species should occur.

## Determination

Wildlife observation and photography are compatible.

## Stipulations Necessary to Ensure Compatibility

- Prohibit vehicle access beyond parking lots.
- Prohibit permanent and overnight blinds.
- Develop trails and viewing areas that have minimal impact on wildlife and their habitats.
- Annually review wildlife observation and photography activities to ensure these activities are compatible.

## Justification

Based on the anticipated biological effects described previously and in the EA and draft CCP, it is determined that wildlife observation and photography at Rainwater Basin Wetland Management District will not interfere with the habitat goals and objectives or purposes for which it was established.

Wildlife observation and photography are priority public uses listed in the Improvement Act. By facilitating these uses, visitors will gain knowledge and an appreciation of fish and wildlife, which will lead to increased public stewardship of wildlife and their habitats. Increased public stewardship will support and complement the Service's actions in achieving the purposes of the district and the mission of the Refuge System.

**Mandatory 15-year Reevaluation Date: 2022**

## 4. DESCRIPTION OF USE: RECREATIONAL FISHING

All of the district's WPAs are open to recreational fishing in accordance with the Migratory Bird Hunting and Conservation Stamp Act. The district will continue to provide for recreational fishing in accordance with state regulations.

Generally, fisheries do not exist at the WPAs. Wetlands in the Rainwater Basin are shallow, temporary or seasonal wetlands that normally are dry during the summer months. It is only during a series of excessively wet years that the wetlands support fish. Fish that have occurred in the wetlands are black bullhead and European carp. Boating with electric motors will be allowed.

## Availability of Resources

The current fishing program will be administered using available resources. The CCP does not call for the implementation of any new fishing programs.

## Anticipated Impacts of Use

Fishing and other human activities cause disturbance to wildlife. All of the WPAs will be open to recreational fishing in accordance with the Migratory Bird Hunting and Conservation Stamp Act.

## Determination

Recreational fishing is compatible.

## Stipulations Necessary to Ensure Compatibility

- Require that fishing follow state and federal regulations.
- Monitor existing use to ensure that facilities are adequate and disturbance to wildlife continues to be minimal.

## Justification

Based on the biological effects addressed previously and in the EA and draft CCP, it is determined recreational fishing will not interfere with the habitat goals and objectives or purposes for district establishment. Fishing is a priority public use as listed in the Improvement Act.

**Mandatory 15-year Reevaluation Date: 2022**

## 5. DESCRIPTION OF USE: RECREATIONAL HUNTING

The district will continue to allow hunting of all species according to state regulations. All of the WPAs are open to recreational hunting in accordance with the Migratory Bird Hunting and Conservation Stamp Act.

## Availability of Resources

Currently, sufficient resources are available to carry out the proposed recreational hunting program.

## Anticipated Impacts of Use

Some wildlife disturbance will occur during recreational hunting activities. All of the WPAs will be open to recreational hunting in accordance with the Migratory Bird Hunting and Conservation Stamp Act.

## Determination

Recreational hunting is compatible.

## Stipulations Necessary to Ensure Compatibility

- Require the use of nontoxic shot, in accordance with current regulations for migratory bird and upland game hunting.
- Prohibit vehicle access beyond parking lots.
- Prohibit permanent and overnight hunting blinds
- Prohibit the use of horses.
- Prohibit camping, overnight use, and fires.
- Require that hunting be in accordance with federal and state regulations.
- Promote sound hunting practices for hunter safety and quality experiences.
- Annually review recreational hunting activities to ensure these activities are compatible.

**Justification**

Hunting on Refuge System lands has been identified as a priority public use in the Improvement Act. Hunting is a legitimate wildlife management tool that can be used to manage populations. Hunting will harvest a small percentage of the renewable resources, which is in accordance with wildlife objectives and principles.

Based on the biological effects described previously and in the EA and draft CCP, it is determined that recreational hunting in the district will not interfere with or detract from the purposes for which the district was established or its habitat goals and objectives.

**Mandatory 15-year Reevaluation Date: 2022**

**6. DESCRIPTION OF PROPOSED PUBLIC USE: TIMBER HARVEST**

The district will continue timber harvest that is conducted under a special use permit by private individuals. Timber harvest will be used as a management tool to reduce the invasion of woody vegetation in grassland and wetland habitats.

**Availability of Resources**

The CCP does not propose any changes in timber harvest activities allowed with special use permits. The resources necessary to administer a timber harvest program will be sufficient at current staffing and budgetary levels.

**Anticipated Impacts of the Use**

Timber harvest will have short-term impacts by causing disturbance to wildlife and disturbance to ground cover. The removal of trees will reduce the effects of shading and robbing of soil moisture, and

thereby encourage warm-season grassland species to grow. The absence of trees will reduce nest and chick mortality for ground-nesting birds that are commonly preyed on by raptors, raccoon, and skunk.

**Determination**

The use of timber harvest (under special use permit) as a habitat management tool is compatible.

**Stipulations Necessary to Ensure Compatibility**

- Require general and special conditions for each permit to ensure consistency with management objectives.
- Restrict the use of vehicles and motorized equipment to the minimum necessary to conduct timber harvest.
- Require trees be cut to ground level and the stumps be treated with approved herbicide to prevent regrowth.
- Require slash to be removed or piled according to the manager's discretion.

**Justification**

Soil and climatic conditions are suitable to grow trees on any land left undisturbed. In addition, larger, mature trees that have timber value are the primary source of seed for new seedlings. Tree removal is vital to the maintenance of quality wetlands and uplands. The spread of saplings will (1) increase the costs of vegetation management that are associated with prescribed burning and (2) reduce haying as a management option. Invasive, cool-season grasses will be commonly associated with shaded woodland areas. Bird predators such as owls, red-tailed hawk, raccoon, and skunk will benefit from shelterbelts and volunteer trees.

**Mandatory 15-year Reevaluation Date: 2022**

Submitted to: Gene Mack  
 Gene Mack, Project Leader      Date  
 Rainwater Basin Wetland      8-20-07  
 Management District, NE

Approved by: Richard A. Coleman 8/21/07  
 Richard A. Coleman, PhD,      Date  
 Assistant Regional Director  
 Region 6, National Wildlife  
 Refuge System, CO

Reviewed by: Rod Krey 8-21-07  
 Rod Krey, Refuge Supervisor      Date  
 Region 6, National Wildlife  
 Refuge System, CO



# Appendix C

## *Preparers*

This CCP is the result of the efforts by members of the planning team for the Rainwater Basin Wetland Management District.

<i>Name</i>	<i>Title</i>	<i>Agency</i>
Mike Artmann	GIS specialist	USFWS
Susann Cayouette	administrative officer	USFWS
Jeff Drahota	wildlife biologist	USFWS
Mark Ely	GIS specialist	USFWS
Bernardo Garza	planning team leader	USFWS
Kyle Graham	private lands biologist	USFWS
Steve Karel	refuge operations specialist	USFWS
Brad Krohn	biological technician	USFWS
Brice Krohn	supervisory range technician	USFWS
Ted LaGrange	wetland program manager	NGPC
Gene Mack	project leader	USFWS
Mindy Meade	private lands biologist	USFWS
Steve Moran	RWBJV coordinator	USFWS
Paul Nelson	maintenance worker	USFWS
Mark Pfof	biological technician	USFWS
Ryan Reker	GIS specialist	USFWS
Ronnie Sanchez	deputy project leader	USFWS
Richard Schroeder	wildlife biologist/ecologist	USGS-BRD
Mark Vrtiska	waterfowl program manager	NGPC
Bruce Winter	prescribed fire specialist	USFWS



# Appendix D

## *Public Involvement*

The Service began preparation of this “Comprehensive Conservation Plan for the Rainwater Basin Wetland Management District” with the preplanning process in September 2005. On November 10, 2005, the Service contacted state and tribal representatives to invite them to participate in the planning process for the district’s CCP. The planning team was formed during the kickoff meeting in Kearney, Nebraska, December 6–7, 2005, with the following team members: Service personnel from the district and the regional office; representatives from the NGPC and RWBJV; and personnel from USGS–BRD (see appendix C, preparers).

### **PUBLIC SCOPING**

The notice of intent to prepare the CCP for the Rainwater Basin Wetland Management District was published in the *Federal Register* on November 30, 2005. The public scoping process was completed in December 2005. Four public open-house meetings were held from 7:00 to 9:00 p.m. during consecutive nights throughout the Rainwater Basin, as follows:

- Kearney, NE; December 5, 2005
- York, NE; December 6, 2005
- Clay Center, NE; December 7, 2005
- Holdrege, NE; December 8, 2005

Notification of dates and times of the public open house meetings was distributed through press releases and a planning update mailed out to everyone in the CCP mailing list.

Of the 51 people who attended these meetings, 38 were non-Service individuals. Written comments were received from 17 individuals. Comments received identified biological, recreational, and economic concerns about management of the district.

Those who attended provided both written and oral comments. They were informed that comprehensive planning was an open process and they could submit their comments at any time and by any means (for example, by letter, telephone, and email) until the time the CCP was final. Additional written comments were received by the planning team via mail.

Over the course of preplanning and scoping, the planning team collected available information on the resources of the district and the surrounding areas

as well as what are the issues facing the district and of importance to the public. This information is summarized in chapter 2 (“The District”) and chapter 3 (“District Resources and Description”).

The planning team revisited the purposes for which the district was created and drafted issues and qualities lists, as well as the vision and goals, for the district during a workshop held in Kearney, Nebraska, October 13–14, 2005.

Between February and April 2006, the planning team developed a range of alternatives that included two alternatives for management of the district. A subsequent assessment of each alternative’s environmental consequences guided the team in choosing the alternative that would best fulfill the purposes, vision, and goals for the district. Once the Service identified the proposed action, the planning team developed the objectives, strategies, and rationales for each of the goals set out by the district.

### **PUBLIC REVIEW OF THE DRAFT CCP**

The Service released the draft CCP and EA for a 30-day public comment period starting July 18, 2007. During this public comment period, the planning team held a public meeting in Hastings, Nebraska (July 31, 2007, from 12:00 to 8:00 p.m.). An announcement of this meeting and the release of the draft CCP and EA for public comment was published in the *Federal Register* on July 18, 2007, as well as in local media. Six members of the public, including a local television station reporter and a congressional aid, attended the public meeting.

The public comment period closed August 18, 2007. Two printed letters, a personal communication with the project leader, three personal comments during the public meeting, and an email message were the only comments received from the public. The following summarizes the comments received and the Service’s responses.

### **COMMENTS AND RESPONSES**

**Comment 1**—The district staff is doing a much better job than in the early part of the 1990s. There are no problems or concerns.

**Response 1**—(None needed.)

**Comment 2**—The grazing and farming programs (as a preparatory technique for restoration to native habitats) administered by district staff have improved waterfowl habitat by changing the plant community from less desirable to more desirable species (for example, from cattails to barnyard grass and reduction in the commonly occurring reed canarygrass and other noxious weeds) at several WPAs.

**Response 2**—The district staff will continue to carry out these and other habitat management techniques that help restore and maintain the habitat conditions required by the migrating and resident wildlife species found in the Rainwater Basin.

**Comment 3**—Food plots should not be considered among the district's habitat management activities because these plots benefit an introduced species (pheasant), plus there is already plenty of leftover grain on lands adjacent to the district's WPAs for pheasant to feed on.

**Response 3**—The district staff will not plant food plots during the implementation of the CCP because food in the form of grain is not a limiting factor in the life or ecology of pheasant or any other species within the Rainwater Basin. More limiting is the diversity and abundance of natural food associated with native plant communities—native plant seeds and insects.

**Comment 4**—The CCP should include information about how wetlands managed by the district help recharge groundwater in the Rainwater Basin.

**Response 4**—There is not adequate information to fully determine the role played by the district's wetlands in the recharging of groundwater in the Rainwater Basin. The CCP addresses the need for further research into the role that the district's wetlands have on groundwater recharge and other ecological benefits to the environment of the Rainwater Basin, such as the quantity and quality of surface and subsurface water resources.

**Comment 5**—Comments provided during the public review period should receive serious consideration and not be not merely a step in an already-completed process.

**Response 5**—All comments provided during the planning process have received serious consideration and have been instrumental in the development of the vision, goals, objectives, and strategies in the final CCP. The Service has adhered to the guidelines and spirit of the NEPA in fully developing and equally analyzing and considering a range of alternatives for the management of the resources of the district. The development and analysis of the range of alternatives and selection of a preferred alternative for this final CCP has not been predecisional.

**Comment 6**—What is to prevent future managers from interpreting this CCP to fit their goals? The

CCP may be changed, creatively interpreted, or ignored if it suits the goals of the decision maker.

**Response 6**—This 15-year CCP is a congressionally mandated plan that district and refuge managers must adhere to when making management decisions. The Refuge System's administrative structure will ensure continuity of management and compliance with goals and objectives in this CCP (for example, through oversight by refuge supervisors).

**Comment 7**—Alternative A is the only realistic option. Alternative B offers lofty goals that are unrealistic with available funding and management capabilities. A third option should be introduced in which alternative B management would be fully implemented and further improvements would be undertaken if funding becomes available. Adding additional land without having additional funding will lead to further degradation of the current WPAs and the degradation of new properties acquired. This CCP means nothing without the long-term commitment of funding and personnel to carry out the plan.

**Response 7**—Through this planning process, the Service has determined that carrying out alternative B with its aim to improve and expand management capabilities through partnerships is realistic and feasible. Addition of more land to the district will only occur if the district staff is capable of managing it in accordance with the goals and objectives in this CCP.

**Comment 8**—There is concern that some Service personnel are not concerned with production of waterfowl, but only with migrating waterfowl, particularly during spring migration. The result of this CCP will depend on current and future interpretation and actions, and those will change with changes of personnel.

**Response 8**—The district is located on the critical bottleneck of the hourglass-shaped Central Flyway. All bird species migrating through the flyway converge in the Rainwater Basin during spring and fall migrations—the basin provides a vital resting and staging point for millions of migratory birds. Many wetland habitats in the basin have been converted to urban and agricultural landscapes. The remaining wetlands, both within and outside the district, are now required to provide all the necessary resting and feeding grounds for all the migrating birds that converge in the Rainwater Basin. Thus, a main concern for the Service is to provide the necessary habitat conditions required by migrating birds. The Service is not solely concerned with providing migrating habitat, but given the overall habitat conditions present throughout the basin, the district staff gives priority to ensuring migrating birds can find the best possible habitat during their migrations. In addition, the district staff works incessantly to restore upland habitats (nesting habitat). As noted in response 6, this 15-year

CCP is a congressionally mandated plan that district and refuge managers must follow when making management decisions, and there is administrative oversight to ensure compliance.

**Comment 9**—The government needs to fully fund “payments in lieu of taxes” on all lands to an amount equal to what a private landowner would have paid at the higher of either agricultural production value or recreational value. Paying the full “in lieu of taxes” could help fund better local law enforcement.

**Response 9**—“Payments in lieu of taxes” are paid by the Service through the provisions of the Revenue Sharing Act. However, the Service’s ability to make these payments depends on designated funding appropriated by Congress. The Service makes the best possible use of the resources that Congress allocates to the agency. The Service agrees that payment in full “in lieu of taxes” could translate into better local law enforcement.

**Comment 10**—The federal government needs to be a leader in funding and managing its own properties, which means funding the management needs of all currently owned lands in the district.

**Response 10**—As noted in response 9, the Service prioritizes management needs and makes the best possible use of the resources that Congress allocates to the agency.

**Comment 11**—Alternative B does not address how funding will be provided other than with partnerships, local cooperation, and volunteerism. This implies that the Service is transferring its funding and staffing to the local communities, but does not allow local decision making about the management actions. This will also decrease funding and staffing for privately managed wildlife habitat.

**Response 11**—Only the Service has decision-making authority for management within the district; the Service cannot transfer this responsibility or authority. The Service is committed to providing adequate funding and personnel to manage lands in the district. This includes finding new, innovative solutions to current and future management, funding, and personnel challenges—partnerships can be some of the solutions. The Service focus on partnerships aims to fulfill the goals and objectives in this CCP. Partnerships that the district staff seek and maintain will be mutually beneficial, depending on the needs of each individual partner.

**Comment 12**—The Service needs to develop and maintain contact information for adjoining landowners and tenants, plus residents within 2 miles of district properties. The Service needs to annually send each of these contacts a letter that outlines district management plans for the properties near them, how to contact the Service if there is a problem arising from such management as burning or livestock grazing. When grazing is being used, the

Service needs to provide the local sheriff’s office and adjoining farmers and residents with the livestock owners’ contact information. The Service needs to use the local and county newspapers, as well as regional and state newspapers, to promote district management and inform the public.

**Response 12**—This CCP (a public document readily available in printed format at the district office and as a ready-to-download PDF document at the Service’s region 6 refuge planning website) outlines the different techniques used by the district’s staff to manage lands administered by the Service. The Service will continue to contact fire departments and other appropriate entities to inform them of management activities on Service lands. The district staff, within its budgetary and personnel capabilities, will attempt to develop better means to keep its neighbors informed. This task will be more easily accomplished once the Service can allocate more personnel to the district.

**Comment 13**—Nebraska is sometimes known as the “Tree Planter State”—it is home of Arbor Day. The Service needs to recognize that this is part of the local culture and heritage and, therefore, preserve planted trees, shelterbelts, windbreaks, and border trees. These treed areas are habitats that provide food, shelter, and nesting and resting areas for nonwaterfowl migratory birds. There could be a win-win situation by protecting large trees, along with carrying out early control of seedling trees instead of destroying the large-tree habitat that takes 30–60 years to replace.

**Response 13**—Most of the land in the Rainwater Basin is in private property and contains vast treed habitats that ensure the preservation of the local culture and heritage. For the district’s WPAs, the Service’s responsibility is to manage for the purpose for which the district was established—for waterfowl and other migratory birds—by restoring wetland and grassland conditions needed by these species. The majority of bird species that migrate through the Central Flyway depend on the mosaic of grasslands and wetlands to provide them with sustenance and rest amidst their long flights. While the current mosaic of extended areas of trees used in shelterbelts, windbreaks, and border trees is conducive to some native and introduced wildlife species, and even to some migrating Neotropical birds, it is not adequate nesting and feeding habitat for most migrating species. Prior to human settlement of the Rainwater Basin, the vastly predominant mosaic of habitats in this area were prairie grasslands dotted with numerous complexes of wetlands. Treed areas were sparse and mostly confined to the margins of rivers and creeks. Shelterbelts provide habitat for some species of wildlife; however, shelterbelts and other trees and shrubs decrease the size of grassland blocks and result in fragmented habitats. Shelterbelts are unnatural in grasslands and provide habitat for both

avian and mammalian predators. Recent studies have shown that many grassland-nesting birds and upland-nesting waterfowl either avoid areas adjacent to trees or have lower nest success due to predation. Furthermore, treed areas tend to deplete surface water from wetlands and harbor migratory bird predators. Service activities at the district focus on restoration of presettlement habitat conditions in the lands administered by the district.

**Comment 14**—Large trees provide habitat for the bald eagle, which is America’s national bird and symbol and is highly regarded in the culture of Native Americans. It is important for the youth and others who live in or visit this area to see this majestic bird alive and free on public land, not just as a picture in a book. Observation of bald eagles is one of the thrills cited by visitors to this area during migration. The bald eagle is being driven from these public lands by current and proposed management activity; it takes many years to replace large-tree habitat.

**Response 14**—As described in response 13, the purpose for which the district was established is to manage WPAs for waterfowl and other migratory birds by restoring wetland and grassland conditions needed by these species. Bald eagles benefit from the presence of the migratory birds that rest and feed in the Rainwater Basin. Bald eagle populations throughout the United States have grown steadily to the point that this species has been taken off the endangered species list. This majestic American symbol uses and will continue to use the basin so long as there are adequate food resources (migrating waterfowl and other migratory species) that sustain its individuals. Bald eagles make use of all adequate habitats, whether public or private lands. There are vast treed areas within the basin that the bald eagles will use for nesting and roosting even after the Service has restored to grassland and wetland all the lands it administers in the district.

**Comment 15**—There is value in the observation of young wildlife; young ducklings and goslings are an attraction to bird watchers and hunters. The Moger WPA produced wood ducks before their nesting trees were removed.

**Response 15**—As described in response 14, wildlife make use of habitats in the Rainwater Basin so long as they provide adequate food, nesting, loafing and resting resources, whether on public or private lands. Geese, while able to use treed areas for nesting, is primarily a prairie-wetland complex species that will continue to use the basin after the Service has restored the lands it administers to a prairie-wetland complex. To keep trees around wetlands, the Service would need to shift its mission, to one of providing nonhistorical habitat for a species such as the wood duck that was not numerous in grasslands of the Central Flyway. All this would be at the expense of those species that historically relied on the open wetland-grassland landscape.

**Comment 16**—The Service must commit itself to meeting the goals and expectations of partners. In the past, partnerships have meant local funding for Service responsibilities. Other government or nongovernmental groups might not provide funding or other assistance if there is no way to be assured that the Service will maintain what has been provided. Past tree plantings funded by nonfederal groups were destroyed when management changed.

**Response 16**—The Service is committed to meeting the goals and expectations it enters into with its partners. The definition that the Service uses to describe a partnership is, “a contract or agreement entered into by two or more individuals, groups of individuals, organizations or agencies in which each agrees to furnish a part of the capital or some in-kind service, such as labor, for a mutually beneficial enterprise.” As noted in response 11, the partnerships that the district staff will seek and maintain will be mutually beneficial, depending on the needs of each partner. Partnerships will aim to fulfill the goals and objectives in this CCP.

**Comment 17**—When partnerships and cooperation are needed, the partners and cooperators must be full partners, where their concerns or ideas are given full consideration. In the past, cooperation with the Service has often meant that a local landowner or government agrees to do something for little or no benefit to the landowner or government.

**Response 17**—Cooperation is essential to effective partnerships; thus, response 16 applies here. The Service considers, respects, and pays attention to the concerns and ideas expressed by its partners and cooperators. In fact, the partnership-driven management scheme for the district finds its roots in ideas and comments provided to the Service by its neighbors, partners, and cooperators.

**Comment 18**—The Service needs to give greater consideration to the concerns of local government and input provided by community leaders. People from the county where the property is located must be given a greater voice, and the Service must yield to local needs if a mutually acceptable compromise cannot be found.

**Response 18**—The Service gives full consideration to the concerns of local government and community leaders. However, neither the Service or the district can deviate from their congressionally mandated mission and purpose. Furthermore, as stated in response 11, only the Service has decision-making authority for management within the district; the Service cannot transfer this responsibility or authority.

**Comment 19**—The Service needs to involve local school and youth groups in construction and maintenance activities at the district. This would develop feelings of pride and ownership in the persons involved. The Little Blue NRD has used this as a method to lessen vandalism on some areas.

**Response 19**—The district staff will seek ways to carry out this helpful idea.

**Comment 20**—The Service needs to contact local schools and offer to provide them guides for field trips to areas in the district.

**Response 20**—The district staff will seek ways to provide local schools with information about school visits and use of the district's facilities. The implementation of this idea will be facilitated when the district is allocated further personnel, especially an outdoor recreation planner.

**Comment 21**—Drought, wet years, burning, grazing, insects, and diseases are some of nature's methods of control that need to be respected. Multi-year drought on private lands has had an impact on the infestation of cattails and rushes. The Service needs to allow basins to go through a drought cycle of more than a few months—many irrigated basins have not undergone a multiyear drought since being purchased. The Service should rotate water-pumping in basins to provide multiyear drought cycles, but not all basins should be left dry at the same time. This management will provide more natural conditions and be more cost effective.

**Response 21**—Drought and extended periods of dry conditions are natural phases of a healthy wetland. However, the large-scale loss of wetlands throughout the Rainwater Basin has made it more vital that the remaining 10% contain water at a frequency higher than what historically occurred. It is part of this CCP to increase the number of WPAs that have water-pumping capabilities; those areas that have had annual water-pumping can be allowed to go through some year-long dry conditions. The added pumping capabilities will increase water management flexibility, allowing a better mix of wet and dry wetlands during migration.

**Comment 22**—Five percent of all lands owned by the Service in each township should be maintained specifically for local habitat interests. The availability of food plots, trees, shrubs, or legumes would diversify the habitat for other wildlife that rely on wetlands. This would provide habitat suitable for upland birds, deer, turkey, songbirds, and raptors.

**Response 22**—As noted in response 13, the purpose for which the district was established is to manage WPAs for waterfowl and other migratory birds by restoring wetland and grassland conditions needed by these species. Providing 5% of the district's land for habitat for nontrust species would mean a 5% decline in habitat for migratory waterfowl and other migratory birds. Based on this CCP, management will continue to provide habitat for upland birds and songbirds. The agricultural lands and shelterbelts that currently dominate the Rainwater Basin landscape provide habitat for deer, turkey, raptors, and other woodland birds.

**Comment 23**—The district's buildings and offices should be built on or adjoining an existing WPA. The sites should be landscaped using plants that occur on the adjoining WPA, and the grounds should be maintained exactly like the WPA. The Service needs to design the facility for use as an outdoor classroom for the public. The Service should consider having facilities built by private owners and leased to the Service.

**Response 23**—These facility comments are very helpful and the Service will consider them during design and location of new district buildings. Construction and location will be influenced considerably by the source and intent of the funds received. For example, the funding source could require the district to co-locate with other Service offices or with another agency.

**Comment 24**—The CCP proposes spending almost \$2 million on equipment that can only be used on a seasonal basis and is best used by skilled operators. It would be better to use contractors, who can complete projects quickly and efficiently. Some small equipment items on the CCP list should be leased or hired on an as-needed basis.

**Response 24**—The district currently uses contractors and rental equipment as much as possible. However, some equipment is needed on a frequent or immediate basis, which makes it practical to own the equipment. In addition, the seasonality of use for some equipment makes it difficult to locate or rent that piece of equipment because it is in high demand at the same time the district would need to lease it.

**Comment 25**—Given the size of the district and the number of WPAs, one or two law enforcement people would not be enough to provide adequate coverage. Redefining the working hours of current and additional employees would help provide a presence during weekends and evening hours to discourage lawbreaking. These employees could call authorities to make the arrests.

**Response 25**—Even the additional law enforcement staff (1) specified to carry out this CCP would not be adequate to address all law enforcement situations throughout the district. All staff are watchful and report violations they observe while doing their normal work in the district. To redefine work hours of current employees would mean to reduce the amount of staff time available to do their main work. For example, a maintenance person who is redirected to patrol areas would not be available to do maintenance. It would also increase the probability of that individual being involved in a life-threatening situation that he or she was not hired for or trained to respond to.

**Comment 26**—The Service needs to provide an honest comparison between the farming income and "new" recreational income. Areas acquired have a history of use by hunters and bird watchers. The

income from new visitors would need to surpass the income from current users and agriculture. New income for the area is welcome, but not if there is a greater loss to current income; it would be better to maintain the current income and be able to attract additional income.

**Response 26**—The income data and projections in the CCP is from a recent socioeconomic report. The Service used a consulting company—with ample experience in the research and analysis of socioeconomic data—to assess the economic impact of the Service’s activities in the Rainwater Basin. The data used in the subsequent socioeconomic analysis is current, accurate, and the best available for the projections stated in the report.

**Comment 27**—The decline of habitat for upland game has reduced the number of hunters using these areas in the fall. The spring goose season has helped offset this change, but does not have the magnitude or duration that upland hunting once had.

**Response 27**—The majority of lands acquired by the Service within the Rainwater Basin was marginal agricultural land containing drained or altered wetlands, which the Service converted to grassland and wetland. The number of hunters using these areas has increased since the land was converted to wildlife habitat. However, the overall decline in upland game habitat over the past decades has reduced the number of hunters in the basin during the fall. This overall decline in upland game habitat has increased the need for district lands to provide recreational opportunities.

**Comment 28**—The CCP does not address the carbon released by the burning and rotting of mature trees removed from borders and windbreaks. Many of these trees are 30–70 years old and have been storing carbon for that period of time. Sediment and tree removal disturb the soil and cause the release of carbon dioxide into the atmosphere. Disking, whether in a farmer’s field or in a wetland, exposes the soil for the release of carbon dioxide; it seems that there would be a greater release of carbon when sediments containing large amounts of organic matter is dug up and exposed to the air.

**Response 28**—The carbon balance or cycle is a complicated matter. Although burning and rotting of mature trees do release carbon, these are natural events since no tree lives forever. The conversion of tilled land into nontilled grassland does sequester carbon. Although reducing the amount of carbon emissions is a concern of the district, as well as every individual, it is not the purpose for which this district was established. The Service recognizes that all management practices have pros and cons that need to be considered.

**Comment 29**—Minimal maintenance roads provide valuable access to hunters and nature watchers, as well as providing necessary means for farmers to

move between fields. Because these roads are less traveled, they are often safer and more direct routes for farmers moving large equipment. While a couple of miles added to the travel distance (after closure of these types of roads) for an automobile may be a minor inconvenience to some, the added distance for slow-moving equipment would impact local farmers.

**Response 29**—Roads being sought for closure by the district are roads that have outlived their usefulness. They are commonly characterized by little or no maintenance and are located within wetlands. Often the request to close a particular road is initiated by adjoining landowners who are tired of being asked to pull stuck vehicles during hunting season. The process for closing any county or township road involves requesting permission from the appropriate government, providing public notice, and conducting a hearing on the matter. The district follows this process. In general, a road remains open if the local people are opposed to closing it. Roads closed to the public on district lands may have arrangements made to give adjoining landowners farming access, if necessary.

**Comment 30**—Efficient and beneficial use of water resources is important to all of us. However, pumping water for wildlife as well as farm irrigation can impact roads and adjoining properties of this basin. The Service’s description of water pumping in wetlands as irrigation would likely not meet Nebraska’s definition of irrigation. Currently, rights to underground water do not exist in Nebraska; the Little Blue NRD has initiated the process to change Nebraska law to provide for wildlife uses of underground water. In areas of conflict between surface rights holders and groundwater users, the surface water rights have priority. This CCP states that pumping is done only where the Service controls the entire wetland. However, residents across the road from Moger WPA experience flooding from heavy rains that flood the wetland. In addition, county roads on the south and west sides of the WPA are flooded by heavy rains. Therefore, any water the Service pumps into the Moger WPA adds to the flood potential. It is the goal of the local residents not to have irrigation water run off of their property and to be able to use most if not all of a normal rain.

**Response 30**—Pumping of any groundwater can have an impact on roads and adjoining properties under certain conditions. The district intends to pump enough water into its wetlands to provide migratory bird habitat, but still leave enough water storage area for normal rains and snowmelt. Since waterfowl and shorebirds prefer shallow water, only a small portion of the wetlands’ storage capacity is filled by pumping. However, heavy storm events can cause flooding. The district works with county officials and adjoining landowners to resolve such concerns; within the last decade, the district has only had a couple of complaints of this type. The more common complaint is flooding of fields and roads

caused by cornstalks being washed off the fields and plugging road culverts, keeping runoff from reaching wetlands.

**Comment 31**—Many of the invasive plant species have flourished under the management practices of the last 45 years. These plants were not a problem when the Service purchased the property, but became a problem due to lack of management. In the early years of the district it was thought that no management was needed, but it is now recognized that was not true.

**Response 31**—As more information is obtained through the years, management and objectives have changed. In recent years, the adaptive resource management approach has enabled the Service to evaluate and make adjustments on a timely basis.

**Comment 32**—In the past, the Service has resisted the efforts of local groups to build observation towers and blinds; everyone needs to work together.

**Response 32**—About 10 years ago, the district worked with one local group to build an observation blind on Massie WPA. That group maintains the blind for the benefit of wildlife enthusiasts who visit the area. As noted in response 11, only the Service has decision-making authority for management within the district; the Service cannot transfer this responsibility or authority. The Service cannot allow local groups to freely build towers and blinds that are not compatible with the Service's mission. For example, a blind that is restrictive in accessibility to all the public would not be compatible.

**Comment 33**—The pumping of warm water in January and early February led to an increased early migration under more severe conditions and that coincided with the most severe fowl cholera outbreaks. When there is no open water due to drought or cold weather, pumping creates unnatural open water and leads to more overcrowding. Natural adaptation to seasonal changes are a better plan.

**Response 33**—Natural adaptation to seasonal changes is a better plan, but that is unrealistic because of wetland losses within the Rainwater Basin, as well as throughout the Central Flyway. Natural runoff that normally reached wetlands has declined due to alterations within watersheds. Water pumping is currently the only way to replace the natural runoff. There is no known direct relationship between water temperature and avian cholera. Studies within the Rainwater Basin have documented that the bacteria does not remain in the water or soil, but is brought into the area each spring by migratory birds. The rate that water is delivered into a wetland and the area over which the water spreads out allows that water to quickly cool to the natural temperature of the wetland. It takes only one night of freezing temperatures to reduce pumped water to the temperature of the wetland.

**Comment 34**—It is surprising that prairie dogs would be mentioned in this plan because it seems that management is centered on waterfowl that are hunted and that other species are tolerated at best. There is no problem with prairie dogs on this area so long as they do not impact local landowners.

**Response 34**—The Service has a responsibility to manage all types of native wildlife. Although the emphasis is on waterfowl and migratory birds, management is done to help sustain all types of wildlife that naturally occur on district lands, including prairie dogs.

**Comment 35**—The Service should respect the needs and views of citizens, landowners, and governments in the communities where the WPAs are located. In addition, the Service needs to earn respect through timely, honest, and open communication with these citizens, landowners, and governments about district plans and actions.

**Response 35**—The Service agrees with this comment.

**Comment 36**—There is interest in the recharge of aquifers; if the Service can include recharge of aquifers in the plan, there may be related money to support district management.

**Response 36**—As noted in response 4, the district staff does not have adequate information to fully determine the role played by the district's wetlands in the recharge of aquifers (groundwater) in the Rainwater Basin. This CCP addresses the need for further research into the role that the wetlands managed by the district have on groundwater recharge and other ecological benefits to the environment of the Rainwater Basin such as the quantity and quality of surface and subsurface water resources.

**Comment 37**—The draft CCP and EA represents a continuing violation of the NEPA and section 7 of the Endangered Species Act due to the Service not doing the following: (1) preparing an environmental impact statement; (2) engaging in intra-Service consultations on the sport hunting program for the district and the Refuge System; and (3) using data from nonconsumptive and nonhunter surveys provided by the Humane Society. An environmental impact statement should be prepared because of sport hunting and overall recreation programs.

**Response 37**—Some citizens are concerned with hunting on units of the Refuge System. The Refuge System is guided by laws enacted by Congress and the president, as well as policy derived from those laws. The Improvement Act identifies hunting as one of six, priority, wildlife-dependent, recreational uses to be facilitated when compatible with the mission of the Refuge System and the purposes of any of its units. Hunting is consistent with the purposes of the district. The district manages lands acquired with Federal Duck Stamp and other funds as WPAs.

Because the source of funding for land acquisition is from Duck Stamps, the district's WPAs are open to public use including hunting. While the units of the Refuge System are managed first and foremost for wildlife, the focus is on perpetuating populations, not individuals. Hunting does adversely affect individual animals, but is allowed when it will not threaten perpetuation of the population being hunted.

The district has engaged in intra-Service, section 7 (of the Endangered Species Act) consultation with the Service's ecological services office in Grand Island, Nebraska, to ensure that the actions delineated in the CCP do not adversely affect any federally listed species. The conclusion made by the ecological services office is that the implementation of the CCP will not adversely affect federally listed species in the Rainwater Basin.

The preferred alternative (CCP) was not a major federal action that would significantly affect the quality of the human environment within the meaning of section 102(2)(C) of the NEPA. Accordingly, the preparation of an environmental impact statement was not required.

**Comment 38**—The Service should mow instead of using cattle to manage vegetation. Although grazing could help waterfowl, grazing has caused nothing but disaster.

**Response 38**—Like all management practices, grazing can produce results that cannot be achieved by other management actions. And like other practices, it must be used appropriately. The district staff has found, through years of observations, that livestock grazing can target specific plant species. Grazing that coincides with the beginning growth on undesirable plant species has caused that species to be overgrazed and reduced in number and health. The result is that other, preferred plant species can better compete. The more common use of grazing within the district is to control reed canarygrass. If left unmanaged, this species can displace all other plant species, resulting in a monotypic stand. For native, seed-producing plants to begin to return, the canarygrass has to be severely grazed; that level of grazing results in bare soil and torn-up roots. To the casual observer, the management practice appears to be a disaster. Unfortunately for the observer, the benefit is not seen until the next year's growing season.

## PLANNING MAILING LIST

The following mailing list was developed for this CCP's planning effort.

### FEDERAL OFFICIALS

U.S. Senator Chuck Hagel, Washington DC;  
Area director, Lincoln, NE

U.S. Senator Ben Nelson, Washington DC;  
Area director, Lincoln, NE

U.S. Representative Adrian Smith, Washington DC;  
Area director, Grand Island, NE

### FEDERAL AGENCIES

Bureau of Reclamation

National Park Service—Homestead National Monument, Beatrice, NE; Lewis and Clark National Trail, Omaha, NE

U.S. Army Corps of Engineers; Kearney, NE

USDA, Forest Service; Chadron, NE

USDA, Natural Resources Conservation Service in Nebraska—Holdrege Service Center, Grand Island Service Center, Clay Center Service Center, Hastings Service Center, Franklin Service Center, Elwood Service Center, York Service Center, Aurora Service Center, Geneva Service Center, Wilbur Service Center

USFWS in Nebraska—Ecological Services, Grand Island, NE; Crescent Lake NWR, North Platte NWR, Fort Niobrara NWR, Valentine NWR

USFWS—Northern Prairie Waterfowl Research Center

USFWS—regional offices of regions 1–7 and 9; National Conservation Training Center

USGS—Fort Collins Science Center, Fort Collins, CO; Northern Prairie Research Center; Jamestown, ND

### NEBRASKA STATE OFFICIALS

Senator Greg Adams, York  
Senator Ray Aguilar, Grand Island  
Senator Carroll Burling, Kenesaw  
Senator Tom Carlson, Holdrege  
Senator Annette Dubas, Fullerton  
Senator Ray Janssen, Nickerson  
Senator Joel Johnson, Kearney  
Senator Russ Karpisek, Wilber  
Senator Chris Langemeier, Schuyler

### STATE GOVERNMENT

NDEQ, Lincoln  
Nebraska Corn Board, Lincoln  
Nebraska Department of Agriculture, Lincoln  
Nebraska Department of Natural Resources, Lincoln  
Nebraska Department of Roads, Lincoln  
Nebraska Department of Water Resources  
Nebraska Emergency Management Agency, Lincoln  
Nebraska Environmental Trust, Lincoln  
Nebraska Forest Service, Lincoln  
Nebraska State Historical Society  
Nebraska State Historic Preservation Office  
NGPC, Lincoln

### LOCAL GOVERNMENT

Buffalo County Weed Board  
Central Nebraska Public Power Irrigation District

Central Platte NRD  
 Clay County Highway Department  
 Fillmore County Roads Department  
 Gosper County Weed Control  
 Hamilton County Board of Supervisors  
 Little Blue NRD  
 Lower Loup NRD  
 Lower Niobrara NRD  
 Meat Animal Research Center  
 Phelps County Road Department  
 Seward County Highway Department  
 Tri-Basin NRD  
 Upper Big Blue NRD

### **LOCAL ORGANIZATIONS AND BUSINESS AND CIVIC GROUPS**

The 1980 Anawalt Family Trust  
 Abengo Energy  
 A C Feedyards Inc.  
 Agnes E. Johnson Farms  
 Audubon Society  
 Boyd Kaiser Trust  
 BSL LLC  
 Burt Partners  
 Clarence W. Schmidt Trust  
 Clay Center Critter Care  
 Crane Meadows Nature Center  
 C & R Hendrickson Farms Inc.  
 Cudmore-Kneifl Construction Company Inc.  
 Daniel R. Stengel Revocable Trust  
 Dannehl Farms Inc.  
 Darleen Nielsen Trust  
 David B. Huber Revocable Trust  
 David High Farms Ltd.  
 Donald D. Lovegrove Trust  
 Donna Linder Trust  
 Dorothy I. Ebert Estate  
 Ducks Unlimited  
 Earl W. Frazier Revocable Trust  
 Flinthill Farms Ltd.  
 Gene Lundeen Inc.  
 Gladys W. Scharmann Trust  
 H-D Management Company  
 Kathleen M. Swartz Revocable Trust No. 1  
 Lauer Ag Inc.  
 Lazy T Milliron Inc.  
 Lipovsky Farm Corporation  
 L & K Irrevocable Living Trust  
 Marvin W. Volzke Trust  
 Max A. Gemar Family Trust  
 M & C Stadler Inc.  
 ME LLC  
 The Nature Conservancy  
 Nebraska Cattlemen  
 Nebraska Environmental Trust  
 Nebraska Farm Bureau  
 Nebraska Farmer  
 Nebraska Groundwater Foundation  
 Nebraska Hunters Connection  
 Nebraska State Historical Society  
 Nielsen Family Farms LLC

Ostgren Construction Company  
 Pheasants Forever  
 Playa Lakes Joint Venture  
 Prairie Plains Institute  
 Rainwater Basin Joint Venture  
 R Lazy K Inc.  
 R L Kaliff Ranch Company  
 Rosetta Nelson Family Farms LLC  
 Ruby L. Real Trust  
 Semco Land Inc.  
 Standard Farms of Nebraska Inc.  
 Stuckey Grandchildren Partnership  
 Submerged Land Co. Inc.  
 Sylvia L. Schmidt Irrevocable Trust  
 Thelma J. Arneson Trust  
 Thunderbird Farms Inc,  
 Triple E Farms Inc.  
 VCK Farms LLC  
 Whooping Crane Trust  
 William Seng Trust

### **NATIONAL ORGANIZATIONS**

American Bird Conservancy, VA  
 American Rivers, Washington DC  
 Defenders of Wildlife, Washington DC  
 Ducks Unlimited, TN  
 Isaac Walton League, MD  
 National Audubon Society; NY, Washington DC  
 National Trappers Association Inc., WV  
 National Wildlife Federation, VA  
 National Wildlife Refuge Association, Washington DC  
 The Nature Conservancy, CO  
 Sierra Club, CA  
 U.S. Humane Society, Washington DC  
 The Wilderness Society, Washington DC  
 Wildlife Management Institute; CO, OR, Washington DC

### **MEDIA**

Aurora News-Register  
 Bertrand Herald  
 Blue Hill Leader  
 Clay County News  
 Doniphan Herald  
 Franklin County Chronicle  
 Friend Sentinel  
 Gibbon Reporter Office  
 GI Family Radio  
 Grand Island Independent  
 Henderson News  
 Holdrege Daily Citizen  
 Kearney Hub  
 KGFW Radio  
 KHAS-TV  
 KOLN/KGIN-TV  
 KRVN Radio  
 Lincoln Star Journal  
 Milford Times  
 Minden Courier  
 Nebraska Signal  
 NTV Network

Omaha World Herald  
Tribune-Newspaper  
Seward County Independent Newspaper  
Shelton Clipper  
York News Times/Advantage

***UNIVERSITIES AND COLLEGES***

University of Nebraska–Lincoln, Cooperative Unit

***INDIVIDUALS***

352 persons

# Appendix E

## *Environmental Compliance*

### **Environmental Action Statement**

U.S. Fish and Wildlife Service, Region 6  
Lakewood, Colorado

Within the spirit and intent of the Council on Environmental Quality's regulations for implementing the National Environmental Policy Act and other statutes, orders, and policies that protect fish and wildlife resources, I have established the following administrative record.

I have determined that the action of implementing the "Comprehensive Conservation Plan for Rainwater Basin Wetland Management District" is found not to have significant environmental effects, as determined by the attached "Finding of No Significant Impact" and the environmental assessment as found with the draft comprehensive conservation plan.

  
\_\_\_\_\_  
J. Mitch King  
Regional Director, Region 6  
U.S. Fish and Wildlife Service  
Lakewood, CO

8/22/07  
\_\_\_\_\_  
Date

  
\_\_\_\_\_  
Richard A. Coleman, PhD  
Assistant Regional Director, Region 6  
National Wildlife Refuge System  
U.S. Fish and Wildlife Service  
Lakewood, CO

8/21/07  
\_\_\_\_\_  
Date

  
\_\_\_\_\_  
Rod Krey  
Refuge Supervisor (KS, NE, ND, SD)  
U.S. Fish and Wildlife Service  
Lakewood, CO

8/21/07  
\_\_\_\_\_  
Date

  
\_\_\_\_\_  
Gene Mack  
Project Leader  
Rainwater Basin Wetland Management District  
Kearney, NE

8/21/07  
\_\_\_\_\_  
Date

## Finding of No Significant Impact

U.S. Fish and Wildlife Service, Region 6  
Lakewood, Colorado

### *Fulfill the Comprehensive Conservation Plan for Rainwater Basin Wetland Management District*

Two management alternatives for the Rainwater Basin Wetland Management District were assessed as to their effectiveness in achieving the district's purposes and their impact on the human environment.

- Alternative A, the “no-action” alternative, would continue current management.
- Alternative B, “integrated partnership approach” (the proposed action), would focus on addressing resource management in a holistic manner, with an emphasis on cooperation, coordination, and better exchange of information. An expanded district staff would work with partners to improve the waterfowl production areas across the landscape of the Rainwater Basin. The emphasis would be on adaptive management—as more information is known, management would be changed to improve effects on the environment for the benefit of migratory birds in the Central Flyway.

Based on this assessment and comments received, I have selected alternative B as the preferred alternative for implementation.

The preferred alternative was selected because it best meets the purposes for which the Rainwater Basin Wetland Management District was established and is preferable to the “no-action” alternative in light of physical, biological, economic, and social factors. The preferred alternative will continue to provide public access for wildlife-dependent recreation, environmental education, and interpretation.

I find that the preferred alternative is not a major federal action that would significantly affect the quality of the human environment within the meaning of Section 102(2)(C) of the National Environmental Policy Act of 1969. Accordingly, the preparation of an environmental impact statement on the proposed action is not required.

The following is a summary of anticipated environmental effects from implementation of the preferred alternative:

- The preferred alternative will not adversely impact endangered or threatened species or their habitat.
- The preferred alternative will not adversely impact archaeological or historical resources.
- The preferred alternative will not adversely impact wetlands nor does the plan call for structures that could be damaged by or that would significantly influence the movement of floodwater.
- The preferred alternative will not have a disproportionately high or adverse human health or environmental effect on minority or low-income populations.
- The state of Nebraska has been instrumental in the development of this plan and has been notified and given the opportunity to review the comprehensive conservation plan and associated environmental assessment.

  
J. Mitch King  
Regional Director, Region 6  
U.S. Fish and Wildlife Service  
Lakewood, CO

  
Date

# Appendix F

## *Black-tailed Prairie Dog Management Plan*

**BLACK TAILED PRAIRIE DOG MANAGEMENT PLAN  
Rainwater Basin Wetland Management District**

**Written and Revised in 2003 by  
Tom Koerner, Deputy Refuge Manager  
&  
Jeff Drahota, Wildlife Biologist**

*Approvals and Concurrence*

Rainwater Basin Wetland Management District:

***Project Leader***

Gene D. Mack

Signature

Date

Mountain-Prairie Regional Office:

***National Wildlife Refuge System***

***Refuge Supervisor, Nebraska, Kansas & Colorado:***

Ron Shupe

Signature

Date

***Regional Wildlife Biologist:***

Wayne King

Signature

Date

### Description

The scientific name for the black-tailed prairie dog is *Cynomys ludovicianus*. "Ludovicianus," is the Latin form of Ludwig or Louis, relating back to the Lewis and Clark expedition of 1804-1806, when prairie dogs were first collected for science.

The prairie dog is a burrowing member of the order Rodentia, the largest group of mammals in the world. An adult black-tailed prairie dog is between 12 and 16 inches long and generally weighs between 1.5 and 2.5 pounds. Its tail is covered with hair and is about one-fourth of the animal's total length. Its body is tan to pale brown in color, its underparts are white to buffy white, and its tail is tipped with black. The prairie dog's legs are short, but its feet are large and have well-developed claws, especially on the forefeet. Its head is broad and rounded, and its eyes are fairly large.

### Distribution and Abundance

The black-tailed prairie dog is one of five species of prairie dogs found in North America. It is the most abundant and widely distributed species and is the only prairie dog found in Nebraska. It is found throughout the Great Plains from southern Canada to just inside Mexico. The western edge of its range is along the Rocky Mountains, and the eastern edge follows the natural boundary between tall and mid-grass prairie. In Nebraska, prairie dogs are found roughly in the western two-thirds of the state. Historic towns have been documented on a number of WPAs throughout the WMD; however, the most active towns are found in the west half of the WMD.

Black-tailed prairie dogs live in colonies or "towns" that range in size from as small as one acre to several thousand acres. The largest prairie dog colony on record was in Texas, and was about 100 miles wide, 250 miles long and contained an estimated 400 million animals. It is estimated that in the late 1800s, some 700 million acres of North American rangeland were inhabited by prairie dogs. Habitat changes and extensive eradication efforts have reduced the acreage by about 90 to 95 percent from historic levels.

We believe that on WPAs, the cessation of regular grazing on these towns led to the eventual loss of the town due to the creation of unsuitable habitat. This factor becomes more important the farther east one travels. Increasing precipitation levels, resulting in higher vegetative growth without grazing may result in less suitable habitat.

### Habitat and Home

Areas of short and mid-grass rangeland that are heavily grazed by livestock are the prairie dog's preferred habitat. Prairie dog colonies are most recognizable by the mounds and holes at their burrow entrances. A colony will typically have 30 to 50 burrow entrances per acre.

The animal's burrow system can be quite complex and extensive. Mounds of excavated soil around the burrow entrance are generally cone-shaped and vary from one to three feet in height and from three to 10 feet in diameter. These mounds serve as lookout points and serve to prevent water from entering the burrow system. Tunnels are generally three to six feet below the surface and about 15

## Introduction

feet long, although burrows have been reported to reach depths of 15 feet. Burrow systems typically include several chambers, including one near the entrance where the prairie dog can sit and listen for activity above ground, and one or more nest chambers where they sleep and care for their young.

## Habits

The fact that prairie dogs live in colonies indicates they are highly social animals. The largest social unit is the colony or town. Towns are often divided into "wards" by topographical barriers such as roads, ridges or trees, and are generally five to 10 acres in size. Although prairie dogs in one ward may be able to see and hear animals of an adjacent ward, movement among wards is unusual. Wards are divided into several smaller prairie dog social units, called "coterie." Each coterie generally consists of one adult male, one to four adult females, and any offspring less than two years old. Members of one coterie defend their territory from invasion by members of other coterie.

Prairie dogs are active during the day, usually from about sunrise to sunset, and during summer they spend about one-third to one-half of the daylight hours feeding. Another third is involved in social interactions with other colony members as well as working on burrows and mounds and responding to alarm calls. The remainder of daylight is spent underground, especially during midday when temperatures above ground are high. The black-tailed prairie dog is active all year. In winter, it remains underground for several days when weather is severe, but comes out on sunny afternoons to look for food and bask in the sun.

Black-tailed prairie dogs exhibit an elaborate communication system. At least 11 separate calls have been identified, and a variety of postures and displays are utilized. Calls range from signals of alarm to "all-clear." Physical contact is another method of prairie dog communication. Mouth-to-mouth contact is used to identify coterie members from strangers, and grooming among coterie members is common.

## Food

Grasses are the preferred food of the prairie dog, and generally makes up about three fourths of its diet. In the fall, broadleaf forbs become more important as green grass is less available. In winter, any available green vegetation is consumed. In the spring and summer, each prairie dog consumes up to two pounds of vegetation per week.

In addition to the vegetation it eats, the prairie dog also clips much of the vegetation within its colony. This is probably done to keep the vegetation clipped short to provide an unobstructed view of approaching predators. Over a period of time, clipping, foraging and digging activities can alter the composition of the vegetation in a prairie dog town. Short, native grass like buffalograss and blue grama is favored when present.

## Reproduction

## Site Description

A prairie dog reaches sexual maturity after its first winter and has one litter per year. Breeding takes place in March and early April, and a litter of 4-6 young is born 30 to 35 days later. Young prairie dogs are born hairless, helpless, and with their eyes closed. They remain underground for about six weeks and first emerge from the den in May or June. They are weaned at this time and begin feeding on green vegetation. They reach adult size by fall.

### Mortality

Although the prairie dog has been known to live for at least eight years in captivity, its average life span in the wild is usually three to four years. In addition to actions of man, the prairie dog faces many natural predators. Badgers are probably the main predators, but coyotes, weasels, golden eagles, hawks, swift fox, and other predators take prairie dogs. Bullsnares and rattlesnakes take young prairie dogs but generally not adults. The black-footed ferret was once a primary prairie dog predator, but it is now considered an endangered species and no wild ferrets have been verified in Nebraska since the 1940s.

A prairie dog is susceptible to a number of diseases, the most notable being plague. Plague is an infectious disease transmitted by the bite of an infected flea. Plague can devastate prairie dog populations, wiping out entire colonies in some areas. This disease was known as "black death" in the 1300s when about one-third of Europe's human population was lost. Although it can be transmitted to humans through the bite of an infected flea, plague has not been found in prairie dogs in Nebraska and is now treatable in humans.

### Importance

In many ways, a prairie dog town can be considered a biological oasis. Many wildlife species associate with prairie dogs. Some species feed on prairie dogs, but others utilize the burrow systems or the unique habitat to fulfill their needs. Vacant burrows are used by cottontail rabbits, several species of small rodents, tiger salamanders, and by burrowing owls. Our most active towns have had successful nesting by burrowing owls. Meadowlarks, grasshopper sparrows, and other birds are found in greater numbers in prairie dog towns than in the surrounding rangeland because they are attracted to the open spaces, where seeds and insects are more accessible. In addition to their importance to other wildlife species, prairie dogs are also important to wildlife observers and photographers.

## **SITE DESCRIPTION**

Originally, the Rainwater Basin in south central Nebraska contained more than 3,900 wetland basins within a 17-county area. The Basin region covers 4,200 square miles of flat to gently rolling Peorial Loess Plains. Wetland basins are generally large, shallow depressions with deep clay layers in the wetland basin—creating an impervious water barrier. The name Rainwater Basin

## Site Description

comes from the basins' ability to go from dry to flooded conditions quickly—following heavy rainstorms and snow melts. Continual siltation problems result from rapid runoff carrying heavy silt loads from agricultural land resulting in poor water quality. Soils surrounding the basins are very fertile, consisting of heavy silt loams and silty clay. Deep beneath the soil lies the Ogallala Aquifer, which is the source of water for the extensive amount of irrigated corn and soybeans. Water from the Platte River, lying north of the region, is delivered by irrigation canals to irrigate the southwest portion of the region. Agricultural and rural development has destroyed more than 90 percent of the original number of wetlands.

The Fish and Wildlife Service began acquiring wetlands in 1963 with the purchase of Massie Waterfowl Production Area (WPA). By 1966, 7,000 acres were acquired and a management office was established in Hastings, Nebraska. In 1976, the office was moved to its present location: Kearney, Nebraska.

The Rainwater Basin Wetland Management District (WMD) currently manages 63 tracts of land, 61 of which are Waterfowl Production Areas (WPA) totaling 23,059 acres. One of the remaining two areas is McMurtrey Wildlife Management Area that was transferred from the military. Its 1052 acres are closed to public use. The other tract is the Platte River National Wildlife Management Area (438 acres). This property is owned by the state of Wyoming and managed through a memorandum of understanding. WPAs are small isolated tracts of land scattered throughout the District. Most WPAs contain only one large wetland. All WPAs are managed as a grassland/wetland ecosystem. Wetland management is focused toward providing optimum waterfowl and shorebird habitats. The uplands are managed for a high diversity of native tall and mid-grass plant species. Thirty-eight FmHA conservation easements totaling 2350 acres are managed by this office as well.

Spring migration is the primary focus of the Rainwater Basin WMD. Each spring, about six million snow geese, one million Canada geese, 90 percent of the mid-continent white-fronted goose population, millions of ducks, and one-half million sandhill cranes use these wetlands and associated Platte River areas. Habitat becomes very critical during this time of year. Extensive pumping and aggressive wetland management are needed to maintain quality habitat for resting and staging.

In addition, we manage habitats to provide habitat for many other species of migratory birds and resident wildlife which utilize these grasslands and adjacent uplands.

### **Soils**

Soils throughout the Rainwater Basin vary. In general, they are characterized as Peorial Loess. The soils formed under tall and mid-grass prairies and are characterized as deep, well drained, and fertile. The wetland or hydric soils are scattered throughout the WMD and are generally believed to have formed through a combination of wind and water erosion. Scouring events created a depression and rainfall events caused the migration of clays to the bottom of the basin. These hydric soils (Massie, Scott, and Fillmore soils) have a clay layer from 1 to 10 feet thick. They create an impermeable layer and restrict movement of water.

### **Surrounding Land Uses**

The major industry and source of income throughout the WMD is crop and livestock production. Nearly every acre has been converted to agricultural use. Small to medium sized communities (25,000 residents) are scattered across the WMD.

### **Current Status of Black-tailed Prairie Dogs on WPAs**

We currently have 5 WPAs with known active black-tailed prairie dog towns (see Figure 1). These are Prairie Dog (see Figure 2), Atlanta (see Figure 2), McMurtrey (see Figure 4), Hultine (see Figure 5), and Clark (see Figure 6) WPAs. These were surveyed in 1999 and again in 2003. McMurtrey, Atlanta, and Prairie Dog WPAs appear to have viable populations at this time. Atlanta WPA currently has the largest dog town at 24 acres yet, most are less than 8 acres.

On Prairie Dog WPA, a number of abandoned burrows were re occupied after initiating grazing on the sites. Likely, individuals from the active burrows relocated into these abandoned burrows.

The town on Atlanta WPA has remained relatively the same size, and has not expanded outside of the electric fence lines placed to control grazing access. The population on McMurtrey NWR, appears to be well established from re-introductions made in 2001.

The towns on Clark and Hultine WPAs established after several years of heavy grazing. The newly established towns on both sites are small, containing less than 12 burrows each in the summer of 2003. They both have the potential to expand into previously occupied burrows.

## **REINTRODUCTION**

## Reintroduction

A number of additional WPAs have evidence that historic black-tailed prairie dog towns (see Figure 7). Old burrows and mounds can still be found. We propose that re-introductions of black-tailed prairie dogs be made into these historic sites. In addition, several high public use sites on Harvard, Massie, Mallard Haven, and Funk WPAs that have the potential for environmental education and interpretation along with establishing a viable population, are proposed release sites. Although no evidence in the form of mounds or burrows exists at these sites, it is believed that historically black-tailed prairie dogs could be found in the immediate vicinity of these sites. Currently, seven release sights including both historic and new dog towns are proposed (see Figure 8) and will be populated as circumstance allow. We propose re-introductions of black-tailed prairie dogs into these sites under the following conditions:

1. All landowners immediately adjacent to the proposed reintroduction site have been contacted to discuss our intent, management options, and concerns.
2. Private rangeland is not located immediately adjacent to the proposed release site and/or an adequate cultivated barrier exists to minimize movement from the re-introduction site and discourage colonization on adjacent private land.
3. Suitable habitat exists at the time of release. Suitable habitat includes poor stands of grass commonly dominated by introduced grasses.

Intended release sites should provide suitable black-tailed prairie dog habitat at the time of release. This would include vegetation 3" or less in height and a minimum of one "burrow" for every four animals released. The vegetation could be grazed, or mowed if grazing were not possible, through the growing season to keep it short. Burrows could be present from a historic town, or they could be drilled in with the 6" auger attachment on the Bobcat. Drill one burrow for every two dogs released. Both of these conditions will reduce the likelihood that the prairie dogs will abandon the site upon release and would likely result in a higher survival rate of released animals.

### Pre-Capture

We are occasionally contacted by individuals wishing to remove prairie dogs from their property. They would be willing to allow these prairie dogs to be live trapped for relocation. If we have a suitable re-introduction site where all three conditions stated above have been met, we may consider accepting these prairie dogs for reintroduction purposes. Each proposal will be evaluated on a case by case basis, giving preference to those sites closest to the proposed reintroduction site. If our staff will be required to conduct the trapping and relocation, time budgets of the staff will be given priority and will only be conducted when our workload allows.

Prior to trapping, the prairie dog town that will be trapped should be surveyed for indications of

## Reintroduction

disease. A walk through should be done looking for signs of dead animals including other species. Remote monitoring of animals should be done using a spotting scope for one hour during midday (at least three hours after sunrise and two hours before sunset) on a sunny day. It is recommended that all burrows be treated 10 days prior to trapping with Delta Dust to kill fleas currently inhabiting the colony. If dead animals are found, two options can be taken: 1) bag the animal and send to the disease lab to determine cause of death; 2) find another trapping sight.

Pre-baiting with should start three days before trapping begins. Live traps can be placed within 3 feet of active burrows and locked open during pre-baiting. Trapping should start after June 1 and can be done until sub-freezing temperatures occur.

Monkey pox was recently a high profile news story with regards to handling of prairie dogs. The origin of the outbreak was traced to a shipment of prairie dogs in the pet trade. These prairie dogs then infected humans which handled them. USDA APHIS immediately placed restrictions on the trade and handling of prairie dogs. At this time, it does not appear that these restrictions apply to the US Fish and Wildlife Service with regards to ongoing research and/or management of wild populations of prairie dogs. This new outbreak should reinforce the need to research the background of any potential capture sites and exclude any that may have had captive black-tailed prairie dogs released on site.

## Capture

New bait but smaller portions should be used during the trapping stage. Traps should be unlocked with bait placed on the pan or just behind the pan. If needed, flagging should be used to mark traps since they may be difficult to monitor from a distance so not to disturb the colony unless animals have been captured. Traps should not be left unlocked if they cannot be checked at least every 24 hours.

Once animals have been captured, they should be dusted with flea powder (the product Seven is recommended) before the trap is moved. Long gloves should be used to prevent fleas from biting around the wrist. Traps can then be placed in the back of a pickup and removed from the site and taken to the quarantine site.

A 24-hour quarantine is required after trapping. Captured animals should be left in the traps they were captured in. They should be placed in a cool and dry area such as a pole barn or well ventilated shed. If a longer quarantine is used, apple slices can be used to sustain the animals and will meet their food and water requirements.

Any dead animals found during the quarantine should be handled with extreme caution. These animals should be sent to the disease lab for analysis. If plague is found, these animals should not

## Management

be used for reintroduction.

### Post Release

After release, the site will be periodically surveyed to determine presence or absence of prairie dogs, if vegetation is conducive to expansion, limiting expansion, or maintaining populations. Currently, 8-10 acre prairie dog towns appear to be manageable. Larger towns could be maintained on a few WPA's as long as the station's mission and goals are not jeopardized. Three or four areas could have up to 30 acre prairie dog towns if further expansion is controllable and the threat of colonization on adjacent land is minimal. It is also necessary to monitor adjacent cropland for potential foraging impacts due to overcrowding, expansion, or changes in forage quantity on WPA's.

### MANAGEMENT

Management includes any activity conducted to control the size of the prairie dog town, maintain the habitat suitability for black-tailed prairie dogs, and/or ensure its long term viability.

Hunting black-tailed prairie dogs on Federal Waterfowl Production Areas in the Rainwater Basin WMD was closed as of 01/01/2003. We propose to leave this closure in place indefinitely. This decision was made based upon the relatively small size of the towns, small populations, and relative isolation from other active towns. Although significant harvest was not believed to be occurring before the closure, changes in local interest in prairie dog shooting along with decreasing opportunities would likely have led to an increase in harvest.

Active prairie dog towns on Atlanta, Prairie Dog, McMurtrey, Clark, and Hultine WPAs have been managed primarily through heavy grazing over the last four seasons. Grazing cooperators are instructed to place water tanks on the town site. This results in short, clipped vegetation which is more suited to maintaining a viable town. We have interseeded buffalo grass and blue grama on these sites, with the expectation that increased grazing pressure will result in establishment of these two, low growing grasses. Although not widely distributed in the mid and tall grass prairie, buffalo grass and blue grama can be found in areas which are heavily grazed. We plan to continue grazing at a rate sufficient to maintain a vegetative height less than 3" on active towns. If grazing is not an option in some years, mowing may be considered as a possible alternative.

## Management

Historic black-tailed prairie dog towns may be re-occupied without re-introductions simply by providing suitable habitat. Prairie dogs, especially the first year class, have been documented to disperse into suitable habitat located several miles away. By grazing these historic towns at a rate sufficient to provide suitable habitat, it is conceivable they may be re-occupied at some point. This is what we believe occurred on Clark and Hultine WPAs.

Our land acquisition program has the potential to acquire new WPAs that may contain active black-tailed prairie dog towns. Should we acquire lands with an active black-tailed prairie dog town, we will manage the sites in a similar fashion to maintain a viable town within a defined boundary.

It has been demonstrated that fence placement alone has been sufficient to prevent the spread of a town outside of the defined boundary on Prairie Dog and Atlanta WPAs (see Figure 2 & 3). We feel that this phenomenon will hold true across the WMD. The grassland vegetation in these precipitation zones grows faster than the prairie dogs can cut it, resulting in unsuitable habitat. If private pasture exists adjacent to the release site, it is conceivable to believe that some prairie dogs would relocate to this pasture and cause damage by their burrowing activities. That is why release under these circumstances will be carefully considered. The majority, however, have adjacent cropland. The annual tillage and tall growth eliminate cropland as potential black-tailed prairie dog habitat.

### Control

In the event that our assumptions are wrong and that the black-tailed prairie dogs spread outside of the defined boundaries, control measures will be taken. All legal and effective methods for prairie dog control may be considered, excluding the use of any toxicants.

Some options considered but are not limited too:

- |               |   |
|---------------|---|
| Box traps     | Although relatively ineffective at capturing large numbers, this may be considered where individual animals may want to be salvaged for some purpose. |
| Leghold traps | Placed in front of burrow entrances, may be relatively effective at reducing a small number of animals.   |

## Management

- Visual barriers The use of snow fence, hay bales, or other visual obstructions have proven successful in some places with reducing or eliminating the spread of prairie dogs from a defined boundary.
- Raptors The installation of raptor perches along the perimeter of prairie dog towns has proven successful at some locations for reducing the spread of the town.
- Shooting Shooting individual animals which have spread outside of the designated boundaries could prove effective. This could be accomplished by US Fish and Wildlife Service staff or by an individual through a Special Use Permit.

**REFERENCES**

Black-tailed Prairie Dog Surveys for 1999 and 2003. Rainwater Basin Wetland Management District. Unpublished report.

Boddicker, Major L. Prevention and Control of Wildlife Damage. B-75 to B-84 Prairie Dogs.

Hynstrom, Scott E., and Dallas R. Virchow. Prairie Dogs and the Prairie Ecosystem.

Luce, Bob, et. al. A Multi-State Conservation Plan for the Black-tailed Prairie Dog, *Cynomys ludovicianus*, in the United States. 2002.

Nebraska Game and Parks Commission Website:  
<http://www.ngpc.state.ne.us/wildlife/pdogs.html>

Van Pelt, William E. The Black-tailed Prairie Dog Conservation Assessment and Strategy.

## References

# 2003 Active Prairie Dog Towns on WPAs



-  All roads
-  WPAs with Prairie Dog towns
-  Rainwater Basin Counties
-  Other WPAs





Figure 1



Figures

# Proposed Black-tailed Prairie Dog Release Areas



-  All roads
-  Proposed WPAs for future Prairie Dog releases
-  Rainwater Basin Counties
-  Other WPAs




Figure 8

# Appendix G

## Section 7 Biological Evaluation

### INTRA-SERVICE SECTION 7 BIOLOGICAL EVALUATION FORM For

Issuance and Implementation of the Comprehensive Conservation Plan for the  
U.S. Fish and Wildlife Service  
Rainwater Basin Wetland Management District



Originating Person: Gene Mack  
Jeff Drahota  
Bernardo Garza  
Telephone Number: (308) 236-5015  
Date: August 31, 2007

- I. Region: 6**
- II. Service Activity (Program):** Rainwater Basin Wetland Management District (WMD), Comprehensive Conservation Plan (CCP).
- III. Pertinent Species and Habitat:**
- A. Listed species and/or their critical habitat within the action area:**  
whooping crane (E), piping plover (T), Interior least tern (E)
- B. Proposed species and/or proposed critical habitat within the action area:**  
none
- C. Candidate species within the action area:**  
none
- IV. Geographic area or station name and action:**  
Rainwater Basin National Wildlife Refuge Complex  
Issuance and Implementation of the CCP for Rainwater Basin WMD
- V. Location: (map attached)**
- A. Ecoregion number and name:** Platte/Kansas Ecoregion
- B. County and State:** Buffalo County, Nebraska. The Rainwater Basin WMD encompasses 17 counties in south-central Nebraska.
- C. Distance and direction to nearest town:** The Rainwater Basin Complex headquarters is located in Kearney, Nebraska.
- D. Species/habitat occurrence:**
- Whooping crane: The Rainwater Basin WMD is a key stopover site for this species migrating through the Central Flyway to their breeding area in the Northwest Territories in the spring and their wintering area on Aransas NWR in the fall. Since

2000 there have been at least eight confirmed fall observations of whooping cranes using Rainwater Basin WMD. Additionally, during recent years, this species has been documented on the Waterfowl Production Areas (WPA) of the complex.

**Piping plover:** The piping plover breeds and forages along river channels and nest on sparsely vegetated sandbars. It also breeds on large, alkaline lakes. The piping plover may be an uncommon visitor during the migration period, but has not been documented to nest within the Rainwater Basin WMD.

**Interior Least tern:** The Interior least tern breeds and forages along river channels and nest on sparsely vegetated sandbars. It has been documented on Rainwater Basin WMD, but this is an anomaly, as the majority of this species' habitat use in Nebraska centers on the major river systems such as Missouri River, Platte River, Niobrara River, and Loup River.

#### **VI. Description of proposed action:**

Implementation of Alternative 2 of the CCP to guide the management of the Rainwater Basin WMD for the next 15 years. Implementation of this CCP comprises implementation of all actions and activities to achieve the stated goals contained in the CCP that will ultimately lead to the fulfillment of the purposes for which Congress established all the units comprising the Rainwater Basin WMD and assist in the fulfillment of the goals of the National Wildlife Refuge System.

#### **VII. Determination of effects:**

##### **A. Explanation of effects of the action on species and critical habitats in Items III. A, B and C:**

**Whooping crane:** Implementing the CCP is not thought to have detrimental effects on this migrant crane. In fact, the continued preservation and management of WPAs for the benefit of wildlife species should enhance loafing, feeding and resting sites for whooping cranes.

**Interior least Tern/**

**Piping plover:** Implementing the CCP is not thought to have detrimental effects on either the Interior least tern or piping plover. In fact, the continued preservation and management of WPAs, especially grazing management may increase this species use of WPAs.


**B. Explanation of actions to be implemented to reduce adverse effects:**

None are necessary. All actions delineated in this CCP are thought to follow and be in accordance with provisions of protection and restoration plans for several species, as delineated by the Service and other federal and State agencies. The WMD staff is well acquainted with provisions that would be invoked and be put into effect to protect federally listed species and species of special concern from any public use or management action by WMD staff or visitors to the WMD.

**VIII. Effect determination and response requested:**

**A. Listed species/designated critical habitat:**

<u>Determination</u>	<u>Response Requested</u>
may affect, not likely to adversely affect species/adversely modify critical habitat	
Whooping Crane	<input checked="" type="checkbox"/> Concurrence
Least Tern	<input checked="" type="checkbox"/> Concurrence
Piping Plover	<input checked="" type="checkbox"/> Concurrence

  
 \_\_\_\_\_  
 Gene Mack, Project Officer

8-31-07  
 \_\_\_\_\_  
 Date

**IX. Reviewing ESO Evaluation:**

- A. Concurrence  Nonconcurrency \_\_\_\_\_
- B. Formal Consultation Required \_\_\_\_\_
- C. Conference Required \_\_\_\_\_
- D. Informal Conference Required \_\_\_\_\_

  
 \_\_\_\_\_  
 Assistant Nebraska Field Supervisor

9/4/07  
 \_\_\_\_\_  
 Date

Ecological Services  
Grand Island, Nebraska



# Appendix H

## List of Plants

The Service maintains a list of plant species at <http://www.fws.gov/nwi/bha/list88.html> that occur in wetlands. The following list of plants for WPAs managed by the Rainwater Basin Wetland Management District was generated using the region 5 map (1,523 plant species) from this website. The USDA Plants website at <http://plants.usda.gov> listed 9,485 plant species for Nebraska. Both the USDA Plants website and the NatureServe Explorer website at <http://www.natureserve.org/explorer> were used to verify scientific names, common names, and habits for those species not listed on the Service website.

### SCIENTIFIC NAME

Scientific name is the genus and species applied to the taxon by the National List of Scientific Plant Names (1982), USDA Plants, NatureServe Explorer (2006).

### NATIONAL RANGE OF INDICATORS

The national indicators reflect the range of estimated probabilities (expressed as a frequency of occurrence) of a species occurring in wetland versus nonwetland across the entire distribution of the species. A frequency, for example, of 67%–99% (facultative wetland) means that 67%–99% of sample plots containing the species randomly selected across the range of the species would be wetland. A question mark (?) following an indicator denotes a tentative assignment based on the botanical literature and not confirmed by regional review. When two indicators are given, they reflect the range from the lowest to the highest frequency of occurrence in wetlands across the regions in which the species is found. A positive (+) or negative (–) sign was used with the facultative indicator categories to more specifically define the regional frequency of occurrence in wetlands. The positive sign indicates a frequency toward the higher end of the category (more frequently found in wetlands), and a negative sign indicates a frequency toward the lower end of the category (less frequently found in wetlands).

### NATIONAL WETLAND INDICATOR (INDICATOR CATEGORIES)

**Obligate Wetland (OBL).** Occur almost always (estimated probability >99%) under natural conditions in wetlands.

**Facultative Wetland (FACW).** Usually occur in wetlands (estimated probability 67%–99%), but occasionally found in nonwetlands.

**Facultative (FAC).** Equally likely to occur in wetlands or nonwetlands (estimated probability 34%–66%).

**Facultative Upland (FACU).** Usually occur in nonwetlands (estimated probability 67%–99%), but occasionally found on wetlands (estimated probability 1%–33%).

**Obligate Upland (UPL).** Occur in wetlands in another region, but occur almost always (estimated probability >99%) under natural conditions in nonwetlands on the region specified. If a species does not occur in wetlands in any region, it is not on the National List.

The wetland indicator categories should not be equated to degrees of wetness. Many obligate wetland species occur in permanently or semipermanently flooded wetlands, but a number of obligates also occur and some are restricted to wetlands that are only temporarily or seasonally flooded. The facultative upland species include a diverse collection of plants that range from weedy species adapted to exist in a number of environmentally stressful or disturbed sites (including wetlands) to species in which a portion of the gene pool (an ecotype) always occurs in wetlands. Both the weedy and ecotype representatives of the facultative upland category occur in seasonally and semipermanently flooded wetlands.

### REGION 5 WETLAND INDICATOR (REGIONAL INDICATOR FOR 5—CENTRAL PLAINS NE, KS, CO [EASTERN])

The regional indicator reflects the estimated probability (likelihood) of a species occurring in wetlands versus nonwetlands in the region. Regional indicators reflect the unanimous agreement of the Regional Interagency Review Panel. If a regional panel was not able to reach a unanimous decision on a species, NA (no agreement) was recorded on the regional indicator field. An NI (no indicator) was recorded for those species for which insufficient information was available to determine an indicator status. A nonoccurrence (NO) designation indicates that the species does not occur in that region. An asterisk (\*) following a regional indicator identifies tentative assignments based on limited information from which to determine the indicator status.

**HABIT**

Habit indicates the plant characteristics and life forms assigned to each species in the National List of Scientific Plant Names (1982) and by the Soil Conservation Service. Family names are listed alphabetically under specific life forms restricted to these families. The habit symbols are combined to describe the life form of the species (for example, ANG means annual native grass and IT means introduced tree).

<i>Habit Symbol</i>	<i>Characteristic or Life Form</i>	
A	Annual	
B	Biennial	
C	Clubmoss	Lycopodiaceae Selaginellaceae
E	Emergent	
F	Forb	
/	Floating	
F3	Fern	Adiantaceae Aspleniaceae Blechnaceae Cyatheaceae Davalliaceae Dennstaedtiaceae Dryopteridaceae Gleicheniaceae Grammitidaceae Hymenophyllaceae Lomariopsidaceae Marattiaceae Ophioglossaceae Osmundaceae Parkeriaceae Polypodiaceae Psilotaceae Pteridaceae Schizaeaceae
G	Grass	Poaceae
GL	Grasslike	Cyperaceae Juncaceae
H	Partly woody	
HS	Half shrub	
H2	Horsetail	Equisetaceae
I	Introduced	
N	Native	
P	Perennial	
+	Parasitic	
P3	Pepperwort	Marsileaceae
Q	Quillwort	Isoetaceae
S	Shrub	
-	Saprophytic	
Z	Submerged	
\$	Succulent	
T	Tree	
V	Herbaceous vine	
W	Waterfern	Azollaceae Salviniaceae
WV	Woody vine	

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Abutilon theophrasti</i>	velvetleaf	UPL–FACU–	UPL	AIF
<i>Achillea millefolium</i>	common yarrow	FACU	FACU	PNF
<i>Agropyron elongatum</i>	tall wheatgrass	FAC,FACU*	NI	PNG*
<i>Agropyron smithii</i>	western wheatgrass	FACU	UPL,FAC–	PNG
<i>Agrostis hyemalis</i> ( <i>A. antecedens</i> )	ticklegrass	FACU	FACU,FACW	PNG
<i>Agrostis gigantea</i>	redtop	FAC+,FACW	FAC+	PNG
<i>Alisma plantago-aquatica</i>	European water plantain	OBL	OBL	PNEF
<i>Alisma subcordatum</i>	American water plantain	OBL	OBL	PNEF
<i>Allium canadense</i>	meadow onion	FACU–,FACU	FACU	PNF
<i>Allium canadense</i> var. <i>lavendulare</i>	meadow onion	FACU–,FACU	FACU	PNF
<i>Alopecurus carolinianus</i>	Carolina foxtail	FACW	FAC+,FACW	ANG
<i>Amaranthus arenicola</i>	sandhill amaranth	UPL,FAC	FACU	ANF
<i>Amaranthus retroflexus</i>	redroot amaranth	FACU	FACU–,FAC–	ANF
<i>Amaranthus rudis</i>	amaranth	FACW	FACU–, FACW	ANF
<i>Ambrosia artemisiifolia</i>	annual ragweed	FACU	FACU–,FACU+	ANF
<i>Ambrosia grayi</i>	woollyleaf burr ragweed	FAC	FAC,FACW	PNF
<i>Ambrosia psilostachya</i>	naked-spike ragweed	FAC	FACU–, FAC	PNF
<i>Ambrosia trifida</i>	great ragweed	FACW	FAC,FACW	ANF
<i>Ammannia coccinea</i>	valley redstem	OBL	FACW+,OBL	ANF
<i>Amorpha canescens</i>	leadplant	FAC*	NI	PNF*
<i>Andropogon gerardii</i>	big bluestem	FAC–	FACU,FAC	PNG
<i>Apocynum cannabinum</i>	clasping-leaf dogbane	FAC	FACU,FAC+	PNF
<i>Argemone</i> spp. ( <i>polyanthemos</i> )	crested pricklypoppy	UPL		PNF
<i>Aristida oligantha</i>	three-awn grass	FACU*	NI	PNG*
<i>Artemisia ludoviciana</i>	white sagebrush	FACU–	UPL,FACU	PNFH
<i>Asclepias stenophylla</i>	slim-leaved milkweed	FACU*	NI	PNF*
<i>Asclepias syriaca</i>	common milkweed	FACW–*	NI	PNF
<i>Asclepias verticillata</i>	whorled milkweed	UPL		PNF
<i>Asclepias viridiflora</i>	woolly milkweed	UPL		PNF
<i>Asclepias viridis</i>	green antelopehorn	UPL		PNF
<i>Aster junciformis</i>	rush aster	OBL	OBL	PNF
<i>Aster</i> spp. (see <i>Symphotrichum</i> )	wild aster	UPL,OBL*	NI	PINF*
<i>Astragalus canadensis</i>	Canadian milkvetch	FACU	FACU,FACW–	PNF
<i>Astragalus crassicaarpus</i>	groundplum milkvetch	UPL		PNF

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Azolla mexicana</i>	Mexican mosquito fern	OBL	OBL	PN/W
<i>Bacopa rotundifolia</i>	disk water-hyssop	OBL	OBL	PNF
<i>Baptisia bracteata</i>	long-bract indigo	UPL		PNF
<i>Berula erecta</i>	water parsnip	OBL	OBL	PIF
<i>Bidens cernua</i>	nodding beggar-ticks	OBL	FACW+,OBL	AIF
<i>Bidens comosa</i>	leafy-bract beggar-ticks	FACW	FACW	ANF
<i>Bidens frondosa</i>	devil's beggar-ticks	FACW	FACW,FACW+	ANF
<i>Bidens vulgata</i>	big devil's beggar-ticks			ANF
<i>Boltonia asteroides</i>	white boltonia	FACW	FACW,OBL	PNF
<i>Bouteloua curtipendula</i>	sideoats grama	FACU*	NI	PNG*
<i>Bouteloua gracilis</i>	blue grama	UPL*	NI	PNG*
<i>Brickellia eupatorioides</i>	false boneset	UPL		PNF*
<i>Bromus inermis</i>	smooth brome	UPL*	NI	PNG*
<i>Bromus japonicus</i>	Japanese brome	FACU	FACU,UPL	AIG
<i>Bromus tectorum</i>	downy brome	UPL*	NI	ANG*
<i>Bryophyte spp.</i>	moss	NI	NI	NI
<i>Buchloe dactyloides</i>	buffalograss	FACU	FACU-,FACU	PNG
<i>Calamagrostis canadensis</i>	bluejoint	OBL	FAC,OBL	PNG
<i>Calamovilfa longifolia</i>	prairie sandreed	UPL		PNG
<i>Callirhoe alcaeoides</i>	light poppymallow	UPL		PNF
<i>Callirhoe involucrata</i>	purple poppymallow	UPL		PNF
<i>Calylophus serrulatus</i>	yellow sundrops	UPL		PNF
<i>Calystegia sepium</i> ( <i>Convolvulus sepium</i> )	hedge false bindweed	FAC		PNFV
<i>Cannabis sativa</i>	hemp	FAC+*	NI	ANF*
<i>Capsella bursa-pastoris</i>	common shepherd's purse	FACU	FACU,FAC	AIF
<i>Cardus nutans</i>	musk thistle	UPL		B/PIF*
<i>Carex bicknellii</i>	Bicknell's sedge	FACU,FACW	FACU	PNGL
<i>Carex brevior</i>	short-beak sedge	FAC	UPL,OBL	PNEGL
<i>Carex cristatella</i>	crested sedge	FACW	FAC,FACW+	PNGL
<i>Carex gravida</i>	heavy sedge	OBL*		PNGL
<i>Carex laeviconica</i>	smooth-cone sedge	OBL	OBL	PNEGL
<i>Carex lanuginosa</i> ( <i>C. pellita</i> )	woolly sedge	OBL	OBL	PNGL
<i>Carex tribuloides</i>	blunt broom sedge	FACW,OBL	FACW	PNGL
<i>Carex vulpinoidea</i>	fox sedge	OBL	OBL	PNEGL

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Carex × stipata</i>	stalk-grain sedge	OBL	OBL	PNGL
<i>Cassia chamaecrista</i>	showy partridge pea	UPL		ANF
<i>Ceanothus americanus</i>	New Jersey tea	UPL		PNF
<i>Ceanothus herbaceus</i>	Jersey tea	UPL		PNS
<i>Celtis occidentalis</i>	hackberry			PNT
<i>Celtis occidentalis</i>	common hackberry	UPL		PNT
<i>Cenchrus longispinus</i>	sandbur	FAC*	NI	ANG*
<i>Chenopodium album</i>	lambsquarters	FAC	FACU,FAC	AIF
<i>Chenopodium desiccatum</i>	aridland goosefoot	UPL		ANF
<i>Chenopodium leptophyllum</i>	narrowleaf goosefoot	NI	UPL,FAC	ANF
<i>Cichorium intybus</i>	chicory			BPIF
<i>Circaea</i> spp.	enchanter's nightshade	FACW,UPL		PNF
<i>Cirsium altissimum</i>	roadside thistle	FAC*	NI	BNF*
<i>Cirsium arvense</i>	Canada thistle	FACU	FACU-,FAC	PIF
<i>Cirsium canescens</i>	prairie thistle	UPL		BNF*
<i>Cirsium flodmanii</i>	Flodman's thistle	NI	FACU?	PNF
<i>Cirsium ochrocentrum</i>	yellowspine thistle	UPL		BPNF
<i>Cirsium</i> spp.	thistles	FAC+*	NI	ABINF*
<i>Cirsium undulatum</i>	wavy-leaf thistle	FACU	FACU,FAC	BPNF
<i>Cirsium vulgare</i>	bull thistle	UPL	UPL,FAC	BIF
<i>Comandra umbellata</i>	umbellate bastard toad-flax	UPL,FACU	UPL	PN+F
<i>Convolvulus</i> spp.	field bindweed	FAC*	NI	PNF*
<i>Convolvulus arvense</i>	field bindweed	FAC*	NI	PNF*
<i>Conyza canadensis</i>	Canada horseweed	FACU-	UPL,FAC	ANF
<i>Conyza ramosissima</i>	dwarf horseweed	FAC*	NI	ANF
<i>Coreopsis tinctoria</i>	golden tickseed	FAC	FACU,FAC	ANF
<i>Crepis runcinata</i>	hawksbeard dandelion	FAC	FACU,FACW	PNF
<i>Cyperus acminatus</i>	short-point flatsedge	OBL	OBL	ABPNGL
<i>Cyperus aristatus</i>	awned flatsedge	OBL	FACW+,OBL	ANGL
<i>Cyperus erythrorhizos</i>	redrooted cyperus	OBL	FACW+,OBL	APNEGL
<i>Cyperus esculentes</i>	chufa	FACW	FAC,FACW	PNGL
<i>Cyperus lupulinus</i>	Houghton flatsedge	FACU		PNG
<i>Dalea candida</i>	white prairie clover	UPL		PNF
<i>Dalea purpurea</i>	violet prairie clover	UPL		PNF

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Dalea villosa</i>	silky prairie clover	UPL		PNFHS
<i>Delphinium carolinianum</i>	Carolina larkspur	UPL		PNF
<i>Delphinium virescens</i>	prairie larkspur	UPL		PNF
<i>Desmanthus illinoensis</i>	prairie bundleflower	FACU	UPL,FAC	PNF
<i>Desmodium canadense</i>	showy tick-trefoil	FAC	FACU,FAC	PNF
<i>Desmodium illinoense</i>	Illinois tick-trefoil	UPL		PNF
<i>Dianthus armeria</i>	Deptford pink	UPL		PNF
<i>Dichanthelium acuminatum</i>	panic grass	FAC,FACW	FAC	PNG
<i>Dichanthelium oligosanthes</i>	Heller's witchgrass	FACU	FACU,FAC	PNG
<i>Digitaria sanguinalis</i>	hairy crabgrass	FACU	FACU-,FAC-	AIG
<i>Echinacea angustifolia</i>	blacksamson echinacea	UPL		PNG
<i>Echinochloa crusgalli</i>	barnyard grass	FACW	FACU,FACW	AIG
<i>Echinochloa muricata</i>	rough barnyard grass	OBL	FAC,OBL	ANG
<i>Elatine rubella</i>	red waterwort			ANF
<i>Eleocharis compressa</i>	flat-stem spikerush	FACW	FACW, FACW+	PNEGL
<i>Eleocharis erythropoda</i>	bald spikerush	OBL	OBL	PNGL
<i>Eleocharis obtusa</i>	blunt spikerush	OBL	OBL	APNEGL
<i>Eleocharis ovata</i>	ovate spikerush	OBL	OBL	ANEGL
<i>Eleocharis palustris</i>	creeping spikerush	OBL	OBL	PNEGL
<i>Eleocharis smallii</i>	Small's spikerush	OBL	OBL	PNGL
<i>Elymus canadensis</i>	nodding wild rye	FACU	FACU, FAC+	PNG
<i>Elymus smithii</i>	western wheatgrass	FACU		PNG
<i>Elymus trachycaulus</i>	slender wheatgrass	FACU		PNG
<i>Elymus virginiana</i>	Virginia wild rye	FAC	FAC,FACW	PNG
<i>Eragrostis spectabilis</i>	purple lovegrass	FACU	UPL,FACU	PNG
<i>Erigeron annuus</i>	white-top fleabane	FACU	FACU,FAC	ANF
<i>Erigeron philadelphicus</i>	Philadelphia fleabane	FACU	FACU,FAC	ANF
<i>Erigeron strigosus</i>	prairie fleabane	FAC	FACU,FAC	ANF
<i>Eupatorium perfoliatum</i>	common boneset	OBL	FACW+,OBL	PNF
<i>Euphorbia cyathophora</i>	fire-on-the-mountain	UPL		NAPF
<i>Euphorbia dentata</i>	toothed spurge	UPL		AIF
<i>Euphorbia maculata</i> ( <i>Chamaesyce maculata</i> )	spotted broomspurge	FACU-	UPL,FACU	ANF
<i>Euphorbia marginata</i>	snow-on-the-mountain	UPL,FACU	FACU	ANF
<i>Euphorbia nutans</i>	eyebane broomspurge	FACU-,FACU	FACU-	AIF

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Festuca arundinacea</i>	Kentucky fescue	FACU	UPL,FACW–	PIG
<i>Festuca saximontana</i>	Rocky Mountain fescue			NPG
<i>Fragaria virginiana</i>	Virgina strawberry	UPL,FAC	FACU	PNF
<i>Fraxinus pennsylvanica</i>	green ash	FACW	FAC,FACW	NT
<i>Gaillardia pulchella</i>	blanketflower	UPL		PNF
<i>Galium aparine</i>	catchweed bedstraw	FACU	FACU,FAC–	ANF
<i>Gaura coccinea</i>	scarlet gaura	UPL		PNS/F
<i>Gaura parviflora</i>	velvet-leaf butterfly-weed	FACU?	NI	ANF
<i>Gleditsia triacanthos</i>	honey-locust	FAC	FACU,FAC	NTS
<i>Glyceria striata</i>	fowl mannagrass	OBL	OBL	PNEG
<i>Glycine max</i>	soybean	UPL*	NI	AI*
<i>Glycyrrhiza lepidota</i>	American licorice	FACU	UPL,FAC+	PNF
<i>Grindelia squarrosa</i>	curly-cup gumweed	FACU–	UPL,FACU	ABPNF
<i>Gutierrezia sarothrae</i>	broom snakeweed	UPL		PNS/F
<i>Hedeoma hispida</i>	rough false pennyroyal	UPL		ANF
<i>Helianthus annuus</i>	common sunflower	FACU	FACU,FAC	ANF
<i>Helianthus grosseserratus</i>	saw-tooth sunflower	FACW	FAC,FACW	PNF
<i>Helianthus maximiliani</i>	Maximilian's sunflower	UPL	UPL,FACU	PNF
<i>Helianthus petiolaris</i>	prairie sunflower	UPL		ANF
<i>Helianthus rigidus</i>	stiff sunflower	UPL		PNF
<i>Hesperis matronalis</i>	dames rocket			IBPF
<i>Hesperostipa spartea</i>	porcupinegrass	UPL		PNG
<i>Hesperostipa viridula</i>	green needlegrass	UPL		NG
<i>Heteranthera limosa</i>	blue mud-plantain	OBL	OBL	ANEF
<i>Heteranthera multiflora</i>	bouquest mud-plantain	OBL		ANF
<i>Heterotheca villosa</i>	hairy goldaster	UPL		PNF
<i>Hieracium longipilum</i>	hairy hawkweed	FACU		PNF
<i>Hordeum jubatum</i>	fox-tail barley	FACW	FAC,FACW	PNG
<i>Hordeum pusillum</i>	little barley	FAC	FACU,FAC	ANG
<i>Hypoxis hirsuta</i>	eastern yellow stargrass	FAC,FACW	FACW	PNF
<i>Ipomoea hederacea</i>	ivyleaf morning-glory	FACU,FAC	FACU	AIV
<i>Ipomoea leptophylla</i>	bush morning-glory	UPL		PNF
<i>Ipomoea pandurata</i>	wild sweet-potato vine	FACU,FAC–	FAC–	PNF
<i>Ipomoea purpurea</i>	common morning-glory	UPL,FAC	FACU	AIV

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Juncus dudleyi</i>	Dudley rush			PNG
<i>Juncus interior</i>	inland rush	FAC	FACU,FACW	PNGL
<i>Juncus</i> spp.	rush	FACW*	NI	PNGL*
<i>Juniperus virginiana</i>	eastern red cedar	FACU-	FACI-,FACU	NT
<i>Kochia scoparia</i>	Mexican summer-cypress	FACU	UPL,FAC	AIF
<i>Koeleria macrantha</i>	prairie Junegrass	UPL		PNG
<i>Kuhnia eupatorioides</i>	false boneset	FAC,FACU*	NI	PNF*
<i>Lactuca canadensis</i>	tall yellow lettuce	FACU	FACU-,FAC+	ABNF
<i>Lactuca ludoviciana</i>	biannual lettuce	FAC	UPL,FAC	BPNF
<i>Lactuca serriola</i>	prickly lettuce	FAC	FACU,FAC	ABIF
<i>Lactuca tatarica</i>	blue lettuce	FAC		BPNF
<i>Leersia oryzoides</i>	rice cutgrass	OBL	OBL	PNG
<i>Lemna minor</i>	lesser duckweed	OBL	OBL	PN/F
<i>Lemna trisulca</i>	star duckweed	OBL	OBL	PN/F
<i>Lepidium densiflorum</i>	common pepperweed	FAC		ABNF
<i>Lepidium virginicum</i>	Virginia pepperweed	UPL,FAC-	FACU	ABNF
<i>Leptochloa fascicularis</i>	bearded sprangletop	OBL	FACW,OBL	ANG
<i>Lespedeza capitata</i>	round-head bushclover	UPL	UPL,FACU	PNF
<i>Liatris punctata</i>	dotted gayfeather	FACU		PNF
<i>Liatris pycnostachya</i>	cattail gayfeather	FAC	FACU,FAC+	PNF
<i>Limosella aquatica</i>	northern mudwort	OBL	OBL	APNEF
<i>Lindernia dubia</i>	yellowseed false pimpernel	OBL	OBL	ANF
<i>Linum rigidum</i>	stiffstem flax			APNF
<i>Lithospermum incisum</i>	narrow-leaved puccoon	UPL		PNF
<i>Lomatium foeniculaceum</i>	wild parsley	UPL		PNF
<i>Lotus corniculatus</i>	bird's-foot trefoil	FACU	FACU-,FAC	PIF
<i>Lotus purshianus</i>	prairie trefoil	FACU*	NI	ANF*
<i>Lotus unifoliolatus</i>	American bird's-foot trefoil			ANF
<i>Ludwigia palustris</i>	marsh seedbox	OBL	OBL	PNEF
<i>Lycopus americanus</i>	American bugleweed	OBL	OBL	PNF
<i>Lygodesmia juncea</i>	rush skeletonplant	UPL		PNF
<i>Lythrum salicaria</i>	purple loosetrife	FACW+,OBL	OBL	PIF
<i>Marsilea vestita</i>	hairy water fern	OBL	OBL	PNEP3
<i>Medicago sativa</i>	alfalfa			APIF

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Melilotus alba</i>	white sweetclover	FACU	FACU–,FACU+	ABIF
<i>Melilotus officinalis</i>	yellow sweetclover	FACU	FACU–,FACU+	ABIF
<i>Mirabilis linearis</i>	narrowleaf four o'clock			PNFHS
<i>Mirabilis nyctaginea</i>	heartleaf four o'clock	UPL	UPL,FACU	PNF
<i>Mollugo verticillata</i>	green carpet-weed	FAC	FAC–,FAC	ANF
<i>Monarda fistulosa</i>	wild bergamot	UPL,FAC+	FACU–	PNF
<i>Morus alba</i>	white mulberry	UPL,FAC	FAC	IT
<i>Morus rubra</i>	red mulberry	FACU	FACU, FAC	NT
<i>Muhlenbergia mexicana</i>	Mexican muhly	FACW	FAC,FACW	PNG
<i>Muhlenbergia racemosa</i>	green muhly	FACW	FACU,FACW	PNG
<i>Myosurus minimus</i>	tiny mouse-tail	FACW	FACW–OBL	ANF
<i>Nepeta cataria</i>	catnip	FACU	FACU–,FACW–	PIF
<i>Oenothera biennis</i>	common evening-primrose	FACU	FACU–,FACU+	BIF
<i>Oenothera villosa</i>	hairy evening-primrose	FAC	FACU,FACW	BPNF
<i>Oligoneuron rigidum</i>	goldenrod	UPL		PNF
<i>Oxalis dillenii</i>	gray-green woodsorrel	FACU*	NI	PNF*
<i>Oxalis stricta</i>	yellow woodsorrel	FACU*		PNF
<i>Oxalis violacea</i>	violet woodsorrel	UPL		PNF
<i>Panicum capillare</i>	witchgrass	FAC	FACU,FAC	ANG
<i>Panicum dichotomiflorum</i>	fall panic grass	FAC	FAC,FACW	ANG
<i>Panicum virgatum</i>	switchgrass	FAC	FAC,FACW	PNG
<i>Parietaria pensylvanica</i>	Pennsylvania pellitory	FAC	FACU–,FACW–	ANF
<i>Penstemon grandiflorus</i>	large-flower beardtongue	UPL		PNF
<i>Phalaris arundinacea</i>	reed canarygrass	FACW+	FACW,OBL	PNG
<i>Phleum pratense</i>	timothy	FACU	FACU	PIG
<i>Phyla cuneifolia</i>	wedgeleaf fog-fruit	FAC		PNS/F
<i>Physalis heterophylla</i>	clammy groundcherry	UPL		PNF
<i>Physalis longifolia</i>	longleaf groundcherry	UPL		PNF
<i>Physalis</i> spp.	groundcherry	FAC*	NI	AP*
<i>Physalis virginiana</i>	Virginia groundcherry	UPL		PNF
<i>Poa compressa</i>	Canada bluegrass	FACU	FACU–,FAC	PIG
<i>Poa pratensis</i>	Kentucky bluegrass	FACU	FACU,FAC–	PNG
<i>Polygala alba</i>	white milkwort			PNF
<i>Polygonum arenastrum</i>	knotweed	UPL*	NI	APNF*

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Polygonum bicornne</i> ( <i>P. pennsylvanicum</i> )	pink smartweed	FACW+	FACW-,OBL	ANEF
<i>Polygonum coccineum</i>	water smartweed	OBL	OBL	PNE/F
<i>Polygonum hydropiper</i>	swamp smartweed	OBL	OBL	PNEF
<i>Polygonum lapathifolium</i>	curlytop knotweed	OBL	FAC,OBL	ANF
<i>Polygonum persicaria</i>	spotted ladysthumb	OBL	FAC,OBL	AIF
<i>Polygonum punctatum</i>	dotted smartweed	OBL	FACW,OBL	PNEF
<i>Polygonum ramosissimum</i>	bushy knotweed	FAC	FACU-,FACW	ANF
<i>Populus deltoides</i>	eastern cottonwood	FAC	FAC,FACW	NT
<i>Portulaca oleracea</i>	common purslane	FAC	FACU,FAC	AN\$F
<i>Potamogeton gramineus</i>	variable pondweed	OBL	OBL	PNZF
<i>Potamogeton nodosus</i>	longleaf pondweed	OBL	OBL	PNZF
<i>Potentilla arguta</i>	tall cinquefoil	FACU	UPL,FACU+	PNF
<i>Potentilla norvegica</i>	Norwegian cinquefoil	FAC	FACU,FAC	ABPNF
<i>Potentilla recta</i>	sulphur cinquefoil	FACU*	NI	PNF*
<i>Prenanthes racemosa</i>	glaucous rattlesnake-root	FAC	FACU-,FACW	PNF
<i>Prunella vulgaris</i>	heal-all	FAC	FACU,FACW	PIF
<i>Prunus americana</i>	American plum	UPL	UPL,FACU	NST
<i>Prunus virginiana</i>	chokecherry	FACU	FACU-,FAC	NST
<i>Psoralea argophylla</i>	silver-leaf scurfpea	FACU*	NI	PNF*
<i>Psoralea tenuiflora</i>	few-flowered scurfpea	UPL		PNF
<i>Psoralidium lanceolatum</i>	lemon scurfpea	UPL		PNF
<i>Pulsatilla patens</i>	pasqueflower	UPL		PNF
<i>Ranunculus flabellaris</i>	yellow water buttercup	OBL	OBL	PNEF
<i>Ranunculus longirostris</i>	longbeak buttercup	OBL	OBL	PNZ/F
<i>Ranunculus pennsylvanicus</i>	Pennsylvania buttercup	FACW,OBL	OBL	APNEF
<i>Ranunculus sceleratus</i>	cursed buttercup	OBL	OBL	APNEF
<i>Ratibida columnifera</i>	upright prairie coneflower	UPL		PNF
<i>Rhus glabra</i>	smooth sumac	UPL		NT
<i>Riccia fluitans</i>	liverwort			
<i>Robinia pseudoacacia</i>	black locust	UPL	UPL,FAC	NT
<i>Rorippa palustris</i>	bog yellowcress	OBL	FAC,OBL	ANEF
<i>Rorippa sessiliflora</i>	stalkless yellowcress	OBL	FACW+,OBL	ANEF
<i>Rorippa sinuata</i>	spreading yellowcress	FACW	FAC+,FACW	PNF
<i>Rosa arkansana</i>	wild prairie rose	NI	FAC?	NSH

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Rosa multiflora</i>	multiflora rose	UPL,FACU	UPL	IS
<i>Rosa woodsii</i>	Woods' rose	UPL,FAC-	FACU	NS
<i>Rudbeckia hirta</i>	black-eyed susan	FACU-,FACU	FACU	BPNF
<i>Rumex altissimus</i>	pale dock	FAC	FAC,FACW+	PNF
<i>Rumex crispus</i>	curly dock	FACW	FACU,FACW	PIF
<i>Sagittaria brevirostra</i>	shortbeak arrowhead	OBL	OBL	PNEF*
<i>Sagittaria calycina</i>	hooded arrowhead	OBL	OBL	PNEF
<i>Sagittaria cuneata</i>	arumleaf arrowhead	OBL	OBL	PNEF*
<i>Sagittaria graminea</i>	grassy arrowhead	OBL	OBL	PNEF
<i>Sagittaria latifolia</i>	broadleaf arrowhead	OBL	OBL	PNEF*
<i>Sagittaria longiloba</i>	longbarb arrowhead	OBL	OBL	PNEF
<i>Sagittaria rigida</i>	sessilefruit arrowhead	OBL	OBL	PNEF
<i>Salix amygdaloides</i>	peach-leaf willow	FACW	FACW	NT
<i>Salsola kali</i>	Russian thistle	FACU-,FACU+	FACU	AIF
<i>Salvia azurea</i>	blue sage	UPL		PNF
<i>Schizachyrium scoparium</i>	little bluestem	FACU	FACU-,FACU+	PNG
<i>Schoenoplectus fluviatilis</i>	river bulrush	OBL	OBL	PNEGL
<i>Schoenoplectus heterochaetus</i>	slender bulrush	OBL	OBL	PNEGL
<i>Schoenoplectus pungens</i>	three-square bulrush	FACW+,OBL	OBL	PNEGL
<i>Scirpus acutus</i>	hard-stem bulrush	OBL	OBL	PNEGL
<i>Scirpus validus</i>	soft-stem bulrush	OBL	OBL	PNEGL
<i>Scutellaria parvula</i>	small skullcap	UPL,FACU	FACU	PNF
<i>Senecio plattensis</i>	prairie groundsel	UPL,FACU	FACU	BPNF
<i>Setaria glauca</i>	yellow foxtail grass	FACU,FAC	FAC	AIG
<i>Setaria pumila</i>	yellow bristle grass	FAC	FACU,FAC	AIG
<i>Setaria viridis</i>	green foxtail	FAC*	NI	ANG*
<i>Silene antirrhina</i>	sleepy silene	UPL		ANF
<i>Silphium integrifolium</i>	wholeleaf rosinweed	UPL		PNF
<i>Silphium laciniatum</i>	compassplant	UPL		PNF
<i>Silphium perfoliatum</i>	cup-plant	FACU,FACW	FAC	PNF
<i>Sinapis alba</i>	white mustard	UPL		AIF
<i>Sinapis arvensis</i>	corn mustard			AIF
<i>Sisyrinchium campestre</i>	blue-eyed grass	UPL		PNF
<i>Sisyrinchium montanum</i>	strict blue-eyed grass	FACU,FACW	FAC	PNF

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Solanum carolinense</i>	Carolina nightshade	UPL,FACU	UPL	NSF
<i>Solanum interius</i>	inland nightshade			ANF
<i>Solanum ptycanthum</i>	black nightshade	FAC+*	NI	ANF*
<i>Solanum rostratum</i>	buffalobur nightshade	FAC*	NI	ANF*
<i>Solidago canadensis</i>	Canada goldenrod	FACU	FACU,FACU+	PNF
<i>Solidago gigantea</i>	giant goldenrod	FAC,FACW	FACW	PNF
<i>Solidago graminifolia</i> ( <i>Euthamia graminifolia</i> )	flat-top goldenrod	UPL		PNF
<i>Solidago missouriensis</i>	Missouri goldenrod	FACU*	NI	PNF
<i>Solidago rigida</i>	stiff goldenrod	FACU	UPL,FACU	PNF
<i>Sonchus arvensis</i>	field sowthistle	UPL,FAC	FAC	PIF
<i>Sonchus oleraceus</i>	common sowthistle	UPL,FACU	FACU	AIF
<i>Sorghastrum nutans</i>	Indiangrass	FACU	UPL,FACW	PNG
<i>Sorghum bicolor</i>	milo	UPL*	NI	AIG*
<i>Sparganium eurycarpum</i>	giant burreed	OBL	OBL	PNEF
<i>Spartina pectinata</i>	prairie cordgrass	FACW	FACW,OBL	PNG
<i>Sphaeralcea coccinea</i>	false red mallow			BPNFHS
<i>Sphenopholis obtusata</i>	prairie wedgegrass	FAC-,FACW+	FACW	APNG
<i>Spirodela polyrrhiza</i>	greater duckweed	OBL	OBL	PNF
<i>Sporobolus asper</i>	tall dropseed	UPL,FACU	FACU	PNG
<i>Sporobolus cryptandrus</i>	sand dropseed	FACU-	UPL,FACU	PNG
<i>Sporobolus vaginiflorus</i>	poverty dropseed	FACU	UPL,FACU	ANG
<i>Stachys palustris</i>	marsh hedgenettle	FACW,OBL	OBL	PIF
<i>Strophostyles leiosperma</i>	slickseed fuzzybean			ANFV
<i>Symphoricarpos occidentalis</i>	western snowberry			PNS
<i>Symphoricarpos orbiculatus</i>	coralberry	UPL,FAC-	FACU-	NS
<i>Symphyotrichum ericoides</i>	white heath aster	FACU	UPL,FACU	PNF
<i>Symphyotrichum lanceolatum</i>	panicled aster	FACW	FACW	PNF
<i>Symphyotrichum novae-angliae</i>	New England aster	FACW	FACW-, FACW	PNF
<i>Symphyotrichum praealtum</i> var. <i>nebraskense</i>	Nebraska aster	FACW	FACW-, FACW	PNF
<i>Tamarix ramosissima</i>	saltcedar	FAC,FACW	FACW	IT
<i>Taraxacum officinale</i>	common dandelion	FACU	FACU-,FACU+	PIF
<i>Teucrium canadense</i> ( <i>T. occidentale</i> )	American germander	FACW	FAC+,FACW	PNEF
<i>Thalictrum dioicum</i>	early meadow-rue	FACU+,FACW	FAC	PNF
<i>Thinopyrum intermedium</i>	intermediate wheatgrass	UPL		PIG

<i>Scientific Name</i>	<i>Common Name</i>	<i>Region 5 Wetland Indicator</i>	<i>National Wetland Indicator</i>	<i>Habit</i>
<i>Thlaspi arvense</i>	field pennycress	FACU?	NI	AIF
<i>Toxicodendron radicans</i>	poison ivy	FACU,FACW	FACU	NWVS
<i>Toxicodendron rydbergii</i>	Rydberg poison ivy	FACU,FACW	FAC	NHS
<i>Tradescantia bracteata</i>	longbract spiderwort	UPL,FAC	FAC	PNF
<i>Tragopogon dubius</i>	goatsbeard	FACU*	NI	BIF*
<i>Tribulus terrestris</i>	puncturevine	UPL		AIF
<i>Trifolium pratense</i>	red clover	FACU	FACU-,FAC	BPIF
<i>Trifolium repens</i>	white clover	FACU	FACU-,FAC	PIF
<i>Triodanis perfoliata</i>	claspleaf Venus' looking-glass	UPL,FAC	FAC	ANF
<i>Tripsacum dactyloides</i>	eastern gama grass	FAC,FACW	FAC	PNG
<i>Typha angustifolia</i>	narrowleaf cattail	OBL	OBL	PNEF
<i>Typha latifolia</i>	broadleaf cattail	OBL	OBL	PNEF
<i>Typha × glauca</i>	hybrid cattail	OBL	OBL	PNEF
<i>Ulmus americana</i>	American elm	FAC,FACW	FAC	NT
<i>Ulmus pumila</i>	Siberian elm	FACU*	NI	IT*
<i>Utricularia vulgaris</i>	bladderwort	OBL	OBL	PNZF
<i>Verbascum thapsus</i>	common mullein	UPL		PNF
<i>Verbena bracteata</i>	prostrate vervain	FACU	UPL,FACW	APNF
<i>Verbena hastata</i>	blue vervain	FAC,FACW+	FACW	PNF
<i>Verbena urticifolia</i>	white vervain	UPL,FAC+	UPL	APNF
<i>Verbena stricta</i>	hoary vervain	FAC,FACU*	NI	PNF*
<i>Vernonia baldwinii</i>	Baldwin's ironweed	UPL,FACW-	FACW-	PNF
<i>Vernonia fasciculata</i>	prairie ironweed	FAC	FAC,FACW	PNF
<i>Veronica peregrina</i>	purslane speedwell	FACU-,OBL	OBL	ANEF
<i>Vicia americana</i>	American purple vetch	FAC?	NI	PNFV
<i>Viola pedatifida</i>	prairie violet	UPL,FACU	FACU	PNF
<i>Viola pratincola</i>	blue prairie violet	FACU,FAC	FAC-	PNF
<i>Wolffia columbiana</i>	Columbian watermeal	OBL	OBL	PN/F
<i>Xanthium strumarium</i>	rough cocklebur	FAC	FAC-, FAC+	ANF
<i>Yucca gloriosa</i>	moundlily yucca	FAC	NI	NT
<i>Zea mays</i>	corn			AIF



# Appendix I

## *List of Amphibians, Reptiles, and Mammals*

### AMPHIBIANS

<i>Common Name</i>	<i>National Status in Nebraska</i>
American toad	critically imperiled
Great Plains toad	secure
Woodhouse's toad	secure
northern cricket frog	secure
Cope's gray treefrog	secure
western chorus frog	secure
Great Plains narrowmouth toad	imperiled
plains leopard frog	secure
bullfrog	secure
northern leopard frog	secure
smallmouth salamander	critically imperiled
tiger salamander	secure

### REPTILES

<i>Common Name</i>	<i>National Status in Nebraska</i>
snapping turtle	secure
smooth softshell	secure
spiny softshell	secure
yellow mud turtle	vulnerable
painted turtle	secure
Blanding's turtle	apparently secure
false map turtle	vulnerable
western box turtle	secure
lesser earless lizard	secure
greater short-horned lizard	vulnerable
sagebrush lizard	critically imperiled
fence/prairie/plateau lizard	secure
six-lined racerunner	secure
five-lined skink	critically imperiled
many-lined skink	secure
Great Plains skink	vulnerable
prairie skink	secure
slender glass lizard	critically imperiled
glossy snake	imperiled
worm snake	secure
racer	secure
ringneck snake	secure
corn snake	apparently secure

<i>Common Name</i>	<i>National Status in Nebraska</i>
western ratsnake	apparently secure
fox snake	secure
western hognose snake	secure
eastern hognose snake	apparently secure
prairie kingsnake	vulnerable
common kingsnake	imperiled
milk snake	secure
smooth green snake	critically imperiled
coachwhip	vulnerable
northern water snake	secure
gopher snake	secure
Graham's crayfish snake	imperiled
brown snake	vulnerable
redbelly snake	critically imperiled
plains black-headed snake	critically imperiled
western terrestrial garter snake	apparently secure
western ribbon snake	imperiled
plains garter snake	secure
common garter snake	secure
lined snake	secure
copperhead	critically imperiled
western rattlesnake	apparently secure

## MAMMALS

<i>Common Name</i>	<i>National Status in Nebraska</i>
Townsend's big-eared bat	critically imperiled
big brown bat	secure
silver-haired bat	secure
eastern red bat	secure
hoary bat	secure
western small-footed myotis	apparently secure
little brown bat	apparently secure
northern myotis	vulnerable
fringed myotis	critically imperiled
fringe-tailed myotis	critically imperiled
long-legged myotis	imperiled
evening bat	vulnerable
eastern pipistrelle	critically imperiled
Brazilian free-tailed bat	unranked
black-tailed prairie dog	apparently secure
woodchuck	apparently secure
eastern fox squirrel	secure
Wyoming ground squirrel	possibly extirpated
Franklin's ground squirrel	secure
spotted ground squirrel	apparently secure
thirteen-lined ground squirrel	secure
least chipmunk	vulnerable

<i>Common Name</i>	<i>National Status in Nebraska</i>
American beaver	secure
plains pocket gopher	secure
northern pocket gopher	apparently secure
hispid pocket mouse	secure
Ord's kangaroo rat	secure
olive-backed pocket mouse	vulnerable
plains pocket mouse	secure
silky pocket mouse	apparently secure
meadow jumping mouse	secure
prairie vole	secure
meadow vole	secure
woodland vole	vulnerable
house mouse	exotic
muskrat	secure
northern grasshopper mouse	secure
white-footed mouse	secure
deer mouse	exotic
Norway rat	exotic
western harvest mouse	secure
plains harvest mouse	apparently secure
hispid cotton rat	vulnerable
southern bog lemming	apparently secure
northern short-tailed shrew	vulnerable
short-tailed shrew	vulnerable
least shrew	apparently secure
prairie shrew	apparently secure
Merriam's shrew	critically imperiled
eastern mole	secure
Virginia opossum	secure
mule deer	secure
white-tailed deer	secure
black-tailed jackrabbit	secure
white-tailed jackrabbit	secure
European rabbit	exotic
eastern cottontail	secure
coyote	secure
swift fox	imperiled
red fox	secure
bobcat	secure
northern river otter	imperiled
long-tailed weasel	secure
black-footed ferret	extirpated
least weasel	secure
American mink	secure
American badger	secure
striped skunk	secure
eastern spotted skunk	secure
northern raccoon	secure



# Appendix J

## List of Birds

The order of bird species follows the AOU Check-list of North American Birds (American Ornithological Union 2004).

Seasonal abundance codes follow each species name, in order of seasons (spring, summer, fall, winter). The letter “*m*” is used to indicate that the species is not present during a particular season. For example, “American coot *acar*” (abundant in spring, common summer, abundant in fall, and rare in Winter) or “California gull *omom*” (occasional in spring, not present in summer or winter, and occasional in fall).

Nesting species documented at the WPAs are marked with an asterisk (\*) in front of the species name. Species accounts may indicate presence during the nesting season, but may not be marked as nesting. For example, frequent nesters in the area such as northern rough-winged swallow are not marked as nesters because WPAs do not provide preferred nesting habitat. For the most part, cavity nesters are excluded (the red-bellied woodpecker prefers mature, natural woodlands) due to the absence of mature or dying trees that typically provide cavities. The district’s WPAs are managed as grassland–playa lake ecosystems. Nonnative trees and shrubs have been removed at most WPAs. Due to remaining shelterbelts at WPAs, a few woodland nesters are marked as nesting if they are known to use shelterbelts.

### SEASONAL ABUNDANCE DEFINITIONS

Seasons are listed below. Seasonal abundance codes for some species such as shorebirds may be misleading because their fall migration starts in July or August (summer). For example, “pectoral sandpiper *cccm*” (common in spring, summer due to peak migration occurring mid-August, and fall; not present in winter).

### SEASONS

Spring (March–May)

Summer (June–August)

Fall (September–November)

Winter (December–February)

### SEASONAL ABUNDANCE

*a* = abundant (occur in large numbers)

*c* = common (certain to be seen in suitable habitat)

*u* = uncommon (present, but not certain to be seen)

*o* = occasional (seen only a few times during the season)

*r* = rare (seen at intervals of 2–5 years)

*h* = hypothetical (within normal range, but never documented)

*x* = outside normal range (but has been documented)

### SWANS, GEESE, AND DUCKS

black-bellied whistling-duck *xxxxm*

bean goose *xmxxx*

greater white-fronted goose *arao*

emperor goose *xmxxx*

snow goose *aoao*

Ross’ goose *crco*

\*Canada goose *auac*

brant *rxxxm*

trumpeter swan *hxmh*

tundra swan *omxxx*

wood duck *cucm*

\*gadwall *cuco*

Eurasian wigeon *rxxxm*

\*American wigeon *aoao*

American black duck *rmrr*

\*mallard *acac*

\*blue-winged teal *acar*

\*cinnamon teal *uoum*

\*northern shoveler *acar*

\*northern pintail *auao*

garganey *xmxxx*

\*green-winged teal *aoao*

canvasback *umum*

\*redhead *cucm*

ring-necked duck *crum*

greater scaup *umrm*

lesser scaup *cocm*

white-winged scoter *xmxxx*

black scoter *xmxxx*

bufflehead *cmcm*

common goldeneye *omum*

Barrow’s goldeneye *mmxxx*

hooded merganser *urum*

common merganser *omoo*

\*ruddy duck *cucm*

### GALLINACEOUS BIRDS

\*ring-necked pheasant *cccc*

sharp-tailed grouse *rhro*

\*greater prairie-chicken *ooou*

\*northern bobwhite *cccc*

**LOONS**

common loon *omrm*

**GREBES**

Clark's grebe *xmrm*  
 \*pied-billed grebe *cccm*  
 horned grebe *uhrm*  
 red-necked grebe *hmrm*  
 \*eared grebe *cocm*  
 western grebe *rmrm*

**PELICANS**

American white pelican *cocm*

**CORMORANTS**

double-crested cormorant *cucm*

**BITTERN, HERONS, AND EGRETS**

\*American bittern *ucum*  
 \*least bittern *uom*  
 \*great blue heron *cccr*  
 great egret *omom*  
 snowy egret *uum*  
 little blue heron *orm*  
 tricolored heron *hrrm*  
 cattle egret *uoum*  
 \*green heron *cucm*  
 \*black-crowned night-heron *cocm*  
 yellow-crowned night-heron *rmrm*

**IBISES AND SPOONBILLS**

white ibis *hxxm*  
 glossy ibis *xmmm*  
 white-faced ibis *uurm*  
 roseate spoonbill *mxmm*

**NEW WORLD VULTURES**

black vulture *mxmm*  
 turkey vulture *cucm*

**OSPREY, KITES, HAWKS, AND EAGLES**

osprey *omom*  
 white-tailed kite *xmxx*  
 Mississippi kite *xmxx*  
 bald eagle *cmco*  
 \*northern harrier *cucu*  
 sharp-shinned hawk *uruu*  
 Cooper's hawk *umur*  
 northern goshawk *rmrr*  
 red-shouldered hawk *mrrx*  
 broad-winged hawk *mxmm*  
 \*Swainson's hawk *cuur*  
 \*red-tailed hawk *cucu*  
 ferruginous hawk *uxuo*  
 rough-legged hawk *uhuu*  
 golden eagle *omoo*

**FALCONS AND CARACARAS**

\*American kestrel *cccc*  
 merlin *umuo*  
 gyrfalcon *mmmr*  
 peregrine falcon *omor*  
 prairie falcon *omou*

**RAILS**

yellow rail *rmrm*  
 black rail *xmmm*  
 king rail *rmrm*  
 \*Virginia rail *uum*  
 \*sora *cccm*  
 purple gallinule *xmmm*  
 \*common moorhen *uurm*  
 \*American coot *acar*

**CRANES**

\*sandhill crane *crum*  
 common crane *xmmm*  
 whooping crane *omom*

**PLOVERS**

black-bellied plover *umum*  
 American golden-plover *umum*  
 snowy plover *rmrm*  
 semipalmated plover *umum*  
 piping plover *rhrm*  
 \*killdeer *cccm*

**STILTS AND AVOCETS**

\*black-necked stilt *rrrm*  
 \*American avocet *urum*

**SANDPIPERS AND PHALAROPES**

greater yellowlegs *cocm*  
 lesser yellowlegs *cucm*  
 solitary sandpiper *ccum*  
 willet *uoum*  
 spotted sandpiper *cccm*  
 \*upland sandpiper *uum*  
 Eskimo curlew *hmhm*  
 whimbrel *omrm*  
 long-billed curlew *rmrm*  
 Hudsonian godwit *umxx*  
 marbled godwit *umrm*  
 ruddy turnstone *rmrm*  
 red knot *xmrm*  
 sanderling *omom*  
 semipalmated sandpiper *aucm*  
 western sandpiper *room*  
 least sandpiper *cccm*  
 white-rumped sandpiper *coom*  
 Baird's sandpiper *acum*  
 pectoral sandpiper *cccm*  
 sharp-tailed sandpiper *mmxm*  
 dunlin *umum*  
 curlew sandpiper *mxmm*  
 stilt sandpiper *accm*

buff-breasted sandpiper *uuum*  
 ruff *xmxx*  
 short-billed dowitcher *uurm*  
 long-billed dowitcher *aucm*  
 \*Wilson's snipe (common) *cocm*  
 American woodcock *rmm*  
 \*Wilson's phalarope *aocm*  
 red-necked phalarope *rmm*  
 red phalarope *mxxm*

## SKUAS, JAEGERS, GULLS, AND TERNS

laughing gull *xmmm*  
 Franklin's gull *arcm*  
 Bonaparte's gull *rmm*  
 mew gull *rmmm*  
 ring-billed gull *acco*  
 California gull *omom*  
 herring gull *umuu*  
 common tern *umum*  
 Forster's tern *cucm*  
 least tern *rrrm*  
 \*black tern *acam*

## PIGEONS AND DOVES

rock pigeon *cccc*  
 European collared-dove *oooo*  
 white-winged dove *mrrm*  
 \*mourning dove *aaaa*  
 Inca dove *mxxm*

## CUCKOOS AND ANIS

black-billed cuckoo *oom*  
 \*yellow-billed cuckoo *uuum*

## BARN OWLS

barn owl *oooo*

## TYPICAL OWLS

eastern screech-owl *uuuu*  
 \*great horned owl *cucu*  
 snowy owl *mmmx*  
 \*burrowing owl *uuum*  
 barred owl *rrrr*  
 long-eared owl *oroo*  
 short-eared owl *uruu*  
 boreal owl *mxxm*

## NIGHTJARS

\*common nighthawk *cccm*  
 chuck-will's-widow *mrrm*

## SWIFTS

chimney swift *aacm*

## HUMMINGBIRDS

ruby-throated hummingbird *oom*  
 broad-tailed hummingbird *hrm*  
 rufous hummingbird *mrrm*

## KINGFISHERS

belted kingfisher *uuur*

## WOODPECKERS

Lewis' woodpecker *rxxm*  
 \*red-headed woodpecker *cccc*  
 red-bellied woodpecker *uuuu*  
 \*downy woodpecker *cccc*  
 hairy woodpecker *cccc*  
 \*northern flicker *cccc*

## TYRANT FLYCATCHERS

olive-sided flycatcher *omom*  
 eastern wood-pewee *oom*  
 yellow-bellied flycatcher *rmm*  
 Acadian flycatcher *rmm*  
 alder flycatcher *rmm*  
 willow flycatcher *orom*  
 least flycatcher *umum*  
 Hammond's flycatcher *mxxm*  
 eastern phoebe *uuum*  
 Say's phoebe *urrm*  
 \*great crested flycatcher *cocm*  
 \*western kingbird *cccm*  
 \*eastern kingbird *cccm*  
 \*scissor-tailed flycatcher *rxrm*

## SHRIKES

\*loggerhead shrike *cccc*  
 northern shrike *mmuu*

## VIREOS

white-eyed vireo *mrrm*  
 \*Bell's vireo *uoum*  
 yellow-throated vireo *rxrm*  
 \*warbling vireo *cucm*  
 Philadelphia vireo *umum*  
 red-eyed vireo *urum*

## CROWS, JAYS, AND MAGPIES

\*blue jay *cccc*  
 pinyon jay *mmmx*  
 \*black-billed magpie *uuuu*  
 \*American crow *aoac*

## LARKS

\*horned lark *acaa*

## SWALLOWS

purple martin *cucm*  
 tree swallow *cocm*  
 violet-green swallow *xmmm*  
 northern rough-winged swallow *cccm*  
 bank swallow *uuum*  
 cliff swallow *ccam*  
 barn swallow *aaam*

**TITMICE AND CHICKADEES**

\*black-capped chickadee *cccc*  
tufted titmouse *xxxm*

**NUTHATCHES**

red-breasted nuthatch *mxmo*  
white-breasted nuthatch *oooo*

**CREEPERS**

brown creeper *mooo*

**WRENS**

Carolina wren *roor*  
Bewick's wren *xmmm*  
\*house wren *cccc*  
winter wren *rmro*  
\*sedge wren *rumr*  
\*marsh wren *uuur*

**KINGLETS**

golden-crowned kinglet *umur*  
ruby-crowned kinglet *omom*

**OLD WORLD WARBLERS**

blue-gray gnatcatcher *orom*

**THRUSHES**

\*eastern bluebird *cucr*  
mountain bluebird *omor*  
Townsend's solitaire *rmro*  
veery *rmmm*  
gray-cheeked thrush *umrm*  
Swainson's thrush *cmum*  
hermit thrush *rmrm*  
wood thrush *rmrm*  
\*American robin *acac*

**MIMIC THRUSHES**

\*gray catbird *uuur*  
northern mockingbird *uumm*  
\*brown thrasher *cccc*

**STARLINGS**

\*European starling *aaac*

**WAGTAILS AND PIPITS**

American (water) pipit *cmcr*  
Sprague's pipit *omom*

**WAXWINGS**

Bohemian waxwing *mmmr*  
cedar waxwing *cucu*

**WOOD WARBLERS**

golden-winged warbler *rmmm*  
Tennessee warbler *cmcm*

orange-crowned warbler *cmcm*  
Nashville warbler *umum*  
northern parula *umrm*  
\*yellow warbler *cccc*  
chestnut-sided warbler *umum*  
magnolia warbler *umum*  
black-throated blue warbler *rmrm*  
yellow-rumped warbler *cmcm*  
black-throated gray warbler *xmmm*  
black-throated green warbler *omom*  
Townsend's warbler *rmrm*  
Blackburnian warbler *omom*  
yellow-throated warbler *rmrm*  
pine warbler *hmrm*  
prairie warbler *rmrm*  
palm warbler *omrm*  
bay-breasted warbler *omom*  
blackpoll warbler *cmrm*  
black-and-white warbler *cmum*  
American redstart *cocm*  
prothonotary warbler *rmrm*  
ovenbird *umum*  
northern waterthrush *umum*  
Louisiana waterthrush *rmhm*  
Kentucky warbler *rmhm*  
Connecticut warbler *rmxh*  
mourning warbler *rmhm*  
MacGillivray's warbler *omrm*  
\*common yellowthroat *cccc*  
hooded warbler *rmrm*  
Wilson's warbler *umum*  
Canada warbler *rmrm*  
yellow-breasted chat *oooo*

**TANAGERS**

summer tanager *rmxm*  
scarlet tanager *omom*

**SPARROWS AND TOWHEES**

green-tailed towhee *rmrm*  
spotted towhee *uoum*  
eastern towhee *umum*  
Cassin's sparrow *xmmm*  
American tree sparrow *aoaa*  
\*chipping sparrow *aoar*  
clay-colored sparrow *chcr*  
Brewer's sparrow *rmrm*  
field sparrow *cucr*  
\*vesper sparrow *cocx*  
lark sparrow *cccc*  
\*lark bunting *ouom*  
Savannah sparrow *cmch*  
\*grasshopper sparrow *cccc*  
Baird's sparrow *umum*  
Henslow's sparrow *rrrm*  
Le Conte's sparrow *umcm*  
Nelson's sharp-tailed sparrow *rmum*  
fox sparrow *omor*  
song sparrow *cucu*  
Lincoln's sparrow *crum*  
\*swamp sparrow *uoum*

white-throated sparrow *cour*  
 Harris' sparrow *cocu*  
 white-crowned sparrow *cruo*  
 golden-crowned sparrow *rmm*  
 dark-eyed junco *cmca*  
 McCown's longspur *rmm*  
 Lapland longspur *amaa*  
 Smith's longspur *omur*  
 chestnut-collared longspur *omom*  
 snow bunting *rmro*

## CARDINALS, GROSBEAKS, AND ALLIES

\*northern cardinal *cccc*  
 rose-breasted grosbeak *uurm*  
 black-headed grosbeak *crcm*  
 \*blue grosbeak *uum*  
 lazuli bunting *xm xm*  
 indigo bunting *uum*  
 painted bunting *rm xm*  
 \*dickcissel *cacm*

## BLACKBIRDS AND ORIOLES

\*bobolink *cccm*  
 \*red-winged blackbird *aca*

\*eastern meadowlark *uuuo*  
 \*western meadowlark *acac*  
 \*yellow-headed blackbird *aocr*  
 rusty blackbird *cour*  
 \*Brewer's blackbird *cuco*  
 \*common grackle *aaao*  
 \*great-tailed grackle *cccr*  
 \*brown-headed cowbird *acar*  
 orchard oriole *cccm*  
 \*Baltimore oriole *cccm*  
 Scott's oriole *mxmm*

## FINCHES

pine grosbeak *xm xm*  
 purple finch *umuo*  
 \*house finch *cccc*  
 red crossbill *mrmo*  
 common redpoll *ormo*  
 pine siskin *uouu*  
 \*American goldfinch *cccc*  
 evening grosbeak *rmr*

## OLD WORLD SPARROWS

\*house sparrow *aaaa*



# Appendix K

## *Rainwater Basin Vegetation Classification*

The National Vegetation Classification System (NVCS) for the United States was used to classify vegetation communities at the Rainwater Basin Wetland Management District in 2004. The NVCS is the system mandated by the USGS–National Park Service Vegetation Mapping Program. In addition, the Federal Geographic Data Committee (FGDC) adopted the NVCS to the formation level as a standard for federal agencies (FGDC 1997). This system provides a national (versus regional, state, or local) vegetation classification system that facilitates resource stewardship by ensuring the same plant associations get the same names throughout the Refuge System. The strengths of the NVCS include the following:

- is vegetation based
- uses a systematic approach to classify a continuum
- emphasizes natural and existing vegetation
- uses a combined physiognomic-floristic hierarchy
- identifies vegetation units based on both qualitative and quantitative data
- is appropriate for mapping at multiple scales

The NVCS was established primarily by The Nature Conservancy and is being implemented and updated by NatureServe in support of the network of natural heritage programs (Grossman et al. 1998). Development and refinement of the classification is an ongoing process and proposed revisions are reviewed both locally and nationally. The Nature Conservancy published two volumes describing the classification of United States vegetation as of April 1997 (Grossman et al. 1998). This publication can be found on the Internet at <<http://www.natureserve.org/publications/icec/index.html>>. NatureServe posts regular updates to the list of plant associations in the United States and Canada on their online database server at <<http://www.natureserve.org/explorer>>.

### **CLASSIFICATION PROCEDURE**

The procedure for classifying vegetation followed guidelines described in the “vegetation classification standard” (FGDC 1997), which was derived from the NVCS. The NVCS is a species-based, hierarchical system with seven levels (Grossman et al. 1998). The highest—“coarse”—levels of the hierarchy have a broad geographic perspective and use physiognomic features to distinguish among groups of plant communities. The lowest—“finest”—levels have a local and site-specific perspective and are based on

floristic features (figure I-1). The finest level (association) was used in the “Rainwater Basin Vegetation and Monitoring Project.”

The association is defined as “a plant community of definite floristic composition, uniform habitat conditions, and uniform physiognomy” (see Flahault and Schroter 1910 in Moravec 1993). Associations are separated from alliances through the use of total floristic composition and are named by the most dominant or indicator species. If two or more dominant species occur in the same stratum, a dash (–) is used between the names. If the species occur in different strata, a slash (/) is used. Parentheses ( ) indicate that a diagnostic species is not always present.

Alliances are physiognomically uniform groups of plant associations that share dominant or diagnostic species, usually found in the uppermost stratum of the vegetation. For forested types, the alliance is roughly equivalent to the “cover type” of the Society of American Foresters. Alliances also include nonforested types.

Unlike classifications based on habitat types or potential vegetation, the NVCS strives to describe existing vegetation, whether natural or cultural vegetation. However, due in part to the conservation focus of The Nature Conservancy and NatureServe, the classification of natural vegetation types is often better developed than that of cultural or modified types. The NVCS is unique in that the association is the basic unit, with the higher levels of the hierarchy representing aggregations of units in the lower levels. This differs from other types that work from the top down.


### **PREPARING THE DATA FOR ANALYSIS**

The vegetation classification for the Rainwater Basin began with the vegetation mapping team completing a literature review of all associations identified in Nebraska and the Great Plains region to identify associations that were present in the basin. From this list, the associations that occurred at WPAs were identified. This was done by bringing together all biologists working in the area and having them identify the known associations, as well as creating map units to delineate species of management concern. The following summaries show the lineage for each plant community for the basin starting with the NVCS community names (see also table I-1).

# Land Cover Classifications

## National Vegetation Classification System (NVCS)

### Physiognomic levels



### Floristic levels



Figure I-1. An example of the NVCS physiognomic-floristic classification hierarchy.

Table I-1.

Formation Name	Alliance Name	Alliance Code	Unique Identifier	Association Name	Association Code	Common Name
<b>Formation Class/Subclass: Shrubland/Deciduous Shrubland</b>						
Temporarily flooded, cold-deciduous shrubland	<i>Salix (exigua, interior)</i> temporarily flooded shrubland	III.B.2.N.d.6	CEGL001203	<i>Salix exigua</i> /mesic graminoids shrubland	d.6	sandbar willow/mesic graminoids shrubland
Temporarily flooded, cold-deciduous shrubland	<i>Cornus sericea</i> temporarily flooded shrubland	III.B.2.N.d.27	CEGL005219	<i>Cornus drummondii</i> – <i>Amorpha fruticosa</i> – <i>Cornus sericea</i> shrubland	d.27	sumac–dogwood shrubland
<b>Formation Class/Subclass: Woodland/Deciduous Woodland</b>						
Temporarily flooded, cold-deciduous woodland	<i>Populus deltoides</i> temporarily flooded woodland	II.B.2.N.b.4	CEGL000659	<i>Populus deltoides</i> –( <i>Salix amygdaloides</i> )/ <i>Salix (exigua, interior)</i> woodland	b.4	cottonwood–peachleaf willow, floodplain woodland
<b>Formation Class/Subclass: Herbaceous Vegetation/Perennial Graminoid Vegetation</b>						
Semipermanently flooded, temperate or subpolar grassland	<i>Carex pellita</i> seasonally flooded herbaceous vegetation	V.A.5.N.k.53	CEGL005272	<i>Carex</i> spp.–( <i>Carex pellita</i> , <i>Carex vulpinoidea</i> ) herbaceous vegetation	k.53	central Midwest, sedge meadow
Semipermanently flooded, temperate or subpolar grassland	<i>Eleocharis palustris</i> seasonally flooded herbaceous vegetation	V.A.5.N.k.61	CEGL001833	<i>Eleocharis palustris</i> herbaceous vegetation	k.61	creeping spikerush, wet meadow
Semipermanently flooded, temperate or subpolar grassland	<i>Pascopyrum smithii</i> intermittently flooded herbaceous vegetation	V.A.5.N.i.1	CEGL002038	<i>Pascopyrum smithii</i> – <i>Buchloe dactyloides</i> –( <i>Phyla cuneifolia</i> , <i>Oenothera canescens</i> ) herbaceous vegetation	i.1	wheatgrass, playa grassland
Semipermanently flooded, temperate or subpolar grassland	<i>Phalaris arundinacea</i> seasonally flooded herbaceous vegetation	V.A.5.N.k.20	CEGL001474	<i>Phalaris arundinacea</i> western herbaceous vegetation	k.20	reed canarygrass, wet meadow
Semipermanently flooded, temperate or subpolar grassland	<i>Polygonum</i> spp.– <i>Echinochloa</i> spp. temporarily flooded herbaceous vegetation	V.A.5.N.j.12	CEGL002039	<i>Polygonum</i> spp.– <i>Echinochloa</i> spp.– <i>Distichlis spicata</i> playa lake, herbaceous vegetation	j.12	playa marsh

Table I-1.

Formation Name	Alliance Name	Alliance Code	Unique Identifier	Association Name	Association Code	Common Name
Semipermanently flooded, temperate or subpolar grassland	<i>Typha (angustifolia, latifolia)</i> – ( <i>Schoenoplectus</i> spp.) semipermanently flooded, herbaceous vegetation	V.A.5.N.1.9	CEGL002389	<i>Typha</i> spp. Great Plains, herbaceous vegetation	1.9	northern Great Plains, cattail marsh
Semipermanently flooded, temperate or subpolar grassland	<i>Typha</i> spp.– ( <i>Schoenoplectus</i> spp., <i>Juncus</i> spp.) seasonally flooded vegetation	V.A.5.N.k.33	CEGL002026	<i>Schoenoplectus tabernaemontani</i> – <i>Typha</i> spp.– ( <i>Sparganium</i> spp., <i>Juncus</i> spp.) herbaceous vegetation	k.33	bulrush–cattail–burreed, shallow marsh
Semipermanently flooded, temperate or subpolar grassland	<i>Potamogeton</i> spp.– <i>Ceratophyllum</i> spp.– <i>Elodea</i> spp. permanently flooded, herbaceous vegetation	V.C.2.N.a.14	CEGL002044	<i>Potamogeton</i> spp.– <i>Ceratophyllum demersum</i> Great Plains herbaceous, central Midwest vegetation	a.14	Great Plains, pondweed, submerged, aquatic wetland
Short-sod, temperate or subpolar grassland	<i>Bouteloua gracilis</i> herbaceous alliance	V.A.5.N.e.9	CEGL001756	<i>Bouteloua gracilis</i> – <i>Buchloe dactyloides</i> herbaceous vegetation	e.9	blue grama–buffalograss, short-grass prairie
Tall-sod, temperate grassland	<i>Andropogon gerardii</i> – ( <i>Sorghastrum nutans</i> ) herbaceous alliance	V.A.5.N.a.2	CEGL002025	<i>Andropogon gerardii</i> – <i>Sorghastrum nutans</i> – <i>Hesperostipa spartea</i> loess hills, herbaceous vegetation	a.2	central tall-grass, big bluestem, loess prairie
Medium- to tall-sod, temperate or subpolar grassland	<i>Schizachyrium scoparium</i> – <i>Bouteloua curtipendula</i> herbaceous alliance	V.A.5.N.c.20	CEGL002035	<i>Schizachyrium scoparium</i> – <i>Bouteloua curtipendula</i> – <i>Bouteloua hirsuta</i> – ( <i>Yucca glauca</i> ) herbaceous vegetation	c.20	loess hills, little bluestem, dry prairie

**Association (Scientific) Name: *Carex* spp. (*Carex pellita*, *Carex vulpinoidea*) Herbaceous Vegetation**

*Translated Scientific Name:* Sedge species (woolly sedge, fox sedge) herbaceous vegetation

*Common Name:* Central Midwest sedge meadow

*Unique Identifier:* CEGLO05272

*Classification Code:* V.A.5.N.k.53

*Association Classification Confidence Level:* Weak

*Association Summary:* This sedge, wet meadow type is found in the central midwestern United States. Stands occur on nearly level floodplains, often in bands surrounding channels, or in basins. Soils are poorly drained, silty and clay loams formed in alluvium. Stands are flooded for much of the growing season, but may dry out in late summer. The vegetation cover is quite dense and may be patchy. The structure is dominated by graminoids 20–60 inches (0.5–1.5 m) tall. Typical species include *Carex cristatella*, *Carex molesta*, *Carex pellita* (= *Carex lanuginosa*), *Carex stipata*, *Carex tribuloides*, and *Carex vulpinoidea* (a dominant in southeast Nebraska meadows). Other frequent emergent graminoids include *Eleocharis* spp., *Juncus interior*, *Juncus torreyi* and *Scirpus atrovirens*. *Leersia oryzoides* may be common where the stand borders a marsh. Forbs are common and may be conspicuous. Among the more common are *Apocynum cannabinum*, *Symphytotrichum lanceolatum* (= *Aster lanceolatus*), *Lycopus americanus*, *Lythrum alatum*, and *Verbena hastata*. *Phalaris arundinacea* may invade this community to the point of excluding many of the native species.

**Association (Scientific) Name: *Bouteloua gracilis*–*Buchloe dactyloides* Herbaceous Vegetation**

*Translated Scientific Name:* Blue grama–buffalograss herbaceous vegetation

*Common Name:* Blue grama–buffalograss, short-grass prairie

*Unique Identifier:* CEGLO01756

*Classification Code:* V.A.5.N.e.9

*Association Classification Confidence Level:* Moderate

*Association Summary:* This blue grama–buffalograss, short-grass prairie type is common across much of the central and southern Great Plains of the United States. Stands occur on flat to rolling uplands. The surface soil may be sandy loam, loam, silt loam, or loamy clay. The subsoil is often finer than the surface soil. This community is characterized by a moderate to dense sod of short grasses with scattered mid-grasses and forbs. The dominant species are *Bouteloua gracilis* and *Buchloe dactyloides*. The foliage of these species is 2.8–7.6 inches (7–19 cm) tall, while the flowering stalks of *Bouteloua gracilis* may reach 18 inches (45 cm). The mid-grasses are usually stunted by the arid conditions and often do not exceed 28 inches (0.7 m). Other short graminoids found in this community are *Bouteloua hirsuta*, *Carex duriuscula*, *Carex inops*, *Carex heliophila*, and *Carex filifolia* (in Nebraska). Several mid-grasses occur regularly, such as *Aristida purpurea*, *Bouteloua curtipendula*, *Pascopyrum smithii*,

*Schizachyrium scoparium*, *Elymus elymoides*, *Sporobolus cryptandrus*, *Hesperostipa comata* (= *Stipa comata*), and *Vulpia octoflora*. Forbs such as *Astragalus* spp., *Gaura coccinea*, *Machaeranthera pinnatifida* var. *pinnatifida*, *Opuntia polyacantha*, *Plantago patagonica*, *Psoraleidum tenuiflorum*, *Ratibida columnifera*, and *Sphaeralcea coccinea* are common throughout this community. Shrubs are very rare except in the southern part of this community's range where scattered individuals may occur. In Oklahoma, other characteristic species include *Ambrosia artemisiifolia*, *Aristida oligantha*, *Machaeranthera tanacetifolia*, *Melampodium leucanthum*, *Muhlenbergia torreyi*, *Sporobolus compositus*, *Sporobolus cryptandrus*, and *Zinnia grandiflora*. In Texas, associated species include *Prosopis glandulosa*, *Bouteloua curtipendula*, and *Sporobolus cryptandrus*.

**Association (Scientific) Name: *Andropogon gerardii*–*Sorghastrum nutans*–*Hesperostipa spartea* Loess Hills Herbaceous Vegetation**

*Translated Scientific Name:* Big bluestem–yellow Indiangrass–porcupine grass, loess hills, herbaceous vegetation

*Common Name:* Central tall-grass, big bluestem, loess prairie

*Unique Identifier:* CEGLO02025

*Classification Code:* V.A.5.N.a.2

*Association Classification Confidence Level:* Moderate

*Association Summary:* This big bluestem, tall-grass prairie type is found in the west-central, tall-grass prairie region of the United States, including the Loess Hills. Stands occur on moderately steep mid- to upper slopes of loess hills and along ridges. It is most common on southern and western aspects. The soil is well-drained, acidic to neutral, and shallow to deep loess (16–40 inches [40–100 cm]). The parent material is a deep loess or glacial till and other deeply weathered substrates. This community is virtually lacking in shrubs and trees. Woody vegetation that is present, such as *Amorpha canescens*, is usually less than 20 inches (0.5 m) tall. The dominant vegetation is tall grasses. Of the dominant species, *Andropogon gerardii*, *Sorghastrum nutans*, and *Hesperostipa spartea* (= *Stipa spartea*) typically exceed 40 inches (1 m). *Schizachyrium scoparium*, also very common, is shorter. In Missouri, some other species that are usually found in this community are *Echinacea pallida*, *Potentilla arguta*, *Silphium laciniatum*, and *Sporobolus compositus* var. *compositus*.

**Association (Scientific) Name: *Schizachyrium scoparium*–*Bouteloua curtipendula*–*Bouteloua hirsuta*–(*Yucca glauca*) Herbaceous Vegetation**

*Translated Scientific Name:* Little bluestem–sideoats grama–hairy grama–(soapweed yucca) herbaceous vegetation

*Common Name:* Loess hills, little bluestem, dry prairie

*Unique Identifier:* CEGLO02035

*Classification Code:* V.A.5.N.c.20

*Association Classification Confidence Level:* Moderate

**Association Summary:** This bluestem-grama grass, dry prairie type is found on the loess bluffs along the east side of Missouri River in the central midwestern United States. The soil is somewhat rapidly drained and very shallow (0–16 inches [0–40 cm]). The vegetative structure is comprised of a single layer of dominant grasses intermixed with forbs. Shrubs, especially *Yucca glauca*, are sometimes present. This community is a short- to midgrass prairie dominated by the bunchgrasses *Andropogon gerardii*, *Bouteloua curtipendula*, and *Schizachyrium scoparium*. *Bouteloua hirsuta* can be common. *Sporobolus cryptandrus*, *Dalea leporina*, *Dalea candida*, *Dalea enneandra*, *Astragalus lotiflorus*, and *Astragalus missouriensis* can also be common. Other herbaceous species include *Pulsatilla patens* ssp. *multifida* (= *Anemone patens*), *Symphyotrichum sericeum* (*Aster sericeus*), *Buchloe dactyloides*, *Bouteloua gracilis*, *Delphinium carolinianum*, *Gaura coccinea*, and *Pediomelum argophyllum*, and the lichens *Dermatocarpon lachneum* and *Psora decipiens*.

**Association (Scientific) Name: *Eleocharis palustris* Herbaceous Vegetation**

**Translated Scientific Name:** Marsh spikerush, herbaceous vegetation

**Common Name:** Creeping spikerush, wet meadow

**Unique Identifier:** C EGL001833

**Classification Code:** V.A.5.N.k.61

**Association Classification Confidence Level:** Strong

**Association Summary:** This spikerush, wet meadow community is found in the central Great Plains of the United States and Canada and in the western United States. Stands occur in small depressions in intermittent streambeds or depression ponds that flood early in the season and may dry out by summer. Stands comprise submersed and emergent rooted vegetation under 40 inches (1 m) tall that is dominated by *Eleocharis palustris*, often in nearly pure stands. Soils are generally fine-textured.

**Association (Scientific) Name: *Pascopyrum smithii*–*Buchloe dactyloides*–(*Phyla cuneifolia*, *Oenothera canescens*) Herbaceous Vegetation**

**Translated Scientific Name:** Western wheatgrass–buffalograss–(wedgleaf frogfruit, spotted evening-primrose) herbaceous vegetation

**Common Name:** Wheatgrass, playa grassland

**Unique Identifier:** C EGL002038

**Classification Code:** V.A.5.N.i.1

**Association Classification Confidence Level:** Moderate

**Association Summary:** This wheatgrass, playa grassland community represents the common vegetation type of playa lake basins (depressional wetlands) under rangeland conditions in the southern and central Great Plains of the United States. In the central plains soils are dense silts and clays, occasionally loess-derived, that flood in winter and dry out by early summer. Perennial herbaceous graminoids and forbs less than 40 inches (1 m) tall dominate the community,

with composition varying depending on water levels. In the central plains *Pascopyrum smithii* is most abundant, with *Agrostis hyemalis*, *Eleocharis palustris*, *Eleocharis macrostachya*, *Elymus virginicus*, and *Hordeum jubatum* locally abundant. *Buchloe dactyloides* can be abundant in grazed sites. Early season ephemeral annuals include *Alopecurus carolinianus*, *Elatine rubella*, *Myosurus minimus*, *Veronica peregrina* ssp. *xalapensis*, and, more westward, *Limosella aquatica* and *Plagiobothrys scouleri*. Perennial forbs including *Ambrosia grayi*, *Phyla cuneifolia*, *Oenothera canescens*, *Rorippa sinuata*, and *Vernonia fasciculata* are conspicuous in places. In the southern plains, species characteristic of the type include *Buchloe dactyloides*, *Distichlis spicata*, and *Panicum obtusum*.

**Association (Scientific) Name: *Potamogeton* spp.–*Ceratophyllum demersum* Great Plains Herbaceous Vegetation**

**Translated Scientific Name:** Pondweed species–coontail, Great Plains herbaceous vegetation

**Common Name:** Great Plains pondweed, submerged aquatic wetland

**Unique Identifier:** C EGL002044

**Classification Code:** V.C.2.N.a.14

**Association Classification Confidence Level:** Weak

**Association Summary:** This community type is found in the Great Plains of the United States in shallow to relatively deep (40 inches [1 m]) freshwater basins or bands in marshes or bays that remain flooded in all but the driest years. Vegetation varies from sparse to dense, with submersed rooted and free-floating macrophytes. Species composition varies with substrate, water depth, water chemistry, turbidity, water temperatures, and other factors, but these are poorly understood. Dominant species in Nebraska include narrow-leaved pondweeds (*Potamogeton foliosus*, *Stuckenia pectinata* (= *Potamogeton pectinatus*), *Potamogeton pusillus*, *Najas guadalupensis*, and *Zannichellia palustris*. *Ceratophyllum demersum* and *Utricularia macrorhiza* can be locally abundant. In quiet bays, *Potamogeton nodosus* and *Lemna* spp. are common. In clear water with sandy bottoms *Chara* spp. may also be common.

**Association (Scientific) Name: *Phalaris arundinacea* Western Herbaceous Vegetation**

**Translated Scientific Name:** Reed canarygrass, western herbaceous vegetation

**Common Name:** Reed canarygrass, wet meadow

**Unique Identifier:** C EGL001474

**Classification Code:** V.A.5.N.k.20

**Association Classification Confidence Level:** Strong

**Association Summary:** This association is reported from throughout Washington, Colorado, Nebraska, Montana, Idaho, and into northeastern Utah, and is likely more widespread in the western United States. Its distribution as a natural type is complicated because this native species is widely cultivated as a forage crop

and has escaped and established in wetlands and riparian areas, displacing the local flora. Elevations range from near sea level to 5,576 feet (1,700 m). Stands are found along riparian areas, pond and lake margins, wet meadows, and intermittent drainages. Soils are commonly fine-textured and may be flooded for brief to extended periods. The vegetation is characterized by a dense, tall herbaceous layer (often greater than 80% canopy cover and 60–80 inches [1.5–2 m] tall) that is dominated by *Phalaris arundinacea*, which tends to occur in monocultures. Associated species may include *Equisetum arvense*, *Muhlenbergia asperifolia*, *Mentha arvensis*, *Schoenoplectus acutus* (= *Scirpus acutus*), and many other species in trace amounts where disturbed. Introduced species such as *Lepidium latifolium*, *Cirsium arvense*, *Sonchus oleraceus*, *Euphorbia esula*, and *Phleum pratense* are common in some stands.

**Association (Scientific) Name: *Polygonum* spp.–*Echinochloa* spp.–*Distichlis spicata* Playa Lake Herbaceous Vegetation**

*Translated Scientific Name:* Smartweed species–barnyard grass species–saltgrass playa lake herbaceous vegetation

*Common Name:* Playa marsh

*Unique Identifier:* CEGLO02039

*Classification Code:* V.A.5.N.j.12

*Association Classification Confidence Level:* Weak

*Association Summary:* This wetland community is found in the central Great Plains of the United States, where it occurs in shallow depressions on gently to moderately sloping topography. Soils are deep to moderately deep loams or clay loams underlain by a dense clay sublayer. Ponds often draw down periodically in these playa-type habitats. Annual herbaceous graminoids and forbs, mostly less than 40 inches (1 m) tall, dominate the exposed mud flats, and species composition and extent of the community fluctuate from site to site and year to year. In Nebraska, graminoids include *Cyperus acuminatus*, *Eleocharis engelmannii*, and *Echinochloa muricata*, and forbs include *Bacopa rotundifolia*, *Coreopsis tinctoria*, *Elatine rubella*, *Heteranthera limosa*, *Limosella aquatica*, *Lindernia dubia*, *Mollugo verticillata*, *Polygonum pennsylvanicum* (= *Polygonum bicorne*), *Polygonum lapathifolium*, *Rumex stenophyllus*, and *Sagittaria calycina*. In Kansas, graminoids include *Hordeum jubatum*, and forbs include *Ambrosia grayi*, *Symphotrichum subulatum* (= *Aster subulatus*), and *Chenopodium berlandieri*. The frequent water fluctuations and thick clay pan prevent establishment of most perennial hydrophytes, such as *Schoenoplectus* spp. (= *Scirpus* spp.) and *Typha* spp.

**Association (Scientific) Name: *Typha* spp. Great Plains Herbaceous Vegetation**

*Translated Scientific Name:* Cattail species, Great Plains herbaceous vegetation

*Common Name:* Northern Great Plains cattail marsh

*Unique Identifier:* CEGLO02389

*Classification Code:* V.A.5.N.1.9

*Association Classification Confidence Level:* Moderate

*Association Summary:* This cattail community type is found throughout the northern Great Plains of the United States and Canada. Stands occur in shallow (less than 20 inches [0.5 m]) or deep depressions, stock ponds, and seepy drainages. The vegetation is dominated by relatively pure stands of *Typha* spp., either *T. latifolia* or *T. angustifolia* or both. Many associates can occur including *Eleocharis* spp. and *Sagittaria latifolia*. This type may simply be a less diverse variation of *Typha* spp.–*Schoenoplectus* spp.–mixed herbs, Great Plains herbaceous vegetation (CEGL002228) that arises in disturbed wetland areas.

**Association (Scientific) Name: *Schoenoplectus tabernaemontani*–*Typha* spp.–(*Sparganium* spp., *Juncus* spp.) Herbaceous Vegetation**

*Translated Scientific Name:* Softstem bulrush–cattail species–(burreed species, rush species) herbaceous vegetation

*Common Name:* Bulrush-cattail-burreed shallow marsh

*Unique Identifier:* CEGLO02026

*Classification Code:* V.A.5.N.k.33

*Association Classification Confidence Level:* Weak

*Association Summary:* This shallow marsh, mixed emergent community ranges broadly over the midwestern United States and adjacent Canada. It is found in basin-like depressions, backwater areas of floodplains, and shallow margins of lakes or ponds. Soils are shallow to deep, very poorly drained, consisting of peats, mucks, or mineral materials, and often found in alluvium. Vegetation varies from zones dominated by tall emergents 40–80 inches (1–2 m) tall to those with hydrophytic annual and perennial forbs less than 40 inches (1 m) tall. In the tall emergent zone, *Schoenoplectus tabernaemontani* (= *Scirpus tabernaemontani*), *Schoenoplectus fluviatilis* (= *Scirpus fluviatilis*), *Schoenoplectus acutus* (= *Scirpus acutus*), *Typha angustifolia*, and *Typha latifolia* may dominate, mixed with a variety of other herbaceous species such as *Leersia oryzoides*, *Eleocharis palustris*, *Juncus* spp., and *Sparganium* spp. The hydrophytic annual and perennial forb zone is dominated by *Alisma subcordatum*, *Alisma plantago-aquatica*, *Sagittaria latifolia*, *Sparganium eurycarpum*, and *Pontederia cordata*, along with *Bacopa rotundifolia* and *Heteranthera limosa*. Occasional floating-leaved aquatics are sometimes present including *Azolla caroliniana*, *Lemna* spp., *Spirodela polyrrhiza*, and *Utricularia macrorrhiza*.

**Association (Scientific) Name: *Populus deltoides*–(*Salix amygdaloides*)/*Salix (exigua, interior)* Woodland**

*Translated Scientific Name:* Eastern cottonwood–(peachleaf willow)/(coyote willow, sandbar willow) woodland

*Common Name:* Cottonwood–peachleaf willow floodplain woodland

*Unique Identifier:* CEGLO00659

*Classification Code:* II.B.2.N.b.4

Association Classification Confidence Level: Moderate

Association Summary: This cottonwood-willow woodland is found widely in the central Great Plains of the United States. Stands occur on recently deposited alluvial material along rivers and streams. The soils are derived from alluvial sand, silt, and clay and are poorly developed. The water table fluctuates with the level of the adjacent river or stream. *Populus deltoides* is the dominant species in this community, although *Salix exigua* and/or *Salix interior* is generally more dominant in the initial stage following a major flood event. *Salix amygdaloides* is rare to codominant. The shrub/sapling layer is conspicuous, especially near the streambank, and consists mainly of *Salix exigua*, *Populus deltoides*, and *Salix amygdaloides*, or occasionally *Salix lutea*. In the more easterly parts of the range, *Salix interior* may replace *Salix exigua*. On the older margins of this community *Fraxinus pennsylvanica* is often found as a sapling or small canopy tree. The herbaceous stratum is variable. Graminoids typical of undisturbed sites include *Carex emoryi*, *Carex pellita* (= *Carex lanuginosa*), *Pascopyrum smithii*, and *Spartina pectinata*. *Equisetum arvense* and *Glycyrrhiza lepidota* are common forbs in these sites. Widely distributed species that are adapted to these sites include *Ambrosia psilostachya*, *Artemisia campestris* ssp. *caudata*, *Artemisia ludoviciana*, *Calamovilfa longifolia*, *Cenchrus longispinus*, *Chamaesyce serpyllifolia* (= *Euphorbia serpyllifolia*), *Euphorbia esula*, *Grindelia squarrosa*, *Helianthus petiolaris*, *Heterotheca villosa*, *Phyla lanceolata* (= *Lippia lanceolata*), *Opuntia macrorhiza*, *Poa pratensis*, and *Sporobolus cryptandrus*. These sites are prone to invasion by exotic grasses and forbs, the most widely established being *Agrostis stolonifera*, *Bromus tectorum*, *Cirsium arvense*, *Bassia scoparia* (= *Kochia scoparia*), *Melilotus* spp., *Taraxacum officinale*, and *Tragopogon dubius*.

**Association (Scientific) Name: *Salix exigua*/Mesic Graminoids Shrubland**

Translated Scientific Name: Coyote willow/mesic graminoids shrubland

Common Name: Sandbar willow/mesic graminoids shrubland

Unique Identifier: CEGLO01203

Classification Code: III.B.2.N.d.6

Association Classification Confidence Level: Strong

Association Summary: This riparian association is found primarily in the central Great Plains, but also occurs in parts of the Rocky Mountains and the Intermountain Region's semidesert areas. It generally occurs along backwater channels and other perennially wet, but less scoured sites such as floodplain swales and irrigation ditches. In Nebraska, this community is found on sandbars, islands, and shorelines of stream channels and braided rivers. The vegetation is characterized by the dominance of *Salix exigua* in a moderately dense tall-shrub canopy with a dense herbaceous layer dominated by graminoids. Other common shrubs include saplings of *Populus deltoides*

or *Salix amygdaloides*, *Salix eriocephala*, *Salix lutea*, and *Amorpha fruticosa*. Tall perennial grasses can appear to codominate the stand when *Spartina pectinata*, *Panicum virgatum*, or other tall grasses are present. Other mesic graminoids such as *Carex* spp., *Eleocharis* spp., *Juncus* spp., *Pascopyrum smithii*, *Schoenoplectus pungens* (= *Scirpus pungens*), and *Sphenopholis obtusata* may be present. Common forb species include *Bidens* spp., *Lobelia siphilitica*, *Lycopus americanus*, *Lythrum alatum*, *Polygonum* spp., and *Xanthium strumarium*. Diagnostic features of this association include the nearly pure stands of *Salix exigua* shrubs, with a dense herbaceous layer of at least 30% cover of mesic graminoids.

**FIELD VEGETATION ASSOCIATION  
PLANT COMMUNITY SUMMARY**

The NVCS clearly defines plant communities that can be discriminated on the landscape and at the association level is of fine enough detail to allow evaluation of management actions.

**NVCS ASSOCIATIONS IN THE RAINWATER BASIN  
WETLAND COMPLEX**

The following is a listing of the NVCS associations as they are used in the GIS data layer for plant communities that occur in the Rainwater Basin. The categories were divided into five representative classes: wet meadow, wetland plants, shrubs, trees, and uplands.

**Wet Meadow**

k.53 *Carex* spp.–(*C. pellita*, *C. vulpinoidea*) herbaceous vegetation (sedge species–[woolly sedge, fox sedge] herbaceous vegetation)

This sedge wet meadow type is found in the central midwestern United States. Stands occur on nearly level floodplains, often in bands surrounding channels, or in basins. Soils are poorly drained silty and clay loams formed in alluvium. Stands are flooded for much of the growing season, but may dry out in late summer. The vegetation cover is quite dense and may be patchy. The structure is dominated by graminoids 20–60 inches (0.5–1.5 m) tall. Typical species include *Carex cristatella*, *Carex molesta*, *Carex pellita* (= *Carex lanuginosa*), *Carex stipata*, *Carex tribuloides*, and *Carex vulpinoidea* (a dominant in southeast Nebraska meadows). Other frequent emergent graminoids include *Eleocharis* spp., *Juncus interior*, *Juncus torreyi* and *Scirpus atrovirens*. *Leersia oryzoides* may be common where the stand borders a marsh. Forbs are common and may be conspicuous. Among the more common are *Apocynum cannabinum*, *Symphytotrichum lanceolatum* (= *Aster lanceolatus*), *Lycopus americanus*, *Lythrum alatum*, and *Verbena hastata*. *Phalaris arundinacea* may invade this community to the point of excluding many of the native species.

**k.61 *Eleocharis palustris* herbaceous vegetation (creeping spikerush, wet meadow)**

This spikerush, wet meadow community is found in the central Great Plains of the United States and Canada and in the western United States. Stands occur in small depressions in intermittent streambeds or depression ponds that flood early in the season and may dry out by summer. Stands are composed of submersed and emergent rooted vegetation under 40 inches (1 m) tall that is dominated by *Eleocharis palustris*, often in nearly pure stands. Soils are generally fine-textured.

**i.1 *Pascopyrum smithii*–*Buchloe dactyloides*–(*Phyla cuneifolia*, *Oenothera canescens*) herbaceous vegetation, western wheatgrass–buffalograss–(wedgeleaf frogfruit, spotted evening-primrose) herbaceous vegetation**

This wheatgrass, playa grassland community represents the common vegetation type of playa lake basins (depressional wetlands) under rangeland conditions in the southern and central Great Plains of the United States. In the central plains, soils are dense silts and clays, occasionally loess-derived, that flood in winter and dry out by early summer. Perennial herbaceous graminoids and forbs less than 40 inches (1 m) tall dominate the community, with composition varying depending on water levels. In the central plains, *Pascopyrum smithii* is most abundant with *Agrostis hyemalis*, *Eleocharis palustris*, *Eleocharis macrostachya*, *Elymus virginicus*, and *Hordeum jubatum* locally abundant. *Buchloe dactyloides* can be abundant in grazed sites. Early season ephemeral annuals include *Alopecurus carolinianus*, *Elatine rubella*, *Myosurus minimus*, *Veronica peregrina* ssp. *xalapensis*, and more westward *Limosella aquatica* and *Plagiobothrys scouleri*. Perennial forbs including *Ambrosia grayi*, *Phyla cuneifolia*, *Oenothera canescens*, *Rorippa sinuata*, and *Vernonia fasciculata* are conspicuous in places. In the southern plains, species characteristic of the type include *Buchloe dactyloides*, *Distichlis spicata*, and *Panicum obtusum*.

### Wetland Plants

**k.20 *Phalaris arundinacea* western herbaceous vegetation (reed canarygrass, western herbaceous vegetation)**

This association is reported from throughout Washington, Colorado, Nebraska, Montana, Idaho, and into northeastern Utah, and is likely more widespread in the western United States. Its distribution as a natural type is complicated because this native species is widely cultivated as a forage crop and has escaped and established in wetlands and riparian areas, displacing the local flora. Elevations range from near sea level to 5,576 feet (1,700 m). Stands are found along riparian areas, pond and lake margins, wet meadows, and intermittent drainages. Soils are commonly fine-textured and may be flooded for brief to extended periods. The vegetation is characterized by a dense, tall herbaceous layer (often >80% canopy cover and 60–80 inches [1.5–2 m] tall) that is dominated by *Phalaris arundinacea*, which tends to occur in monocultures. Associated species may include

*Equisetum arvense*, *Muhlenbergia asperifolia*, *Mentha arvensis*, *Schoenoplectus acutus* (= *Scirpus acutus*), and many other species in trace amounts where disturbed. Introduced species such as *Lepidium latifolium*, *Cirsium arvense*, *Sonchus oleraceus*, *Euphorbia esula*, and *Phleum pratense* are common in some stands.

**j.12 *Polygonum* spp.–*Echinochloa* spp.–*Distichlis spicata* playa lake, herbaceous vegetation (smartweed species–barnyard grass species–saltgrass, playa lake, herbaceous vegetation)**

This wetland community is found in the central Great Plains of the United States, where it occurs in shallow depressions on gently to moderately sloping topography. Soils are deep to moderately deep loams or clay loams underlain by a dense clay sublayer. Ponds often draw down periodically in these playa-type habitats. Annual herbaceous graminoids and forbs, mostly less than 40 inches (1 m) tall, dominate the exposed mud flats, and species composition and extent of the community fluctuate from site to site and year to year. In Nebraska, graminoids include *Cyperus acuminatus*, *Eleocharis engelmannii*, and *Echinochloa muricata*. Forbs include *Bacopa rotundifolia*, *Coreopsis tinctoria*, *Elatine rubella*, *Heteranthera limosa*, *Limosella aquatica*, *Lindernia dubia*, *Mollugo verticillata*, *Polygonum pensylvanicum* (= *Polygonum bicorne*), *Polygonum lapathifolium*, *Rumex stenophyllus*, and *Sagittaria calycina*. In Kansas, graminoids include *Hordeum jubatum* and forbs include *Ambrosia grayi*, *Symphytotrichum subulatum* (= *Aster subulatus*), and *Chenopodium berlandieri*. The frequent water fluctuations and thick clay pan prevent establishment of most perennial hydrophytes, such as *Schoenoplectus* spp. (= *Scirpus* spp.) and *Typha* spp.

**i.9 *Typha* spp. Great Plains herbaceous vegetation (CEGL002389) (cattail species, Great Plains herbaceous vegetation)**

This cattail community type is found throughout the northern Great Plains of the United States and Canada. Stands occur in shallow (less than 20 inches [0.5 m]) or deep depressions, stock ponds, and seepy drainages. The vegetation is dominated by relatively pure stands of *Typha* spp., either *T. latifolia* or *T. angustifolia* or both. Many associates can occur, including *Eleocharis* spp. and *Sagittaria latifolia*. This type may simply be a less diverse variation of *Typha* spp.–*Schoenoplectus* spp.–mixed herbs, Great Plains herbaceous vegetation (CEGL002228) that arises in disturbed wetland areas.

**k.33 *Schoenoplectus tabernaemontani*–*Typha* spp.–(*Sparganium* spp., *Juncus* spp.) herbaceous vegetation (softstem bulrush–cattail species–[burreed species, rush species] herbaceous vegetation)**

This shallow marsh, mixed emergent community ranges broadly over the midwestern United States and adjacent Canada. It is found in basin-like depressions, backwater areas of floodplains, and shallow margins of lakes or ponds. Soils are shallow to deep, very poorly drained, consisting of peats, mucks, or mineral materials, and often found in alluvium.

Vegetation varies from zones dominated by tall emergents 40–80 inches (1–2 m) tall to those with hydrophytic annual and perennial forbs less than 40 inches (1 m) tall. In the tall emergent zone, *Schoenoplectus tabernaemontani* (= *Scirpus tabernaemontani*), *Schoenoplectus fluviatilis* (= *Scirpus fluviatilis*), *Schoenoplectus acutus* (= *Scirpus acutus*), *Typha angustifolia*, and *Typha latifolia* may dominate, mixed with a variety of other herbaceous species such as *Leersia oryzoides*, *Eleocharis palustris*, *Juncus* spp., and *Sparganium* spp. The hydrophytic annual and perennial forb zone is dominated by *Alisma subcordatum*, *Alisma plantago-aquatica*, *Sagittaria latifolia*, *Sparganium eurycarpum*, *Pontederia cordata*, along with *Bacopa rotundifolia* and *Heteranthera limosa*. Occasional floating-leaved aquatics are sometimes present, including *Azolla caroliniana*, *Lemna* spp., *Spirodela polyrrhiza*, and *Utricularia macrorrhiza*.

**a.14 *Ceratophyllum demersum* Great Plains herbaceous vegetation (Great Plains pondweed, submerged aquatic wetland)**

This community type is found in the Great Plains of the United States in shallow to relatively deep (40 inches [1 m]) freshwater basins or bands in marshes or bays that remain flooded in all but the driest years. Vegetation varies from sparse to dense, with submersed rooted and free-floating macrophytes. Species composition varies with substrate, water depth, water chemistry, turbidity, water temperatures, and other factors, but these are poorly understood. Dominant species in Nebraska include narrow-leaved pondweeds (*Potamogeton foliosus*, *Stuckenia pectinata* (= *Potamogeton pectinatus*), *Potamogeton pusillus*), *Najas guadalupensis*, and *Zannichellia palustris*. *Ceratophyllum demersum* and *Utricularia macrorrhiza* can be locally abundant. In quiet bays, *Potamogeton nodosus* and *Lemna* spp. are common. In clear water with sandy bottoms *Chara* spp. may also be common.

## Trees

**b.4 *Populus deltoides*–(*Salix amygdaloides*)/*Salix* (*exigua*, *interior*) woodland (eastern cottonwood–[peachleaf willow]/[coyote willow, sandbar willow] woodland)**

This cottonwood-willow woodland is found widely in the central Great Plains of the United States. Stands occur on recently deposited alluvial material along rivers and streams. The soils are derived from alluvial sand, silt, and clay and are poorly developed. The water table fluctuates with the level of the adjacent river or stream. *Populus deltoides* is the dominant species in this community, although *Salix exigua* and/or *Salix interior* is generally more dominant in the initial stage following a major flood event. *Salix amygdaloides* is rare to codominant. The shrub/sapling layer is conspicuous, especially near the streambank, and consists mainly of *Salix exigua*, *Populus deltoides*, and *Salix amygdaloides*, or occasionally *Salix lutea*. In the more easterly parts of the range, *Salix interior* may replace *Salix exigua*. On the older margins of this community

*Fraxinus pennsylvanica* is often found as a sapling or small canopy tree. The herbaceous stratum is variable. Graminoids typical of undisturbed sites include *Carex emoryi*, *Carex pellita* (= *Carex lanuginosa*), *Pascopyrum smithii*, and *Spartina pectinata*. *Equisetum arvense* and *Glycyrrhiza lepidota* are common forbs in these sites. Widely distributed species that are adapted to these sites include *Ambrosia psilostachya*, *Artemisia campestris* ssp. *caudata*, *Artemisia ludoviciana*, *Calamovilfa longifolia*, *Cenchrus longispinus*, *Chamaesyce serpyllifolia* (= *Euphorbia serpyllifolia*), *Euphorbia esula*, *Grindelia squarrosa*, *Helianthus petiolaris*, *Heterotheca villosa*, *Phyla lanceolata* (= *Lippia lanceolata*), *Opuntia macrorrhiza*, *Poa pratensis*, and *Sporobolus cryptandrus*. These sites are prone to invasion by exotic grasses and forbs, the most widely established being *Agrostis stolonifera*, *Bromus tectorum*, *Cirsium arvense*, *Bassia scoparia* (= *Kochia scoparia*), *Melilotus* spp., *Taraxacum officinale*, and *Tragopogon dubius*.

## Shrubs

**d.6 *Salix exigua*/mesic graminoids shrubland (coyote willow/mesic graminoids shrubland)**

This riparian association is found primarily in the central Great Plains, but also occurs in parts of the Rocky Mountains and Intermountain Region's semidesert areas. It generally occurs along backwater channels and other perennially wet, but less scoured, sites such as floodplain swales and irrigation ditches. In Nebraska, this community is found on sandbars, islands, and shorelines of stream channels and braided rivers. The vegetation is characterized by the dominance of *Salix exigua* in a moderately dense tall-shrub canopy with a dense herbaceous layer dominated by graminoids. Other common shrubs include saplings of *Populus deltoides* or *Salix amygdaloides*, *Salix eriocephala*, *Salix lutea*, and *Amorpha fruticosa*. Tall perennial grasses can appear to codominate the stand when *Spartina pectinata*, *Panicum virgatum* or other tall grasses are present. Other mesic graminoids, such as *Carex* spp., *Eleocharis* spp., *Juncus* spp., *Pascopyrum smithii*, *Schoenoplectus pungens* (= *Scirpus pungens*), and *Sphenopholis obtusata*, may be present. Common forb species include *Bidens* spp., *Lobelia siphilitica*, *Lycopus americanus*, *Lythrum alatum*, *Polygonum* spp., and *Xanthium strumarium*. Diagnostic features of this association include the nearly pure stands of *Salix exigua* shrubs, with a dense herbaceous layer of at least 30% cover of mesic graminoids.

## Uplands

**e.9 *Bouteloua gracilis*–*Buchloe dactyloides* herbaceous vegetation (blue grama–buffalograss)/purple three-awn, sideoats grama, sixweeks fescue**

This blue grama–buffalograss, short-grass prairie type is common across much of the central and southern Great Plains of the United States. Stands occur on flat to rolling uplands. The surface soil may be sandy

loam, loam, silt loam, or loamy clay. The subsoil is often finer than the surface soil. This community is characterized by a moderate to dense sod of short grasses with scattered mid-grasses and forbs. The dominant species are *Bouteloua gracilis* and *Buchloe dactyloides*. The foliage of these species is 2.8–7.6 inches (7–19 cm) tall, while the flowering stalks of *Bouteloua gracilis* may reach 18 inches (45 cm). The mid-grasses are usually stunted by the arid conditions and often do not exceed 28 inches (0.7 m). Other short graminoids found in this community are *Bouteloua hirsuta*, *Carex duriuscula*, *Carex inops* ssp. *heliophila*, and *Carex filifolia* (in Nebraska). Several mid-grasses occur regularly, such as *Aristida purpurea*, *Bouteloua curtipendula*, *Pascopyrum smithii*, *Schizachyrium scoparium*, *Elymus elymoides*, *Sporobolus cryptandrus*, *Hesperostipa comata* (= *Stipa comata*), and *Vulpia octoflora*. Forbs such as *Astragalus* spp., *Gaura coccinea*, *Machaeranthera pinnatifida* var. *pinnatifida*, *Opuntia polyacantha*, *Plantago patagonica*, *Psoralidium tenuiflorum*, *Ratibida columnifera*, and *Sphaeralcea coccinea* are common throughout this community. Shrubs are very rare except in the southern part of this community's range where scattered individuals may occur. In Oklahoma, other characteristic species include *Ambrosia artemisiifolia*, *Aristida oligantha*, *Machaeranthera tanacetifolia*, *Melampodium leucanthum*, *Muhlenbergia torreyi*, *Sporobolus compositus*, *Sporobolus cryptandrus*, and *Zinnia grandiflora*. In Texas, associated species include *Prosopis glandulosa*, *Bouteloua curtipendula*, and *Sporobolus cryptandrus*.

**a.2 *Andropogon gerardii*–*Sorghastrum nutans*–*Hesperostipa spartea* loess hills herbaceous vegetation (big bluestem–yellow Indiagrass)**

Stands occur on moderately steep mid- to upper slopes of loess hills and along ridges. It is most common on southern and western aspects. The soil is well-drained, acidic to neutral, and shallow to deep loess (16–40 inches [40–100 cm]). The parent material is a deep loess or glacial till and other deeply weathered substrates. This community is virtually lacking in shrubs and trees. Woody vegetation that is present, such as *Amorpha canescens*, is usually less than 20 inches (0.5 m) tall. The dominant vegetation is tall grasses. Of the dominant species, *Andropogon gerardii*, *Sorghastrum*

*nutans*, and *Hesperostipa spartea* (= *Stipa spartea*) typically exceed 40 inches (1 m). *Schizachyrium scoparium*, also very common, is shorter. In Missouri, some other species that are usually found in this community are *Echinacea pallida*, *Potentilla arguta*, *Silphium laciniatum*, and *Sporobolus compositus* var. *compositus*.

**c.20 *Schizachyrium scoparium*–*Bouteloua curtipendula*–*Bouteloua hirsuta*–(*Yucca glauca*) herbaceous vegetation (little bluestem–sideoats grama)–hairy grama–(soapweed yucca) herbaceous vegetation**

This bluestem-grama grass, dry-prairie type is found on the loess bluffs along the east side of Missouri River in the central midwestern United States. The soil is somewhat rapidly drained and very shallow (0–16 inches [0–40 cm]). The vegetative structure comprises a single layer of dominant grasses intermixed with forbs. Shrubs, especially *Yucca glauca*, are sometimes present. This community is a short- to midgrass prairie dominated by the bunchgrasses *Andropogon gerardii*, *Bouteloua curtipendula*, and *Schizachyrium scoparium*. *Bouteloua hirsuta* can be common. *Sporobolus cryptandrus*, *Dalea leporina*, *Dalea candida*, *Dalea enneandra*, *Astragalus lotiflorus*, and *Astragalus missouriensis* can also be common. Other herbaceous species include *Pulsatilla patens* ssp. *multifida* (= *Anemone patens*), *Symphotrichum sericeum* (*Aster sericeus*), *Buchloe dactyloides*, *Bouteloua gracilis*, *Delphinium carolinianum*, *Gaura coccinea*, and *Pediomelum argophyllum*, and the lichens *Dermatocarpon lachneum* and *Psora decipiens*.

## VEGETATION TRANSECT NOMENCLATURE

The final method of delineating plant communities was to use common name plant associations found within the basin. These communities were grouped into 11 categories. These categories then have representative plant communities commonly found in the basin.

Field codes that provide land managers with common names that they are familiar with were added to this hierarchy, allowing easy data recording during transect data collection. Table I-2 shows all of the local plant associations in the basin.

**Table I-2. Rainwater Basin vegetation mapping polygon descriptions.**

<i>Field Code</i>	<i>Examples or Includes</i>	<i>NVCS Code</i>
annual weed	sunflower, foxtail	weeds
building	McMurtrey, Cottonwood	infrastructure
Canada thistle	dominated by Canada thistle	noxious weed
cattail	native and hybrid	1.9
cedar tree	planted and volunteer cedars with multiple trees averaging 6 feet or taller and <32.5 feet (10 m) basal proximity	introduced
cropland	farmed, row crops	crop
introduced forb	perennial forbs, alfalfa, sweetclover, vetch	introduced
invasive cool-season plant	downy brome, Japanese brome, Kentucky bluegrass, smooth brome, intermediate	introduced
leafy spurge	mapped all patches regardless of size	noxious weed
moist-soil plant	annual early successional plants	j.12
musk thistle	dense patches that can be mapped	noxious weed
native grassland	untilled remnant prairie with mostly native, warm- or cool-season species	a.2, c.20, e.9
newly seeded vegetation	<2 years old with annual weeds or wild rye	a.2, c.20, e.10
parking lot	—	infrastructure
reed ( <i>Phragmites</i> spp.)	all mapped patches of reed ( <i>Phragmites</i> spp.)	noxious weed
planted, high-diversity, seeded vegetation	seeded native grass with 70+ native species included in mix	a.2, c.20, e.9
planted, native, cool-season vegetation	dike mix, other small patches	a.2, c.20, e.9
planted, native, warm-season vegetation	five species, old plantings with low diversity	a.2, c.20, e.9
prairie dog town	active dog towns	e.9
reed canarygrass	—	k.20
road	McMurtrey, well roads	infrastructure

**Table I-2. Rainwater Basin vegetation mapping polygon descriptions.**

<i>Field Code</i>	<i>Examples or Includes</i>	<i>NVCS Code</i>
bulrush ( <i>Scirpus</i> spp.)	<i>Schoenoplectus fluviatilis</i> (three square)	k.33
shrub	American plum, chokecherry, dogwood, sumac	d.6, d.27
tree	cottonwood, green ash, willow	b.4
water or mud flat	dirt, water, mud, void of vegetation (submergent plant communities)	a.14, or other dominant plants
wet meadow	rushes, sedges, or western wheatgrass	k.53, k.61, i.1



# Appendix L

## *Fire Management Program*

The U.S. Fish and Wildlife Service has management and administrative responsibility, including fire management, for approximately 24,000 acres of waterfowl production areas in the Rainwater Basin Wetland Management District.

### **FIRE—A CRITICAL NATURAL PROCESS**

In prairie ecosystems of the Great Plains, vegetation has evolved under periodic disturbance and defoliation from herbivores and fire, with minor weather events such as drought. This periodic disturbance is what kept the ecosystem diverse and healthy while maintaining significant biodiversity for thousands of years.

Historically natural fire, which includes Native American ignitions, has played an important disturbance role in many ecosystems: (1) removal of fuel accumulations; (2) decrease in undesirable plant communities; and (3) reduction in encroachment potential including trees, stimulating regeneration, cycling critical nutrients, and providing a diversity of habitats for wildlife. Higgins (1984) pointed out that 73% of historical lightning fires were started between July and August; the remaining 27% were started between April and June. Native American-set fires occurred mostly from February to June (most in April) and from July to November (most in October) (Higgins 1986).

When fire is excluded on a broad scale (over several decades) as it has been in many areas, the unnatural accumulation of living and dead fuel can contribute to degraded plant communities and wildlife habitats. These fuel accumulations often change the fire regime characteristics. Fuel accumulations have created the potential for uncharacteristically severe wildland fires in many areas across the country. These catastrophic wildland fires often pose risks to the safety of the public and firefighters. In addition, wildland fires threaten property and resource values such as wildlife habitat, grazing opportunities, timber, soils, water quality, and cultural resources.

Return of fire is essential for healthy vegetation and wildlife habitat in grassland and wetland ecosystems. When integrated back into an ecosystem, fire can help restore and maintain healthy systems. To facilitate fire's natural role in the environment, fire must be integrated into land and resource management plans on a broad scale.

Fire can do the following:

- improve waterfowl habitat through reduction of plant density, removal of organic material, and maintenance of early successional vegetation
- promote sediment removal in wetlands by wind scouring
- sustain biological diversity
- improve soil fertility
- improve the quality and amount of livestock forage
- reduce invasive plant communities including nonnative trees
- reduce the susceptibility of plants (caused by moisture and nutrient stress) to insects and disease
- improve water yield from off-site areas

### **WILDLAND FIRE MANAGEMENT POLICY AND GUIDANCE**

In 2001, the Secretaries of Interior and Agriculture completed and approved an update of the 1995 Federal Fire Policy. The 2001 Federal Wildland Fire Management Policy directs federal agencies to achieve a balance between (1) fire suppression to protect life, property, and resources and (2) fire use to regulate fuels and maintain healthy ecosystems. In addition, it directs agencies to use the appropriate management response for all wildland fires regardless of the ignition source. This policy provides eight guiding principles that are fundamental to the success of the fire management program:

- Firefighter and public safety is the first priority in every fire management activity.
- The role of wildland fires as an essential ecological process and natural change agent will be incorporated into the planning process.
- Fire management plans (FMPs), programs, and activities support land and resource management plans and their implementation.
- Sound risk management is a foundation for all fire management activities.
- Fire management programs and activities are economically viable, based on values to be protected, costs, and land and resource management objectives.

- FMPs and activities are based on the best available science.
- FMPs and activities incorporate consideration of public health and environmental quality. Federal, state, tribal, local, interagency, and international coordination and cooperation are essential.
- Standardization of policies and procedures among federal agencies is an ongoing objective.

Fire management considerations, guidance, and direction should be addressed in land use, resource management plans (such as a CCP). FMPs are step-down processes from the land use plans and habitat plans, with more detail on fire suppression, fire use, and fire management activities.

## MANAGEMENT DIRECTION

The Rainwater Basin Wetland Management District will protect life, property, and other resources from wildland fire by safely suppressing all wildland fires. Fire is an important management tool that can be used to accomplish habitat management objectives. If not used properly, fire can also quickly damage or destroy natural resources, equipment, buildings, and property and hurt or kill those that work with it. Prescribed fire and manual or mechanical fuels treatments will be used to reduce hazardous fuels and on district lands to reduce the intensity and severity of wildland fires. Special attention will be given to wildland–urban interface areas, both on Service-owned and adjacent lands, to reduce the risk of wildland fires to communities and improvements.

Prescribed fire and manual or mechanical fuel treatments will be used in an ecosystem management context for habitat management and to protect federal and private property. Fuel reduction activities will be applied where needed, especially in areas with a higher proportion of residences that may be considered “wildland–urban interface” areas. The prescribed fire program is outlined in the Rainwater Basin Fire Monitoring Plan.

All aspects of the fire management program will be conducted in a manner consistent with applicable laws, policies, and regulations. The district will maintain an FMP and carry out the plan to accomplish resource management objectives. Prescribed fire and fuel treatments will be applied in a scientific way under selected weather and environmental conditions to restore and maintain desired habitat conditions and control nonnative vegetation and the spread of woody vegetation. Up to approximately 6,500 acres of grasslands and wetlands will be treated annually to help accomplish habitat management objectives.

### **FIRE MANAGEMENT GOAL**

Use fire as an ecosystem process within wetland and grassland habitats and reach the needed level of prescribed fire management at the WPAs to control

invasive plants, encourage desirable native plants, and maintain productive wetlands that can benefit migratory birds.

### **Fire Management Objective A**

Through the duration of this CCP, increase the use of prescribed fire to 5,000–7,000 acres. Its use will shift from being a management tool to control woody and other undesirable invasive plants to being a management tool to maintain healthy grassland and wetland habitats.

### **Fire Management Objective B**

Through the duration of this CCP, increase the public’s awareness and support of fire as a management tool.

### **Fire Management Objective C**

Through the duration of this CCP, increase the number of fire-qualified partners and interagency prescribed burning.

### **Fire Management Strategies**

- Use strategies and tactics that consider public and firefighter safety as well as resource values at risk.
- Apply fire at a rate and intensity that takes the district from a restoration need (every 1–3 years) to a historical level of fire frequency (every 5–7 years).
- Apply fire in a mosaic pattern that leaves portions of treated WPAs unburned.
- Allow fire to travel through select shelterbelts to reduce cool-season grasses and litter.
- Work with district partners to provide demonstrations, written information, and other methods of communication to inform the public of the benefits of prescribed fire.
- Work with professional instructors and local colleges and universities to conduct classes in basic firefighter training.
- Work with the appropriate agencies to develop interagency agreements that will allow mutual assistance on prescribed burns.
- Develop detailed prescribed burn plans that describe the following:
  - burn units and their predominant vegetation;
  - the primary objectives for the units and specific objectives of the fire
  - acceptable range of results
  - site preparation requirements
  - weather requirements
  - safety considerations and measures to protect sensitive features
  - burn-day activities
  - communications and coordination for burns
  - ignition techniques

- smoke management procedure
- postburn monitoring

Methods (manual or mechanical means, timing, and monitoring) for wildland fire suppression, wildland fire use, and prescribed fire can be found in a more detailed list in the district's step-down FMP.

### Fire Management Rationale

Fire frequency in south-central Nebraska has been estimated to have occurred once every 5–7 years. Madden et al. (1999) assessed the effects of fire on grassland. They found that maximal grassland bird diversity is best attained by creating a mesic, mixed-grass prairie with areas of varying fire return intervals. Grassland birds such as Baird's sparrow, bobolink, grasshopper sparrow, and western meadowlark responded to burned areas. Other species such as common yellowthroat and clay-colored sparrow preferred prairie unburned for 8–10 years. The settlement of the Rainwater Basin has suppressed fire across the landscape.

Current fire at the WPAs is not frequent enough to control invading trees and shrubs. Past burns at the WPAs have shown that it takes three to four consecutive spring burns to remove woody plant invasion. This level of application is needed at approximately 20 WPAs, covering over 3,000 acres.

One of the problems that keep the district from reaching a greater fire frequency is the limited personnel available. A burn that is close to heavy fuel such as a shelterbelt requires a large fire crew to conduct the burn. Removal of shelterbelts will not only benefit grassland wildlife but will (1) eliminate

the need to establish fire lines, (2) reduce the needed size of a fire crew, and (3) reduce hazardous fuels.

Prescribed fire temporarily reduces air quality by reducing visibility and releasing several components through combustion. The four major components are carbon monoxide, carbon dioxide, hydrocarbons, and particulates. Varying amounts of particulate content are generated in different types of fuels, for example, wildlife habitat improvement burns versus fuel reduction burns. The district will meet the Clean Air Act emission standards by adhering to the requirements of the Nebraska State Implementation Plan during all prescribed fire activities.

### FIRE MANAGEMENT ORGANIZATION, CONTACTS, AND COOPERATION

Region 6 has established qualified, technical oversight and support for fire management using the "fire management district" approach. Under this approach, an established modeling system (based on the fire management workload of a group of refuges and possibly even that of interagency partners) has determined an appropriate fire management staffing organization. The fire management workload consists of (1) historical wildland-fire suppression activities and (2) historical and planned fuels treatment workload. Depending on funds, fire management staff and support equipment may be located at the district or at other units in the district and shared between all units.

Wherever possible, fire management activities will be conducted in a coordinated and collaborative manner with federal and nonfederal partners.



# Appendix M

## *Rainwater Basin Joint Venture Private Lands Program: Best Management Practices Eligibility Criteria and Cost Share*

### **Rainwater Basin Joint Venture Private Lands Program Best Management Practices Eligibility Criteria and Cost Share**



**Compiled by the Rainwater Basin Joint Venture  
Private Lands Workgroup**

**Approved by the Rainwater Basin Joint Venture Management Board  
March 14, 2006**



The Rainwater Basin Joint Venture Private Lands Workgroup developed the following programs and criteria to assist in the delivery and evaluation of Joint Venture (JV) programs on private lands. The proposed programs and criteria were developed with the JV objectives in mind, and will help ensure consistent and effective wetland habitat benefits through JV offerings. The Private Lands Work Group also recognizes that many wetland projects will not fit all of the criteria for a certain program, and will evaluate those projects with unique situations and information on a case-by-case basis to determine the program eligibility, applicable practices, and landowner benefits.

### **(1) Wetland Stewardship Program**

This program targets non-cropped wetland areas *only* and provides incentives to landowner's that maintain and manage these habitats.

#### **Wetland Criteria:**

- 1) Wetland is functioning properly. No hydrologic restoration is needed.
- 2) Vegetative community is mostly native species with less than 5% of the wetland dominated by invasives (e.g. trees, reed canary grass, noxious weeds).
- 3) Wetland has not been cropped since 1985.
- 4) Eligible wetland area will be determined by state and federal biologists using tools such as, but not limited to, historic photos, hydric soil maps, topography information, presence of hydric vegetation, etc.
- 5) Areas determined to be cropland *are not* eligible for stewardship payment.

#### **Practices:**

- ✓ Landowner will maintain wetland hydrology, no hydrologic alteration is allowed.
- ✓ Landowner will continue to manage and maintain the native plant community.
- ✓ Landowner will manage and control invasive species.

#### **Benefits:**

- 1) A \$25/acre/year stewardship payment for the hydric footprint (wetland area) over the 10-year agreement period with a \$500 minimum/year.
- 2) Landowner continues to manage the land.
- 3) Landowner controls hunting access.
- 4) This program includes technical assistance from JV partners during the agreement period.
- 5) No minimum wetland size.

### **(2) Restoration and Management Program**

This is a 10-year program that focuses on *restoring and managing* quality wetland habitat.

#### **Wetland Criteria:**

- 1) Landowner must be willing to restore hydrology to the maximum extent feasible within the basin and the vegetative community on non-cropped areas via pit filling, ditch plugs, water

control structures, fill or sediment removal, tree removal, installation of a variable flow tail water recovery system, etc.

- 2) Landowner enters a 10-year agreement that allows the Joint Venture Partners to manage the wetland for the duration of the agreement.

**Practices:**

- ✓ Landowner is given the first opportunity to perform actual management on the property.
- ✓ Reimbursement Rates:
  - Haying/Shredding: \$10/acre
  - Grazing Incentives: \$10/acre/year

Grazing must be concentrated in the wetland with preferred grazing during the growing season. Grazing payments only made when JV asks cooperator to graze wetland with specific timing/rate/goals.

These grazing cost-share incentives may be possible:

  - i. Fencing
    1. Standard electric
    2. High tensile electric
    3. Permanent barbed wire
  - ii. Water source for cattle
  - iii. Mineral blocks/tubs
  - Prescribed Fire: \$15/acre or donation to local fire department.
  - Disking: \$15/acre/pass with minimum of 2 passes using a standard farm disk.
  - Heavy Disking: \$35/acre/pass with a minimum of 2 passes using a >30” heavy construction disk.
  - Chemical Applications: Negotiated based on chemical cost. Labor and equipment rates determined by NRCS docket.
- ✓ Landowner has the right to defer management actions to a private contractor. The JV will help facilitate hiring the contractor to accomplish the objectives.

**Benefits:**

- 1) Landowner receives a “land use payment” for wetland restoration and vegetation management according to the following table:

Current Landuse	Fm, Fo	Sc, Sd	Ma, M,	Buffer
Cropped	\$60/A	\$50/A	\$50/A	\$60/A
Non-cropped	\$50/A	\$50/A	\$50/A	\$50/A

Payment is in exchange for the right of JV partners to restore the wetland and direct management on the project area for the 10-year agreement period. Land use payments are made annually and are not contingent on the amount of management during that year. They are contingent on the landowner’s willingness to participate. Wetland area may be over 25 acres in size; however land use payment will be capped at \$1,250/year (based on available funding). Management cost share will be available for the entire wetland area.

- 2) Landowner is allowed to continue farming if the area has previously been cropped.  
*Exception:* grasslands, pasture, or native prairie may not be brought into production.
- 3) Landowner may perform management on the project area without requesting permission from the JV. Grazing wetlands is encouraged as a way to supplement farm income.
- 4) Landowner controls hunting access. Access payments may be available from other partners.
- 5) Includes technical assistance from JV partners during the agreement period.
- 6) Center pivots are permitted as long as wheel crossing(s) do not interfere with wetland hydrology.

**(3) Hydrology Restoration Program (HRP-Pilot)**

The purpose of this program is to restore wetland hydrology to the fullest extent possible targeting temporary and seasonal wetlands. To participate, the landowner would agree to restore wetland hydrology and sign a 10-year agreement in exchange for an annual land use payment. Due to funding limitations, two focus areas have been identified for this *pilot program*. Maps of the focus areas have been made available to staff working directly with private landowners in these areas.

**Wetland Criteria:**

- 1) Only high priority wetlands (as determined by the RWB-JV Wetland Prioritization Model) are eligible. If all landowners owning the highest priority wetlands (red) are enrolled and funding still remains, landowners owning wetlands in the second highest priority level (salmon) will be contacted.
- 2) During the pilot period, basins that are 20 acres or smaller will be targeted.
- 3) Wetlands with severely altered hydrology (e.g. drains, pits) will be targeted first. These wetlands should have either no, or a low functionality.
- 4) Hydrology within the basin must be restored.

**Practices:**

- ✓ JV partners will provide economic and technical assistance necessary to restore the wetland.

**Benefits:**

- 1) Annual payments begin once restoration is complete. Payment rates follow the HRP table below.
- 2) JV partners provide restoration cost share.
- 3) No use restrictions for the project area.
- 4) Landowner controls hunting access.

**HRP-Pilot Payment Rates**

<b>Current Land Use</b>	<b>Fm, Fo</b>	<b>Sc, Sd</b>	<b>Ma, M,</b>	<b>Buffer</b>
Irrigated cropped	\$128/A	\$96/A	\$64/A	\$128/A
Dryland cropped	\$60/A	\$50/A	\$50/A	\$60/A
Pasture managed	\$15/A	\$15/A	\$15/A	\$15/A

#### **(4) Seasonal Habitat Improvement Program (SHIP)**

This program targets cropped wetland and allows the landowner to maintain cropping during the growing season. SHIP takes advantage of an opportunity to provide wetland habitat during times that the producer is not growing a crop. Ponding water on hydric soils during the winter months has little effect on the crop production of these sites. On hydric soils most crop loss occurs during May through July primarily due to saturation after rainfall events even if wetland drainage has occurred. This program provides a financial incentive to landowners who provide at least the opportunity to pond water on cropped areas during the off season.

##### **Wetland Criteria:**

- 1) Wetland or area to be flooded must be drained, but be capable of holding water utilizing a control structure.
- 2) On cropland, the landowner must allow natural runoff to pond in the area immediately following harvest. Water on the cropped area may be released after March 31 of each year. On pastureland, the landowner must allow natural runoff to pond on the area beginning November 1. Water may then be released after April 30 of each year. Instances where a SHIP occurs on both cropland and pasture, the water control structure can be adjusted on March 31 so that water is not held on the cropland acres.
- 3) The landowner understands that during the duration of this 10-year agreement, there is an inherent risk associated with late rains after March 31 that may prevent him from planting during some years.

##### **Practices:**

- ✓ Water control structure installation and associated dirt work.
- ✓ Cropland areas should have boards placed in control structure after harvest and remain in place at least until March 31.
- ✓ Pastureland should have boards placed in control structure on November 1 and remain in place at least until April 30.

##### **Benefits:**

- 1) The landowner, in consultation with the JV Partners, will determine an acceptable pool elevation based on a topographic survey.
- 2) Annual payment of \$50/acre/year for cropland area determined by landowner. This establishes an agreed upon pool elevation. Annual payment of \$25/acre/year for grassland and pastureland.
- 3) Includes technical assistance and restoration cost share provided by JV partners during the agreement period.
- 4) Center pivots are permitted.
- 5) Landowner controls hunting access.

#### **(5) CRP 23A Incentive Bonus Program - Conservation Reserve Enhancement Program**

This is a one-time signing bonus to promote enrollment and full restoration of wetlands under CP-23A. This program differs from the Hydrology Restoration Program in that landowners lose the right to crop the property for the duration of the CREP agreement.

**Wetland Criteria:**

- 1) Must sign up for CRP 23A (CREP would be included).
- 2) Cropland areas are eligible for incentive if the hydrology is restored (full hydrology restoration on applicant's property).
- 3) No prioritization needed in order to get as many wetlands restored as possible utilizing USDA funding.
- 4) Bonus payment paid on March 31 following completion of restoration.
- 5) Existing wetlands in the RWB that are currently in CRP are eligible for a bonus payment if the wetland hydrology is restored. Payment is based on # years remaining/# years in contract x \$500/acre (e.g. 8 yrs remain on a 10 yr contract =  $0.8 \times \$500 = \$400/\text{acre}$ ).

**Practices:**

- ✓ JV partners, in conjunction with the USDA, will provide economic and technical assistance necessary to restore the wetland.
- ✓ Eligible practices can be found in the CRP contract.

**Benefits:**

- 1) One-time payment of up to \$500/acre. All acres enrolled in the CRP CP23a contract are eligible for the payment. This includes uplands and wetlands. Bonus payment will not be made until restoration is complete. The bonus payment will be made on March 31 following restoration.
- 2) Cost share is available to assist with restoration costs not payable by the USDA.
- 3) Pivot can cross the CRP unless otherwise indicated by the USDA.
- 4) Landowner controls hunting access.

**(6) Short Term Vegetation Management Program**

This program differs from the Restoration Management Program in that it does not require wetland restoration. The Short Term Vegetation Management Program does not provide an annual land use payment, or any type of land access fee. The length of the agreement is negotiable.

**Wetland Criteria:**

- 1) Trees, cattail, reed canary grass, or river bulrush must be a component of the plant community.
- 2) Landowner allows access to the property for management treatment.
- 3) Landowner signs a negotiated-length agreement that allows the Joint Venture Partners access for treatment.
- 4) Financial assistance is limited to reimbursement for treatments only (no incentive payments).
- 5) Grazing must be concentrated in the wetland with preferred grazing during the growing season.

**Practices:**

- ✓ Landowner agrees to allow any or all of the following: disking, chemical application, haying/shredding, prescribed burning, grazing, silt removal, pit filling, and tree removal activities.

Reimbursement Rates:

- Haying/Shredding: \$10/acre
- Intensive Wetland Grazing: \$10/acre/year
- Prescribed Burning: \$15/acre
- Disking: \$15/acre/pass with minimum of 2 passes using a standard farm disk.
- Heavy Disking: \$35/acre/pass with a minimum of 2 passes using >30" heavy construction disk.
- Chemical Applications: Negotiated based on specific chemical cost; labor and equipment rates determined by NRCS docket.

**Benefits:**

- 1) Will add functional value to the wetland.
- 2) Additional grazing incentives may be possible such as fencing, installation of a water source, or purchase of mineral block(s).

**(7) General Joint Venture Cost Share Activities**

The Joint Venture partners accept certain practices on their own, or in combination with the programs outlined above. Cost share is available for these activities if they provide benefits to wildlife habitat with emphasis on migratory birds, or improve watershed hydrology with direct benefits to a protected wetland. A 10-year agreement is required for cost share assistance.

**Wetland Improvements**

- 1) Sediment Control Structures
- 2) Fencing to encourage long term grazing in wetlands
- 3) Grazing (includes deferments)
- 4) Variable Flow Tail Water Recovery System
- 5) Close and Remove Roads
- 6) Vegetation Management
- 7) Junk Pile Removal
- 8) Water Development (wells, pivots, pipe, livestock)
- 9) Re-size/add/replace culverts
- 10) Water Control Structures
- 11) Wetland Creation
- 12) Wetland Seeding
- 13) Earthwork
  - ✓ Pit fills
  - ✓ Drain fills
  - ✓ Silt/sediment removal
  - ✓ Filling road/drainage ditches
  - ✓ Power line burial
  - ✓ Pipeline burial
  - ✓ Dikes/berms

- ✓ Tile drain removal

### **Watershed Improvements**

- 1) Pit fills
- 2) Resize/add/replace culverts
- 3) Buffers
- 4) Grassland Restoration
- 5) Sediment Control Structures
- 6) Variable Flow Tail Water Recovery System
- 7) Removing Flow Restrictions
- 8) Terrace Removal
- 9) Close and Remove Roads
- 10) Tile Drain Removal

## **(8) Easements**

An easement program remains under development until enough properties are secured that an actual program can be evaluated and implemented. JV easements will be prioritized for functioning wetlands without hydrologic modification that are not eligible for NRCS's Wetland Reserve Program. The goal of easement offerings, as with all other JV programs, is to meet wetland protection needs that cannot be met through existing programs. At this time we will follow this format:

### **A. Hydrologic Restoration/Protection**

Provide for on-site restoration of the wetland hydrology, prevent future hydrologic alterations and non-agricultural land use, and place no restrictions on agricultural land use.

### **B. Pasture/Protection Easement**

Provide for on-site restoration of the wetland hydrology, prevent development of any kind, prevent excavation or filling which impacts hydrology, limits land use to pasture, but does not dictate management.

### **C. Protection/Restoration/Management**

Provide for on-site restoration of the wetland hydrology, prevent any hydrologic alterations and non-agricultural land use, and place restrictions on land use and activities for the purpose of assuring maximum wetland habitat values.

The value of the easement will be determined through an appraisal of the value of those rights to be purchased. The minimum length of the easement terms will be 25 years and will be agreed to by the landowner and the entity holding the easement. The USFWS has expressed an interest in holding perpetual restoration easements (example A above). The easement must not restrict the landowner's agricultural rights to the land. In other words, if he is cropping it now he may still try to crop it once the wetland is restored. These easements would have to be perpetual in length.

# Appendix N

## *Wetland Management District Ditch and Tile Maintenance Policy*

---

This policy applies to existing constructed ditches or tiles that come onto waterfowl production areas (WPAs) where no reservation of a drainage easement exists in the WPA title or deed. If there is a drainage reservation in the deed, the Service will follow the terms of that reservation.

- No new wetland or upland drainage facility will be allowed within a WPA.
  - Existing drainage cannot be improved beyond the original construction.
    - Tile may not be replaced with a larger tile.
    - Ditches may not be cleaned out beyond their original depth, width, or length.
    - Ditches may not be replaced with tile lines except where the tile is installed at the same or higher elevation than the original ditch bottom or in other rare exceptions to solve severe erosion.
  - All materials cleaned out of a ditch will be removed from the WPA.
  - All construction sites at WPAs will be seeded down to a grass mix specified by the Service.
  - Cleanout activities will not be allowed during the waterfowl-breeding season (April 1–August 1).
  - If silt deposition is a concern, the Service will request that a grassed waterway or silt basin be installed upstream of Service property to help reduce future siltation.
- Cleanout of natural (never ditched) drainage ways will not be allowed.
  - Ditch and tile maintenance work at WPAs will only be done after the wetland district manager has approved the project and issued a special use permit. (Note: Compatibility determinations are not necessary since the Service does not control maintenance of the system; the Service only controls the timing and scope of maintenance.)
  - Landowners may still be subject to the Swampbuster Provisions and the U.S. Army Corps of Engineers' rules on maintenance and abandonment of ditches.
  - Mowing or spraying of approved herbicide in a ditch after August 1 may be permitted in lieu of excavation.
  - If the ditch has not been cleaned or a tile has not functioned for more than 25 years, or the watershed above the ditch has been substantially altered since the Service bought the property (significant increase in flows or degradation of water quality), a formal right-of-way request maybe required as determined by the wetland district manager.



# Appendix 0

## *RONs and SAMMS Projects*

### **Refuge Operating Needs System (RONs)**

<i>Project Number</i>	<i>Project Description</i>	<i>First-year Need (\$1000s)</i>	<i>Recurring Base Need (\$1000s)</i>	<i>Personnel (FTEs)</i>
97005	Control exotic species invasion.	132	30	0
97007	Improve water management on new waterfowl production areas.	341	30	0
00001	Establish water rights on refuge lands.	109	5	0
00004	Livestock confinements and intensive agriculture's effect on water quality.	162	10	0
97012	Increase public use of refuge lands.	140	75	1
00008	Private lands coordinator for Nebraska Sandhills.	169	104	1
99002	Archaeological review of refuge lands.	70	0	0
99005	Livestock fencing.	78	23	0
97002	Wetland restoration on refuge lands.	158	66	1
97010	Endangered species restoration.	140	75	1
99001	Law enforcement and property protection.	140	75	1
99004	Interpretive and recreational access to public lands.	99	10	0
00003	Snow goose impact on migratory bird populations within Rainwater Basin (NE).	258	10	0
97001	Habitat and population surveys.	70.5	38	0.5
97009	Wetland restoration on private lands.	161.5	90	1
00006	Expand prescribed fire program.	140	75	1
06006	Water delivery to create wetland habitat in CO, NE, ND, MT, and KS.	380	0	0

**Service Asset Maintenance Management system (SAMMS)**

<i>Project Number</i>	<i>Project Description</i>	<i>Cost (\$1000s)</i>
<b>Deferred Maintenance</b>		
00105716	Repair worn pump engine and gear head at Smith unit, Harvard WPA.	32
02120249	Replace water-pumping station at Clark WPA.	51
02120162	Replace boundary fences at Clay County WPAs.	42
00105719	Replace worn pump engine and gear head on well at Hultine WPA.	30
02120273	Replace water-pumping station at Spring WPA, south.	51
02120166	Replace fences at six WPAs in Fillmore County.	28
02120254	Repair water-pumping station at Cottonwood WPA.	37
02120271	Replace water-pumping station at Heron WPA.	51
02120169	Replace fences at three WPAs in Phelps County.	73
00105713	Replace worn pump engine and gear head on well at Prairie Dog WPA.	35
00105710	Replace worn pump engine and gear head on well at Massie WPA.	28
00105714	Repair worn pump engine and gear head on east well at McMurtrey WPA.	32
02120170	Replace fences at three WPAs in Franklin County.	57
02120199	Replace water-pumping station at Mallard Haven WPA.	26
00105715	Repair worn pump engine and gear head on well at Harvard WPA, north.	30
00105717	Replace worn pump engine and gear head on Knudson well at Harvard WPA.	38
02120201	Repair worn pump and engine on south well at Krause WPA.	29
02120200	Repair worn-out pump and engine on north pumping station at Krause WPA.	26
02120198	Replace worn water-pumping station at Hanson WPA.	47
02120257	Replace water-pumping station at Johnson WPA, east.	39
00105712	Replace worn pump engine and gear head on well at Eckhardt WPA.	30
00105718	Replace worn pump engine and gear head on well at Mallard Haven WPA.	17
00105711	Replace worn pump engine and gear head on north well at Youngson WPA.	22
02120206	Replace water-pumping station at Lindau WPA.	49
98105704	Repair rutted gravel road.	43
02120195	Repair water-pumping station at north well at Hultine WPA.	33
02120158	Repair earthen dike at Springer WPA.	37
98109671	Repair dike.	278
<b>Heavy Equipment</b>		
91105702	Replace worn front-end loader.	200
04133918	Replace Bobcat skid steer loader.	40
99105705	Replace worn 1981 dump truck.	51
01116415	Replace 1990 well maintenance truck.	51
01116225	Replace 1967 bulldozer.	195
01116255	Replace Clark 4x4 hinged tractor-dozers.	275

**Service Asset Maintenance Management system (SAMMS)**

<i>Project Number</i>	<i>Project Description</i>	<i>Cost (\$1000s)</i>
99105707	Replace worn equipment truck.	110
04133810	Replace 1966 Cuece 1/2 Kaiser jeep.	50
01116249	Replace WABCO earthmoving scraper.	264
01116227	Replace 1996 Caterpillar bulldozer.	195
01116250	Replace Westinghouse model earthmoving scraper.	264
01116411	Replace worn John Deere tractor.	90
05139000	Replace 2003 Freightliner semi-truck.	200
05139017	Replace 1979 flatbed dump truck.	60
<b>Small Equipment</b>		
01116271	Replace 1992 Dodge pickup.	30
01116409	Replace 1993 Chevrolet Blazer.	30
01116407	Replace worn Jeep.	31
01116273	Replace 1994 Ford 1/2-ton pickup.	29
01116219	Replace worn heavy-duty disc used for invasive plant control.	19
04133805	Replace 2001 F-150 Ford pickup.	31
04133817	Replace 2002 Chevrolet Silverado 1/2-ton pickup.	31
04133809	Replace 2001 F-450 Ford fire truck.	40
04133820	Replace 2002 F-250 Ford pickup.	37
04133811	Replace 2002 Chevrolet Suburban.	37
04133829	Replace 2002 F-450 Ford flatbed truck.	37
04133816	Replace 2002 Chevrolet Impala automobile.	22
04133830	Replace 2002 F-250 Ford pickup.	31
04133831	Replace 2004 F-350 Ford pickup.	37
99105693	Replace worn 12-foot farm disc.	22
99105696	Replace worn 4x4 ATV.	14
98105703	Replace worn backhoe utility tractor.	38
01116212	Replace 1991 Panther 16-foot airboat.	24
01116216	Replace crane on well repair truck.	14
01116241	Replace 16-inch Crisafulli pump.	15
01116244	Replace mobile 12-inch gator pump.	15
01116254	Replace John Deere tractor, model 6400.	58
01116256	Replace lowboy trailer.	18
01116259	Replace airboat trailer.	9
01116260	Replace airboat trailer.	8
01116262	Replace worn dump trailer.	31
01116266	Replace 1998 Ford 1-ton pickup.	31

<b>Service Asset Maintenance Management system (SAMMS)</b>		
<i>Project Number</i>	<i>Project Description</i>	<i>Cost (\$1000s)</i>
01116267	Replace 1999 Ford 1-ton fire engine.	37
01116406	Replace 1995 Ford pickup.	31
01116410	Replace 1998 Dodge 4x4 pickup.	31
01116412	Replace 2001 Dodge 4x4 pickup.	37
01116413	Replace 2000 Ford crew cab truck.	37
01116414	Replace John Deere tractor, model 5310.	42
04133800	Replace 1997 sport utility vehicle.	30
05138982	Replace 2001 Ford 4x4 fire truck.	36
05138987	Replace 2001 Ford 4x4 2-ton truck.	40
05138989	Replace John Deere tractor, model 5410.	65
05138990	Replace Hyster forklift, model H60XL.	60
05138991	Replace incinerator.	25
05138995	Replace American Eagle forklift, model AE8122.00.	100
05138997	Replace 2002 Chevrolet 2-ton truck.	40
05138999	Replace Caterpillar road grader.	220
05139001	Replace semi-truck trailer, lowboy model.	45
05139004	Replace 2003 mule ATV.	8
05139005	Replace 2004 Ford crew cab truck.	40
05139007	Replace 2004 Ford super crew 4x4 truck.	45
05139010	Replace 2005 Ford Hybrid Escape automobile.	30
05139012	Replace 2005 Ford cab/chassis.	28
05139014	Replace 2005 Ford super cab 4x4 truck.	35
05139016	Replace 1979 flatbed dump truck.	50
05139018	Replace 1998 Chevrolet 3/4-ton truck.	35
<b>Roads and Parking Lots (TEA 21)</b>		
03126571	PE/CN* (parking lots 9001–3, 9009–16, 9033, 9121–23, 9071–74).	306
03126572	PE/CN (parking lots 9054–61, 9120, 9077–82, 9084–90).	380
03126573	PE/CN (parking lots 9062–69, 9091–9102, 9116–19).	470
03126574	PE/CN (route 100, 0.3 mile; parking lots 9004–08, 9017–32, 9034–42, 9044–53).	595
<b>Small Construction</b>		
01116196	Replace aged storage shed damaged by storm.	90
04133864	Replace fence at Peterson WPA.	41
02120274	Replace water-pumping station at Rauscher WPA.	51
<b>Large Construction</b>		
00109809	Construct new office building [p/d/cc].	2400

\* PE/CN = preliminary engineering and construction.

# Bibliography

- [AOU] American Ornithological Union. 2004. American Ornithological Union's Check-list of North American Birds. 7th ed. [Place of publication unknown]. [Number of pages unknown].
- Anderson, J.T.; Smith, L.M. 1998. Protein and energy production in playas: implications for migratory bird management. *Wetlands* 18:437–46.
- . 1999. Carrying capacity and diel use of managed playa wetlands by nonbreeding waterbirds. *Wildlife Society Bulletin* 27:281–91.
- Anderson, J.T.; Smith, L.M. 2000. Invertebrate response to moist-soil management of playa wetlands. *Ecological Applications* 10(2):550–8.
- Ankney, C.D.; Afton, A.D. 1988. Bioenergetics of breeding northern shovelers: diet, nutrient reserves, clutch size, and incubation. *Condor* 90:459–72.
- Ankney, C.D.; Alisauskas, R.T. 1991. Nutrient-reserve dynamics and diet of breeding female gadwalls. *Condor* 93:799–810.
- Ankney, C.D.; MacInnes, C.C. 1978. Nutrient reserves and reproductive performance of female lesser snow geese. *Auk* 95:459–71.
- Austin, J.E.; Richert, A.L. 2001. A comprehensive review of observational and site evaluation data of migrant whooping cranes in the United States, 1943–99. U.S. Geological Survey Report. Jamestown, ND: U.S. Geological Survey, Northern Prairie Wildlife Research Center. 156 p.
- Bakker, K.K. 2000. Avian occurrence in woodlands and grasslands on public areas throughout eastern South Dakota [PhD dissertation]. Brookings, SD: South Dakota State University. [Number of pages unknown].
- . 2003. The effect of woody vegetation on grassland nesting birds: an annotated bibliography. In: *The Proceedings of the South Dakota Academy of Sciences*; [Place and date of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 82:119–41.
- Baldassarre, G.A.; Bolen, E.G. 2006. Waterfowl ecology and management. 2nd ed. [City of publication unknown], FL: Krieger Publishing Company. 567 p.
- Baldassarre, G.A.; Whyte, R.J.; Quinlan, E.E. [et al.] 1983. Dynamics and quality of waste corn available to postbreeding waterfowl in Texas. *Wildlife Society Bulletin* 11:25–31.
- Ballard, B.M.; Thompson, J.E.; Petrie, M.J. [et al.] 2004. Diet and nutrition of northern pintails wintering along the southern coast of Texas. *Journal of Wildlife Management* 68(2):371–82.
- Baxter, W.L.; Wolfe, C.W. 1973. Life history and ecology of the ring-necked pheasant in Nebraska. Lincoln, NE: Nebraska Game and Parks Commission. 58 p.
- BBC Research & Consulting. 2006. Rainwater Basin Wetland Management District socioeconomic impact analysis: preliminary report. Unpublished report on file at Rainwater Basin Wetland Management District, NE. 9 p.
- Benning, D.S. 1987. Coordinated mid-continent white-fronted goose survey. Washington DC: U.S. Department of the Interior, Fish and Wildlife Service. U.S. Fish and Wildlife Service annual report. 7 p.
- Bergan, J.F.; Smith, L.M. 1993. Survival rates of female mallards wintering in the Playa Lakes Region. *Journal of Wildlife Management* 57:570–7.
- Bishop, A.A. 2005. Rainwater Basin Joint Venture waterfowl habitat use model (Rainwater Basin Joint Venture GIS layer). Unpublished report on file at Rainwater Basin Wetland Management District, Grand Island, NE. 27 p.
- Bishop, A.A.; Reker, R. 2006. Landcover within the RWB: a GIS layer created with 2004 data. Unpublished reports on file at Rainwater Basin Wetland Management District, Grand Island, NE.
- Bowyer, M.W.; Stafford, J.D.; Yetter, A.P. [et al.] 2005. Moist-soil plant seed production for waterfowl at Chautauqua National Wildlife Refuge, Illinois. *The American Midland Naturalist*: 154(2):331–41.
- Brennan, E.K. 2006. Local and landscape level variables influencing migratory bird abundance, diversity, behavior, and community structure in Rainwater Basin wetlands [PhD dissertation]. [Place of publication unknown]: Texas Technological University. 180 p.
- Brown, S.; Hickey, C.; Harrington, B.; editors. 2000. United States Shorebird Conservation Plan. Manomet, MA: Manomet Center for Conservation Sciences. 60 p.
- Burger, L.D.; Burger, L.W., Jr.; Faaborg, J. 1994. Effects of prairie fragmentation predation on artificial nests. *Journal of Wildlife Management* 58(2):249–54.
- Butcher, Greg. [Date unknown]. Central mixed-grass prairie executive summary. *Partners in Flight*. <[http://www.blm.gov/wildlife/pl\\_34sum.htm](http://www.blm.gov/wildlife/pl_34sum.htm)> [Date of access unknown].
- Checkett, J.M.; Drobney, R.D.; Petrie, M.J. [et al.] 2002. True metabolizable energy of moist soil seeds. *Wildlife Society Bulletin* 30(4):1113–19.

- Clark, R.J. 1975. A field study of the short-eared owl, *Asio flammeus* (Pontoppidan), in North America. *Wildlife Monographs* 47:1–67.
- Clayton, K.M.; Schmutz, J.K. 1999. Is the decline of burrowing owls *Speotyto cunicularia* in prairie Canada linked to changes in Great Plains ecosystems? *Bird Conservation International* 9:163–85.
- Connelly, J.W.; Gratson, M.W.; Reese, K.P. 1998. Sharp-tailed grouse (*Tympanuchus phasianellus*). In: Poole, A.; Gill, F.; editors. *The Birds of North America*, No. 354 [doi:10.2173/bna.354]. Philadelphia: The Academy of Natural Sciences; Washington DC: The American Ornithologists' Union. [Pages unknown].
- Cox, R.R., Jr. 1998. Spring-staging ecology of greater white-fronted geese, lesser snow geese, and northern pintails in the central Platte River valley and Rainwater Basin of Nebraska. Northern Prairie Wildlife Research Center preliminary progress report. [Unpublished report]. 4 p.
- . 1999. Spring-staging ecology of waterfowl in Nebraska—then versus now. *Waterfowl* 2000 12(2):18. Jamestown, ND: Northern Prairie Wildlife Research Center Online. <<http://www.npwrc.usgs.gov/resource/birds/ss ecol/index.htm>> (Version 02DEC99) [Date of access unknown].
- Cox, R.R., Jr.; Davis, B.E. 2002. Habitat use, movements, and survival of female northern pintails during spring migration in Nebraska and subsequent potential breeding-site selection in the Prairie Pothole Region [progress report]. Bismarck, ND: U.S. Geological Survey, Northern Prairie Wildlife Research Center. 150 p.
- Cox, R.R., Jr.; Samuel, M.D.; Church, K.E. 1999. Body condition of adult lesser snow geese killed by avian cholera [progress report]. 20 p.
- Dai, X.; Boutton, T.W.; Hailemichael, M. [et al.] 2006. Soil carbon and nitrogen storage in response to fire in a temperate mixed-grass savanna. *Journal of Environmental Quality* 35:1620–8.
- Davis, S.K. 2004. Area sensitivity in grassland passerines: Effects of patch size, patch shape, and vegetation structure on bird abundance and occurrence in southern Saskatchewan. *The Auk* 121(4):1130–45.
- DeJong, J.R. 2001. Landscape fragmentation and grassland patch size effects on non-game grassland birds in xeric mixed-grass prairies of western South Dakota. Brookings, SD: South Dakota State University. [Pages unknown].
- Delnicki, D.; Reinecke, K.J. 1986. Mid-winter food use and body weights of mallards and wood ducks in Mississippi. *Journal of Wildlife Management* 50:43–51.
- Dewey S.L. 1986. Effects of the herbicide atrazine on aquatic insect community structure and emergence. *Ecology* 67:148–62.
- Dodson, S.I.; Merritt, C.M.; Shannahan, J.P. [et al.] 1999. Low exposure concentrations of atrazine increase male production in *Daphnia pulex*. *Environmental Toxicology and Chemistry* 18(7): 1568–73.
- Drahota, J. 2000. Results of ground survey 2000 [unpublished report]. On file at Rainwater Basin Wetland Management District. 7 p.
- . 2006. A case for managing Rainwater Basin wetlands [unpublished report]. On file at Rainwater Basin Wetland Management District. 11 p.
- Drahota, J.; Bishop, A.; Walters, R. 2004. Rainwater Basin wetland complex south-central Nebraska 2003–2004 vegetation mapping and monitoring project [unpublished report]. On file at Rainwater Basin Wetland Management District, Kearney, NE. 53 p.
- Dubovsky, J.A.; Kaminski, R.M. 1994. Potential reproductive consequences of winter-diet restriction in mallards. *Journal of Wildlife Management* 58(4): 780–6.
- Duebbert, H.F.; Lokemoen, J.T. 1977. Upland nesting of American bitterns, marsh hawks, and short-eared owls. *The Prairie Naturalist* 9(3–4):33–40.
- Dugger, B.D.; Dugger, K.M. 2002. Long-billed curlew (*Numenius americanus*). In: Poole, A.; Gill, F.; editors. *The Birds of North America*, No. 628 [doi:10.2173/bna.628]. Philadelphia, PA: The Academy of Natural Sciences; Washington DC: The American Ornithologists' Union. [Pages unknown].
- Ekstein, J.D.; Hygnstrom, S.E. 1996. Fate of wetlands associated with the Central Nebraska Irrigation Canal System. *Great Plains Research* 6:41–60.
- Ermacoff, N. 1968. Marsh and habitat management practices at the Mendota Wildlife Area. [Place of publication unknown]: California Department of Fish and Game. Game Management Leaflet No. 12. 10 p.
- Evans, R.D.; Wolfe, C.W., Jr. 1967. Waterfowl production in the Rainwater Basin area of Nebraska. *Journal of Wildlife Management* 33(4):788–94.
- Farmer, A.H.; Parent, F. 1997. Effects of landscape on shorebird movements at spring migration stopovers. *Condor* 99: 698–707.
- Farmer, A.H.; Wiens, J. 1999. Models and reality: time-energy trade-offs in pectoral sandpiper migration. *Ecology* 80:2566–80.
- Farrar, J. 1982. The Rainwater Basin, Nebraska's vanishing wetlands. NEBRASKAland. March [updated and reprinted 1988 February]. Nebraska Game and Parks Commission. 15 p.
- . 1996a. Nebraska's Rainwater Basin. NEBRASKAland. March. Nebraska Game and Parks Commission. 18–35.
- . 1996b. The Troester tunnel: notes from the field. NEBRASKAland. May. Nebraska Game and Parks Commission. 6–7.

- Federal Geographic Data Committee. 1997. FGDC vegetation classification and information standards. Reston, VA: [Publisher name unknown]. [Pages unknown].
- Fleskes, J.P.; Jarvis, R.L.; Gilmer, D.S. 2002. September–March survival of female northern pintails radiotagged in San Joaquin Valley, California. *Journal of Wildlife Management* 31:212–9.
- Frankforter, J.D. 1996. Nebraska wetland resources. In: Fretwell, J.D.; Williams, J.S.; Redman, P.J.; editors. National water summary of wetland resources. [Place of publication unknown]: U.S. Department of the Interior, U.S. Geological Survey. U.S. Geological Survey Water Supply Paper 2425. 261–6.
- Fredrickson, L.H.; Reid, F.A. 1988. Waterfowl use of wetland complexes. [Place of publication unknown]: U.S. Department of the Interior, Fish and Wildlife Service. U.S. Fish and Wildlife Service Leaflet 13.2.1. [Pages unknown].
- Fredrickson, L.H.; Taylor, T.S. 1982. Management of seasonally flooded impoundments for wildlife. Washington DC: U.S. Department of the Interior, Fish and Wildlife Service. U.S. Fish and Wildlife Service Resource Publication 148. 29 p.
- Fritcher, S.C.; Rumble, M.A.; Flake, L.D. 2004. Grassland bird densities in several stages of mixed-grass prairie. *Journal of Range Management* 57:351–7.
- Fuhlendorf, S.D.; Engle, D.M. 2001. Restoring heterogeneity on rangelands: ecosystem management based on evolutionary grazing patterns. *Bioscience* 51: 625–32.
- . 2004. Application of the fire-grazing interaction to restore a shifting mosaic on tallgrass prairie. *Journal of Applied Ecology* 41:604–14.
- Gabbert, A.E.; Leif, A.P.; Purvis, J.R. [et al.] 1999. Survival and habitat use by ring-necked pheasants during two disparate winters in South Dakota. *Journal of Wildlife Management* 63:711–22.
- Gaiser, E.E.; Lang, K.L. 1998. Distribution of cladoceran zooplankton among prairie pothole wetlands in northwest Iowa. *Lake and Reservoir Management* 14(1):37–51.
- Gersib, R.A. 1991. Nebraska wetlands priority plan. [Place of publication unknown]: Nebraska Game and Parks Commission. [Pages unknown].
- Gersib, R.A.; Cornely, J.; Trout, A. [et al.] 1990. Concept plan for waterfowl habitat protection, Rainwater Basin area of Nebraska. Lincoln, NE: Nebraska Game and Parks Commission. 71 p.
- Gersib, R.A.; Dinan, K.F.; Kauffeld, J.D. [et al.] 1992. Looking to the future: an implementation plan for the Rainwater Basin Joint Venture. Lincoln, NE: Nebraska Game and Parks Commission. 56 p.
- Gersib, R.A.; Elder, B.; Dinan, K.F. [et al.] 1989a. Waterfowl values by wetland type within Rainwater Basin wetlands with special emphasis on activity time budget and census data. Lincoln, NE: Nebraska Game and Parks Commission; Grand Island, NE: U.S. Department of the Interior, Fish and Wildlife Service. 105 p.
- Gersib, R.A.; Raines, R.R.; Rosier, W.S. [et al.] 1989b. A functional assessment of selected wetlands within the Rainwater Basin area of Nebraska. Lincoln, NE: Nebraska Game and Parks Commission. 41 p.
- Gilbert, M.C. 1989. Ordination and mapping of wetland communities in Nebraska's Rainwater Basin region. CEMRO Environmental Report 89-1. Omaha, NE: U.S. Army Corps of Engineers, Omaha District. 105 p.
- Gleason, H.R.; Euliss, N.H., Jr.; Hubbard, D.E. [et al.] 2003. Effects of sediment load on emergence of aquatic invertebrates and plants from wetland soil egg and seed banks. *Wetlands* 23:26–34.
- Go, V.; Garey, J.; Wolff, M.S. [et al.] 1999. Estrogenic potential of certain pyrethroid compounds in the MCF-7 human breast carcinoma cell line. *Environmental Health Perspectives* 107(3):173–7.
- Gordon, C.C. 1989. The relationship of wetland characteristics to avian cholera (*Pasteurella multocida*) outbreaks in the Rainwater Basin area of Nebraska [master's thesis]. Brookings, SD: South Dakota State University. 126 p.
- Gordon C.C.; Flake, L.D.; Higgins, K.F. 1997. Trace metals in water and sediments of wetlands in the Rainwater Basin area of Nebraska. In: The Proceedings of the South Dakota Academy of Sciences; [Place and date of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 76:253–61.
- Grant, R.A. 1965. The burrowing owl in Minnesota. *The Loon* (3):2–17.
- Grossman, D.H.; Faber-Langendoen, D.; Weakley, A.W. [et al.] 1998. International classification of ecological communities: terrestrial vegetation of the United States. Vol. 1 of The National Vegetation Classification Standard. Arlington, VA: The Nature Conservancy. [Pages unknown].
- Haug, E.A.; Millsap, B.A.; Martell, M.S. 1993. Burrowing owl (*Speotyto cuniculari*) In: Poole, A.; Gill, F.; editors. The Birds of North America, No. 61 [doi:10.2173/bna.61]. Philadelphia: The Academy of Natural Sciences; Washington DC: The American Ornithologists' Union. [Pages unknown].
- Haukos, D.A.; Smith, L.M. 1993. Moist-soil management of playa lakes for migrating and wintering ducks. *Wildlife Society Bulletin* 21:288–98.
- Hayes T.B.; Haston, K.; Tsui, M. [et al.] 2002. Atrazine induced hermaphroditism at 0.1 ppb in American leopard frogs (*Rana pipiens*): laboratory and field evidence. *Environmental Health Perspectives* 111: 568–75.
- Heitmeyer, M.E. 1988. Body composition of female mallards in winter in relation to annual cycle events. *Condor* 90:669–80.

- Heitmeyer, M.E.; Fredrickson, L.H. 1981. Do wetland conditions in the Mississippi delta hardwoods influence mallard recruitment? In: Transactions, North American Wildlife and Natural Resources Conference; [Date and location of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 46:44–57.
- Helzer, C.J. 1996. The effects of wet meadow fragmentation on grassland birds [master's thesis]. Lincoln, NE: University of Nebraska–Lincoln. [Pages unknown].
- Helzer C.J.; Jelinski, D.E. 1999. The relative importance of patch area and perimeter-area ratio to grassland breeding birds. *Ecological Applications* 9(4):1448–58.
- Herkert, J.R.; Reinking, D.L.; Wiedenfeld, D.A. [et al.] 2003. Effects of prairie fragmentation on the nest success of breeding birds in the mid-continental United States. *Conservation Biology* 17(2):587–94.
- Herkert, J.R.; Szafoni, R.E.; Kleen, V.M. [et al.] 1993. Habitat establishment, enhancement and management for forest and grassland birds in Illinois. Springfield, IL: Illinois Department of Conservation, Division of Natural Heritage. Natural Heritage Technical Publication 1. [Pages unknown].
- Higgins, K.F. 1984. Lightning fires in grasslands in North Dakota and in pine savanna lands in nearby South Dakota and Montana. *Journal of Range Management* 37:100–3.
- . 1986. Interpretation and compendium of historic fire accounts in the northern Great Plains. Washington DC: U.S. Department of the Interior, Fish and Wildlife Service. Resource Publication 161. [Pages unknown].
- Hughes, J.P. 1996. The effect of vegetative structure and landscape composition on avian abundance and reproductive success in CRP fields in northeastern Kansas [master's thesis]. Manhattan, KS: Kansas State University. [Pages unknown].
- Jessen, R.L. 1970. Mallard population trends and hunting losses in Minnesota. *Journal of Wildlife Management* 34:93–105.
- Johnsgard, P.A. 1983. The grouse of the world [PhD dissertation]. Lincoln, NE: University of Nebraska–Lincoln. [Pages unknown].
- . 1997. The birds of Nebraska and adjacent plains states. Lincoln, NE: Nebraska Game and Parks Commission. Nebraska Ornithologist's Union Occasional Papers, No. 6. [Pages unknown].
- Johnson, R.G.; Temple, S.A. 1990. Nest predation and brood parasitism of tallgrass prairie birds. *Journal of Wildlife Management* 54:106–11.
- Jorde, D.G. 1981. Winter and spring staging ecology of mallards in south central Nebraska [master's thesis]. [City of publication unknown], ND: University of North Dakota. 116 p.
- Jorde, D.G.; Krapu, G.L.; Crawford, R.D. 1983. Feeding ecology of mallards wintering in Nebraska. *Journal of Wildlife Management* 47:1044–53.
- Jorde, D.G.; Krapu, G.L.; Crawford, D.C. [et al.] 1984. Effects of weather on habitat selection and behavior of mallards wintering in Nebraska. *The Condor* 86:258–65.
- Jorgensen, J.G. 2004. An overview of shorebird migration in the eastern Rainwater Basin, Nebraska. Lincoln, NE: Nebraska Game and Parks Commission. Nebraska Ornithologist's Union Occasional Papers, No. 8. [Pages unknown].
- Kaminski, R.M.; Gluesing, E.A. 1987. Density and habitat-related recruitment in mallards. *Journal of Wildlife Management* 51:141–8.
- Kantrud, H.A. 2006. Cattail management symposium—history of cattails on the prairies: wildlife impacts. <<http://www.npwrc.usgs.gov/resource/plants/cattail/kantrud.htm>> [Access date unknown].
- Keech, C.F.; Dreeszen, V.H. 1959. Geology and ground-water resources of Clay County, Nebraska. Washington DC: U.S. Department of the Interior, U.S. Geological Survey. U.S. Geological Survey Water Supply Paper 1468. 57 p.
- Kendeigh, S.C.; Dol'nik, V.R.; Gavrilov, V.M. 1977. Avian energetics. In: Pinowski, J.; Kendeigh, S.C.; editors. Granivorous birds in ecosystems. New York: Cambridge University Press. 127–204.
- Kennedy, T.A.; Naeem, S.; Howe, K.M. [et al.] 2002. Biodiversity as a barrier to ecological invasion. *Nature* 417:636–8.
- Kim, I.Y.; Shin, J.H.; Kim, H.S. [et al.] 2004. Assessing estrogenic activity of pyrethroid insecticides using in vitro combination assays. *The Journal of Reproduction and Development* 50(2):245–55.
- Koerner, J.W.; Bookout, T.A.; Denarik, K.E. 1974. Movements of Canada geese color-marked near southwestern Lake Erie. *Journal of Wildlife Management* 38:275–89.
- Krapu, G.L. 1981. The role of nutrient reserves in mallard reproduction. *Auk* 98:29–38.
- Krapu, G.L.; Brandt, D.A.; Cox, R.R., Jr. 2004. Less waste corn, more land in soybeans, and the switch to genetically modified crops: trends with important implications for wildlife management. *Wildlife Society Bulletin* 32:127–36.
- Krapu, G.L.; Reinecke, K.J.; Jorde, D.G. [et al.] 1995. Spring staging ecology of midcontinent greater white-fronted geese. *Journal of Wildlife Management* 59:736–46.
- Kuzila, M.S. 1984. Genesis and morphology of soils in and around large depressions in Clay County, Nebraska [PhD dissertation]. Lincoln, NE: University of Nebraska–Lincoln. [Pages unknown].
- . 1994. Inherited morphologies of two large basins in Clay County, Nebraska. *Great Plains Research* 4:51–63.
- Kuzila, M.S.; Lewis, D.T. 1993. Soils in rain basins of south central Nebraska—properties, genesis and classification. *Soil Science American Journal* 37:155–61.

- LaGrange, T.G. 1985. Habitat use and nutrient reserves dynamics of spring migratory mallards in central Iowa [master's thesis]. [City of publication unknown], IA: Iowa State University. 81 p.
- . 2005. A guide to Nebraska's wetlands and their conservation needs. Lincoln, NE: Nebraska Game and Parks Commission. 59 p.
- LaGrange, T.G.; Dinsmore, J.J. 1988. Nutrient reserve dynamics of female mallards during spring migration through central Iowa. In: *Waterfowl in Winter*. Minneapolis: University of Minnesota Press. 287–97.
- Larson, D.L.; McDonald, S.; Fivizzani, A.J. [et al.] 1998. Effects of the herbicide atrazine on *Ambystoma tigrinum* metamorphosis: duration, larval growth, and hormonal response. *Physiological Zoology* 71(6): 671–9.
- Laubhan, M.L.; Fredrickson, L.H. 1992. Estimating seed production of common plants in seasonally flooded wetlands. *Journal of Wildlife Management* 56(2):329–37.
- Laubhan, M.L.; Gleason, R.A.; Knutsen, G.A. [et al.] 2006. A preliminary biological assessment of Long Lake National Wildlife Refuge Complex, North Dakota. Washington DC: U.S. Department of the Interior, Fish and Wildlife Service. Biological Technical Publication, BTP R6006-2006. [Pages unknown].
- Laubhan, Murray K.; Koerner, T.M.; Sprenger, M.D. [et al.] 2005. A preliminary biological assessment of Lacreek National Wildlife Refuge, Martin, South Dakota, USA. Jamestown, ND: U.S. Department of the Interior, U.S. Geological Survey. [Pages unknown].
- Lavergne, S.; Molofsky, J. 2004. Reed canarygrass (*Phalaris arundinacea* L.) as a biological model in the study of plant invasions. *Critical Reviews in Plant Sciences* 23:415–29.
- Loesch, C.R.; Kaminski, R.M. 1989. Winter body-weight patterns of female mallards fed agricultural seeds. *Journal of Wildlife Management* 53:1081–7.
- Madden, E.M. 1996. Passerine communities and bird-habitat relationships on prescribe-burned, mixed-grass prairie in North Dakota [master's thesis]. Bozeman, MT: Montana State University. [Pages unknown].
- Madden, E.M.; Hansen, A.J.; Murphy, R.K. 1999. Influence of prescribed fire history on habitat and abundance of passerine birds in northern mixed-grass prairie. *Canadian Field-Naturalist* 113(4): 627–40.
- McCarthy, C.; Pella, T.; Link, G. [et al.] 1997. Greater prairie chicken nesting habitat, Sheyenne National Grassland, North Dakota. [Place of publication unknown]: U.S. Department of Agriculture, Forest Service. General Technical Report RM-GTR-298. [Pages unknown].
- McKee, G.M.; Ryan, R.; Mechlin, L.M. 1998. Predicting greater prairie-chicken nest success from vegetation and landscape characteristics. *Journal of Wildlife Management* 62:314–21.
- McMurtrey, M.D.; Craig, R.; Schildman, G. 1972. Nebraska wetland survey. [City of publication unknown], NE: Nebraska Game and Parks Commission. Habitat Work Plan K-71. 78 p.
- Messmer, T.A. 1990. Influence of grazing treatments on non-game birds and vegetation structure in south central North Dakota [PhD dissertation]. Fargo, ND: North Dakota State University. [Pages unknown].
- Merrill, M.D.; Chapman, K.A.; Poiani, K.A. [et al.] 1999. Land-use patterns surrounding greater prairie-chicken leks in northwestern Minnesota. *Journal of Wildlife Management* 63:189–98.
- Miller, M.R.; Reinecke, K.J. 1984. Proper expression of metabolizable energy in avian energetics. *Condor* 86(4):396–400.
- Mollhoff, W.J. 2001. The Nebraska breeding bird atlas, 1984–89. Lincoln, NE: Nebraska Game and Parks Commission. Nebraska Ornithologist's Union Occasional Papers, No. 7. 233 p.
- Moon, J.A.; Haukos, D.A. 2006. Survival of female northern pintails wintering in the Playa Lakes region of northwestern Texas. *Journal of Wildlife Management* 70:777–83.
- Moravec, J. 1993. Syntaxonomic and nomenclatural treatment of Scandinavian-type associations and sociations. *Journal of Vegetation Science* 4:833–8.
- [NASS] National Agricultural Statistics Service. 2004. Agricultural chemical usage 2003 field crops summary. <[http://www.nass.usda.gov/Statistics\\_by\\_Subject/Environmental/index.asp](http://www.nass.usda.gov/Statistics_by_Subject/Environmental/index.asp)> [Access date unknown].
- [NDEQ] Nebraska Department of Environmental Quality. 2006. Title 117, Nebraska surface water quality standards. In: *Water quality standard for wetlands*. [Place of publication unknown]. [Pages unknown]. Chapter 4.
- [NGPC] Nebraska Game and Parks Commission. 1977–99 unpublished reports. Lincoln, NE. [Pages unknown].
- . 1997–99 unpublished reports. Lincoln, NE. [Pages unknown].
- . 2000–03 unpublished reports. Lincoln, NE. [Pages unknown].
- Nelson K.J.; Esmoil, B.J. 1999. Background contaminants evaluation of Fort Niobrara and Valentine National Wildlife Refuge. Grand Island, NE: U.S. Department of the Interior, Fish and Wildlife Service. U.S. Fish and Wildlife Service Contaminant Report. [Pages unknown].
- Olson, R.A.; Flake, L.D. 1975. Nesting of ring-necked pheasants in eastern South Dakota. In: *The Proceedings of the South Dakota Academy of Science*; [Place and date of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 54:126–36.

- Pampush, G.J.; Anthony, R.G. 1993. Nest success, habitat utilization and nest-site selection of longbilled curlews in the Columbia Basin, Oregon. *Condor* 95:957–67.
- Pederson, R.L.; Jorde, D.G.; Simpson, S.G. 1989. Northern Great Plains. In: Smith, L.M.; Pederson, R.L.; Kaminski, R.M.; editors. *Habitat management for migrating and wintering waterfowl in North America*. Lubbock, TX: Texas Technological University Press. 281–310.
- Petrie, S.A.; Rogers, K.H. 2004. Nutrient-reserve dynamics of white-faced whistling-ducks breeding in a semiarid South Africa. *Canadian Journal of Zoology* 82:1082–90.
- Price, J.I.; Brand, C.J. 1984. Persistence of *Pasteurella multocida* in Nebraska wetlands under epizootic conditions. *Journal of Wildlife Diseases* 20(2):90–4.
- Prince, H.H. 1979. Bioenergetics of postbreeding dabbling ducks. In: Bookhout, T.A.; editor. *Waterfowl and wetlands—an integrated review. Proceedings of a symposium of the North Central Section of The Wildlife Society*; [Date and location of symposium unknown]. La Crosse, WI: La Crosse Printing. 103–17.
- Prose, B.L. 1987. *Habitat suitability index models: plains sharp-tailed grouse*. Washington DC: Natural Ecology Center. U.S. Fish and Wildlife Service Biological Report 82. [Pages unknown].
- Rabbe, M.; Drahota, J.; Bishop, A. 2004. Estimating duck use-days in the Rainwater Basin wetlands—a baseline model for management implications [unpublished report]. On file at Rainwater Basin Wetland Management District, Kearney, NE. [Pages unknown].
- Raines, R.R.; Gilbert, M.C.; Gersib, R.A. [et al.] 1990. Regulatory planning for Nebraska's Rainwater Basin wetlands (advanced identification of disposal areas). Prepared for the Rainwater Basin advanced identification study. Kansas City, KS: U.S. Environmental Protection Agency, region 7; Omaha, NE: U.S. Army Engineer District. 46 p.
- [RWB JV] Rainwater Basin Joint Venture. 1993. *Water management options for wetland development in the Rainwater Basin*. [Place of publication unknown]: [Publisher name unknown]. 16 p.
- . 1994. *Best management practices for Rainwater Basin wetlands*. [Place of publication unknown]: [Publisher name unknown]. Public Lands Work Group. 41 p.
- Raveling, D.G. 1978. Dynamics of distribution of Canada geese in winter. In: *Transactions, North American Wildlife and Natural Resources Conference*; [Date and place of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 42:206–25.
- Raveling, D.G.; Crews, W.E.; Klimstra, W.D. 1972. Activity patterns of Canada geese during winter. *Wilson Bulletin* 84:278–95.
- Raveling, D.G.; Heitmeyer, M.E. 1989. Relationships of population size and recruitment of pintails to habitat conditions and harvest. *Journal of Wildlife Management* 53:1088–103.
- Redmond, R.L.; Bicak, T.K.; Jenni, D.A. 1981. An evaluation of breeding season census techniques for long-billed curlews (*Numenius americanus*). *Studies in Avian Biology* 6:197–201.
- Reid, F.R.; Kelley, J.R., Jr.; Taylor, T.S. [et al.] 1989. Upper Mississippi Valley wetlands: refuges and moist-soil impoundments. In: Smith, L.M.; Pederson, R.L.; Kaminski, R.M.; editors. *Habitat management for migrating and wintering waterfowl in North America*. Lubbock, TX: Texas Technological University Press. 181–202.
- Reinecke, K.J.; Kaminski, R.M.; Moorehead, D.J. [et al.] 1989. Mississippi alluvial valley. In: Smith, L.M.; Pederson, R.L.; Kaminski, R.M.; editors. *Habitat management for migrating and wintering waterfowl in North America*. Lubbock, TX: Texas Technological University Press. 203–47.
- Renken, R.B.; Dinsmore, J.J. 1987. Nongame bird communities on managed grasslands in North Dakota. *Canadian Field-Naturalist* 101(4):551–57.
- Richert, A. 1999. *Multiple scale analyses of whooping crane habitat in Nebraska* [master's thesis]. Lincoln, NE: University of Nebraska–Lincoln. [Pages unknown].
- Rohwer, F.C. 1984. Patterns of egg laying in prairie ducks. *Auk* 101:603–5.
- Samuel, M.D. 1995. *Physical and biological factors associated with avian cholera in wetland ecosystems* [program report]. Madison, WI: National Wildlife Health Center. [Pages unknown].
- Samuel, M.D.; Shadduck, D.J.; Goldberg, D.R. [et al.] 2005. Avian cholera in waterfowl: the role of lesser snow and Ross's geese as disease carriers in the Playa Lakes region. *Journal of Wildlife Diseases* 41(1):48–57.
- Schildman, G.; Hurt, J. 1984. Update of Rainwater Basin wetland survey. Survey of habitat work plan K-83. W-15-R-40. [Place of publication unknown]: Nebraska Game and Parks Commission. 13 p.
- Schneider, R.; Humpert, M.; Stoner, K. [et al.] 2005. *The Nebraska Natural Legacy Project: A comprehensive wildlife conservation strategy*. Lincoln, NE: Nebraska Game and Parks Commission. 245 p.
- Schwarz, M.S.; Echols, K.R.; Wolcott, M.J. [et al.] 2004. Environmental contaminants associated with swine concentrated animal feeding operation and implications for McMurtrey National Wildlife Refuge. Grand Island, NE: U.S. Department of the Interior, Fish and Wildlife Service. U.S. Fish and Wildlife Service Contaminant Report. [Pages unknown].
- Sharpe, R.S.; Silcock, W.R.; Jorgensen, J.G. 2001. *Birds of Nebraska: their distribution and temporal occurrence*. Lincoln, NE: University of Nebraska Press. 520 p.

- Sharpley, A.N.; Daniel, T.; Sims, T. [et al.] 1999. Agricultural phosphorus and eutrophication. Washington DC: U.S. Department of Agriculture, Agricultural Research Service. ARS-149. [Pages unknown].
- Sherfy, M.H.; Kirkpatrick, R.L. 1998. Nutritional value and management of waterfowl and shorebird foods in Atlantic coastal moist-soil impoundments [master's thesis]. <<http://scholar.lib.vt.edu/theses/available/etd-033199-130907/unrestricted/chapter1.pdf>> [Access date unknown]. [Pages unknown].
- Singleton, J.R. 1951. Production and utilization of waterfowl food plants on the east Texas Gulf Coast. *Journal of Wildlife Management* 15(1):46–56.
- Smith, B.J.; Higgins, K.F. 1990. Avian cholera and temporal changes in wetland numbers and densities in Nebraska's Rainwater Basin area. *Wetlands* 10:1–5.
- Smith, B.J.; Higgins, K.F.; Gritzner, C.F. 1989. Land use relationships to avian cholera outbreaks in the Nebraska Rainwater Basin area. *Prairie Naturalist* 21(3):125–36.
- Smith, B.J.; Higgins, K.F.; Tucker, W.L. 1990. Precipitation, waterfowl densities and mycotoxins: their potential effect on avian cholera epizootics in the Nebraska Rainwater Basin area. In: *Transactions of North American Wildlife and Natural Resources Conference*; [Date and place of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 55:269–82.
- Smith, L.M. 1989. Effects of grazing and burning on nutritive quality of cattail in playas. *Journal of Aquatic Plant Management* 27:51–3.
- . 1990. Waterfowl habitat management and research in North America. In: *Transactions of the 19th International Union of Game Biologist Congress*; [Date of congress unknown]; Trondheim, Norway. [Place of publication unknown]: [Publisher name unknown]. 2:468–76.
- . 1998. Research needs for the Rainwater Basin of Nebraska: a hierarchical approach. Lincoln, NE: Nebraska Game and Parks Commission. 22 p.
- Snyder, W.D. 1984. Ring-necked pheasant nesting ecology and wheat farming on the high plains. *Journal of Wildlife Management* 48:878–88.
- Solberg, J.W. 1992. Coordinated mid-continent white-fronted goose survey. [Place of publication unknown]: U.S. Department of the Interior, Fish and Wildlife Service. U.S. Fish and Wildlife Service annual report. 6 p.
- Sporrong, J.M. 2001. Response of nongame birds and terrestrial invertebrates to restoration of upland grasslands in the Rainwater Basin region, Nebraska [master's thesis]. Eau Claire, WI: University of Wisconsin–Eau Claire. 108 p.
- Starks, P.J. 1984. Analysis of rainbasin depressions of Clay County, Nebraska [master's thesis]. Omaha, NE: University of Nebraska. 90 p.
- Steinauer, G.; Rolfsmeier, S. 2003. Terrestrial natural communities of Nebraska. [Place of publication unknown]: Nebraska Game and Parks Commission. 143 p.
- Stutheit, R.G.; Gilbert, M.C.; Whited, P.M. [et al.]; editors. 2004. A regional guidebook for applying the hydrogeomorphic approach to assessing wetland functions of Rainwater Basin depressional wetlands in Nebraska. ERDC/EL TR-04-4. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 118 p.
- Taylor, J.P.; Smith, L.M. 2005. Migratory bird use of belowground foods in moist-soil managed wetlands in the Middle Rio Grande Valley, New Mexico. *Wildlife Society Bulletin* 33(2):574–82.
- Taylor, M.W.; Wolfe, C.W.; Baxter, W.L. 1978. Land use change and ring-necked pheasants in Nebraska. *Wildlife Society Bulletin* 6:226–30.
- Trautman, C.G.; Dahlgren, R.B.; Seubert, J.L. 1959. Pheasant nesting. *South Dakota Conservation Digest* 26:18–21.
- Traylor, Susan S. 2000. Waterfowl migration in south central Nebraska, February–March 2000 [unpublished report]. Lincoln, NE: University of Nebraska; Nebraska Game and Parks Commission. [Pages unknown].
- [UNL] University of Nebraska–Lincoln. 2004. Biology and treatment recommendations for field crop pests. Lincoln, NE: Department of Entomology. <<http://entomology.unl.edu/fldcrops/pestipm.htm#Item2>> [Access date unknown].
- . 2006a. Groundwater map—drought 2000–2006. Lincoln, NE: Institute of Agriculture and Natural Resources, School of Natural Resources, Conservation and Survey Division. <<http://csd.unl.edu/csd-esic/GWMapArchives/2006GWMaps/drought2000to2006-300dpi.tif>> accessed 25 June 2007.
- . 2006b. Groundwater map—predevelopment to spring 2006. Lincoln, NE: Institute of Agriculture and Natural Resources, School of Natural Resources, Conservation and Survey Division. <<http://csd.unl.edu/csd-esic/GWMapArchives/2006GWMaps/predev06-300dpi.tif>> accessed 25 June 2007.
- U.S. Census Bureau. 2000–2007. State and county quickfacts: Nebraska. <<http://quickfacts.census.gov/qfd/states/31000.html>> [Access date unknown].
- (USDOE) U.S. Department of Energy. 1999. Carbon sequestration research and development. Report No. DOE/SC/FE-1. U.S. Department of Energy, Office of Science, Office of Fossil Energy. <[www.osti.gov/bridge/servlets/purl/810722-9s7bTP/native/810722.pdf](http://www.osti.gov/bridge/servlets/purl/810722-9s7bTP/native/810722.pdf)>. 289 p.
- [USFWS] U.S. Department of the Interior, Fish and Wildlife Service. 1954. Wetlands inventory of Nebraska. U.S. Fish and Wildlife Service, Billings, MT. 31 p.
- . 1977–99 unpublished reports. Grand Island, NE. [Pages unknown].

- . 1981. The Platte River ecology study special research report. Jamestown, ND: Northern Prairie Wildlife Research Center Online. <<http://www.npwrc.usgs.gov/resource/habitat/plriveco/index.htm>> (Version16JUL97) [Access date unknown].
- . 1983–93. Unpublished annual reports. Rainwater Basin Wetland Management District, Kearney, NE. [Pages unknown].
- . 1995. updated 2005. Migratory nongame birds of management concern in the United States: the 1995 list. Washington DC: Office of Migratory Bird Management. 22 p.
- . 2000. [No title]. <[http://mountain-prairie.fws.gov/reference/briefing\\_book\\_nd\\_2000.pdf](http://mountain-prairie.fws.gov/reference/briefing_book_nd_2000.pdf)> [Access date unknown].
- . 2001. The biological integrity, diversity, and environmental health policy. <<http://policy.fws.gov/library01fr3809.pdf>> [Access date unknown].
- . 2006. Livestock confinement [Rainwater Basin GIS layer]. Data on file at Rainwater Basin Wetland Management District.
- [USFWS et al.] U.S. Department of the Interior, Fish and Wildlife Service; Canadian Wildlife Service; Mexican Ministry of Environment, Natural Resources, and Fisheries. 1998. North American waterfowl management plan, 1998 update—expanding the vision. Washington DC: [Publisher name unknown]. 32 p.
- [USFWS and NGPC] U.S. Department of the Interior, Fish and Wildlife Service; Nebraska Game and Parks Commission. 1986. Rainwater Basin of Nebraska migratory bird habitat acquisition plan [unpublished plan]. On file at Rainwater Basin Wetland Management District. 30 p.
- [USEPA] U.S. Environmental Protection Agency. 2003. National pollutant discharge elimination system permit regulation and effluent limitation guidelines and standards for concentrated animal feeding operations (CAFOs). Federal Register 68(29):7177–269. <<http://cfpub.epa.gov/npdes/af/cafofinalrule.cfm>> [Access date unknown].
- van der Burg, Max Post. 2005. Factors affecting songbird nest survival and brood parasitism in the Rainwater Basin region of Nebraska [master's thesis]. Lincoln, NE: University of Nebraska–Lincoln. 71 p.
- Volkert, W.K. 1992. Response of grassland birds to a large-scale prairie planting project. The Passenger Pigeon 54(3):191–6.
- Warner, K.D.; Nieman, D.J. 1999. [Unpublished preliminary report]. Saskatoon, SK: Canadian Wildlife Service. 4 p.
- White, D.H.; James, D. 1978. Differential use of fresh water environments by wintering waterfowl of coastal Texas. Wilson Bulletin 90:99–111.
- Wiens, J.A. 1973. Pattern and process in grassland bird communities. Ecological Monographs 43:236–70.
- Windingstad, R.M.; Hurt, J.J.; Trout, A.K. [et al.] 1984. Avian cholera in Nebraska's Rainwater Basin. In: Transactions of North American Wildlife and Natural Resources Conference; [Date and location of conference unknown]. [Place of publication unknown]: [Publisher name unknown]. 49:577–83.
- Wood, W.W. 2000. Ground-water recharge in the southern high plains of Texas and New Mexico. [Place of publication unknown]: [Publisher name unknown]. U.S. Geological Survey Publication FS-127-99. 4 p.
- Zinkl, J.G.; Dey, N.; Hyland, J.M. [et al.] 1997. An epornitic of avian cholera in waterfowl and common crows in Phelps County, Nebraska, in the spring, 1975. Journal of Wildlife Diseases 13:194–8.

